







5.	The value of $\cos\left(\frac{\pi}{6} + \cot^{-1}(-\sqrt{3})\right)$ is  (A) -1 (B) $\frac{-\sqrt{3}}{2}$ (C) 0 (D) 1	
Ans	(A) -1	1
6.	If p and q are respectively the order and degree of the differential equation $\frac{d}{dx}\left(\frac{dy}{dx}\right)^3 = 0$ , then (p - q) is  (A) 0 (B) 1 (C) 2 (D) 3	
Ans	(B) 1	1
7.	The function $f(x) = x^2 - 4x + 6$ is increasing in the interval  (A) (0, 2) (B) $(-\infty, 2]$ (C) [1, 2] (D) $[2, \infty)$	
Ans	(D) $[2, \infty)$	1
8.	The line $x = 1 + 5\mu$ , $y = -5 + \mu$ , $z = -6 - 3\mu$ passes through which of the following point ?  (A) (1, -5, 6) (B) (1, 5, 6) (C) (1, -5, -6) (D) (-1, -5, 6)	
Ans	(C) (1, -5, -6)	1
9.		

	<p>The area of the shaded region (figure) represented by the curves <math>y = x^2</math>, <math>0 \leq x \leq 2</math> and <math>y</math>-axis is given by</p>  <p>(A) <math>\int_0^2 x^2 dx</math>                      (B) <math>\int_0^2 \sqrt{y} dy</math></p> <p>(C) <math>\int_0^4 x^2 dx</math>                      (D) <math>\int_0^4 \sqrt{y} dy</math></p>													
Ans	(D) $\int_0^4 \sqrt{y} dy$	1												
10.	<p>If E and F are two events such that <math>P(E) &gt; 0</math> and <math>P(F) \neq 1</math>, then <math>P(\overline{E}/\overline{F})</math> is</p> <p>(A) <math>\frac{P(\overline{E})}{P(\overline{F})}</math>                      (B) <math>1 - P(\overline{E}/F)</math></p> <p>(C) <math>1 - P(E/F)</math>                      (D) <math>\frac{1 - P(E \cup F)}{P(\overline{F})}</math></p>													
Ans	(D) $\frac{1 - P(E \cup F)}{P(\overline{F})}$	1												
11.	<p>The probability distribution of a random variable X is given by :</p> <table border="1" data-bbox="220 1422 1029 1534"> <tbody> <tr> <td>X</td> <td>-4</td> <td>-3</td> <td>-2</td> <td>-1</td> <td>0</td> </tr> <tr> <td>P(X)</td> <td>0.1</td> <td>0.2</td> <td>0.3</td> <td>0.2</td> <td>0.2</td> </tr> </tbody> </table> <p>Then E(X) of distribution is</p> <p>(A) -1.8                      (B) -1</p> <p>(C) 1                      (D) 1.8</p>	X	-4	-3	-2	-1	0	P(X)	0.1	0.2	0.3	0.2	0.2	
X	-4	-3	-2	-1	0									
P(X)	0.1	0.2	0.3	0.2	0.2									
Ans	(A) -1.8	1												
12.	<p>If projection of <math>\vec{a} = \alpha \hat{i} + \hat{j} + 4\hat{k}</math> on <math>\vec{b} = 2\hat{i} + 6\hat{j} + 3\hat{k}</math> is 4 units, then <math>\alpha</math> is</p> <p>(A) -13                      (B) -5</p> <p>(C) 13                      (D) 5</p>													

Ans	(D) 5	1
13.	<p>The equation of a line parallel to the vector <math>3\hat{i} + \hat{j} + 2\hat{k}</math> and passing through the point (4, -3, 7) is :</p> <p>(A) <math>x = 4t + 3, y = -3t + 1, z = 7t + 2</math></p> <p>(B) <math>x = 3t + 4, y = t + 3, z = 2t + 7</math></p> <p>(C) <math>x = 3t + 4, y = t - 3, z = 2t + 7</math></p> <p>(D) <math>x = 3t + 4, y = -t + 3, z = 2t + 7</math></p>	
Ans	(C) $x = 3t + 4, y = t - 3, z = 2t + 7$	1
14.	<p>If a line makes angles of <math>\frac{3\pi}{4}, \frac{\pi}{3}</math> and <math>\theta</math> with the positive directions of <math>x, y</math> and <math>z</math>-axis respectively, then <math>\theta</math> is</p> <p>(A) <math>\frac{-\pi}{3}</math> only</p> <p>(B) <math>\frac{\pi}{3}</math> only</p> <p>(C) <math>\frac{\pi}{6}</math></p> <p>(D) <math>\pm \frac{\pi}{3}</math></p>	
Ans	No option is correct. Full marks may be awarded for attempting the question.	1
15.	<p>A factory produces two products X and Y. The profit earned by selling X and Y is represented by the objective function <math>Z = 5x + 7y</math>, where <math>x</math> and <math>y</math> are the number of units of X and Y respectively sold. Which of the following statement is correct ?</p> <p>(A) The objective function maximizes the difference of the profit earned from products X and Y.</p> <p>(B) The objective function measures the total production of products X and Y.</p> <p>(C) The objective function maximizes the combined profit earned from selling X and Y.</p> <p>(D) The objective function ensures the company produces more of product X than product Y.</p>	
Ans	(C) The objective function maximizes the combined profit earned from selling X and Y	1
16.		

	<p>If A denotes the set of continuous functions and B denotes set of differentiable functions, then which of the following depicts the correct relation between set A and B ?</p> <p>(A)  (B) </p> <p>(C)  (D) </p>	
Ans	<p>(B) </p>	1
17.	<p>Four friends Abhay, Bina, Chhaya and Devesh were asked to simplify <math>4AB + 3(AB + BA) - 4BA</math>, where A and B are both matrices of order <math>2 \times 2</math>. It is known that <math>A \neq B \neq I</math> and <math>A^{-1} \neq B</math>.</p> <p>Their answers are given as :</p> <p>Abhay : <math>6AB</math>  Bina : <math>7AB - BA</math>  Chhaya : <math>8AB</math>  Devesh : <math>7BA - AB</math></p> <p>Who answered it correctly ?</p> <p>(A) Abhay (B) Bina  (C) Chhaya (D) Devesh</p>	
Ans	(B) Bina	1
18.	<p>If A and B are square matrices of order m such that <math>A^2 - B^2 = (A - B)(A + B)</math>, then which of the following is always correct ?</p> <p>(A) <math>A = B</math> (B) <math>AB = BA</math>  (C) <math>A = 0</math> or <math>B = 0</math> (D) <math>A = I</math> or <math>B = I</math></p>	
Ans	(B) $AB = BA$	1







	<p>Then <math>3\cos^2\alpha = 1</math></p> $\Rightarrow \cos\alpha = \frac{1}{\sqrt{3}}$ <p>The unit vector along the vector <math>\vec{a} = \frac{1}{\sqrt{3}}(\hat{i} + \hat{j} + \hat{k})</math></p> $\vec{a} = 5(\hat{i} + \hat{j} + \hat{k})$ <p style="text-align: center;">OR</p>	<p>1</p> <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p>
25 (b) Ans	<p><math>R(\vec{x}) \quad P(\vec{\alpha}) \quad Q(\vec{\beta})</math></p> $\frac{QR}{QP} = \frac{3}{2}$ <p>Hence, R divides PQ, externally, in the ratio 1:3.</p> <p>The Position vector of R = <math>\vec{x} = \frac{\vec{\beta} - 3\vec{\alpha}}{1 - 3} = \frac{3\vec{\alpha} - \vec{\beta}}{2}</math></p>	<p>1</p> <p>1</p>
SECTION-C		
This section comprises 6 Short Answer (SA) type questions of 3 marks each.		
26	<p>(a) If <math>y = \log\left(\sqrt{x} + \frac{1}{\sqrt{x}}\right)^2</math>, then show that <math>x(x+1)^2 y_2 + (x+1)^2 y_1 = 2</math>.</p> <p style="text-align: center;">OR</p> <p>(b) If <math>x\sqrt{1+y} + y\sqrt{1+x} = 0, -1 &lt; x &lt; 1, x \neq y</math>, then prove that <math>\frac{dy}{dx} = \frac{-1}{(1+x)^2}</math>.</p>	
26(a) Ans	<p>The given function can be written as</p> $y = 2 \log(x+1) - \log x$ $\Rightarrow y_1 = \frac{2}{x+1} - \frac{1}{x} = \frac{x-1}{x(x+1)}$ $\Rightarrow (x+1)y_1 = \frac{x-1}{x} = 1 - \frac{1}{x}$ $\Rightarrow (x+1)y_2 + y_1 = \frac{1}{x^2}$ $\Rightarrow x(x+1)^2 y_2 + x(x+1)y_1 = 1 + \frac{1}{x}$ $\Rightarrow x(x+1)^2 y_2 + x(x+1)y_1 = 1 + 1 - (x+1)y_1$	<p>1</p> <p>1</p>

	$\Rightarrow x(x+1)^2y_2 + (x+1)^2y_1 = 2$ <p style="text-align: center;">OR</p>	1
26(b) Ans	$x\sqrt{1+y} + y\sqrt{1+x} = 0$ $\Rightarrow x\sqrt{1+y} = -y\sqrt{1+x}$ $\Rightarrow x^2(1+y) = y^2(1+x)$ $\Rightarrow (x-y)(x+y) + xy(x-y) = 0$ $\Rightarrow (x-y)(x+y+xy) = 0$ $x \neq y \Rightarrow x+y+xy = 0$ $\Rightarrow y = \frac{-x}{1+x}$ $\Rightarrow \frac{dy}{dx} = \frac{-1}{(x+1)^2}$	          1/2          1          1/2          1
27.	Let R be a relation on set of real numbers $\mathbb{R}$ defined as $\{(x, y) : x - y + \sqrt{3}$ is an irrational number, $x, y \in \mathbb{R}\}$ Verify R for reflexivity, symmetry and transitivity.	
Ans	Let $x \in \mathbb{R}$ . Then we know that $x - x + \sqrt{3} = \sqrt{3}$ , which is an irrational number. $\Rightarrow (x, x) \in R$ Hence, R is reflexive. We have $\sqrt{3}, 2 \in \mathbb{R}$ such that $\sqrt{3} - 2 + \sqrt{3} = 2(\sqrt{3} - 1)$ , which is an irrational number $\Rightarrow (\sqrt{3}, 2) \in R$ . But, $2 - \sqrt{3} + \sqrt{3} = 2$ , which is a rational number. Hence, $\Rightarrow (2, \sqrt{3}) \notin R$ . . Therefore, R is not symmetric. Let $-\sqrt{3}, \sqrt{3}, 2 \in \mathbb{R}$ such that $(-\sqrt{3}, \sqrt{3}), (\sqrt{3}, 2) \in R$ . But, $(-\sqrt{3}, 2) \notin R$ Therefore, R is not transitive.	          1          1          1

28.

Solve the following linear programming problem graphically :

Minimise  $Z = 2x + y$

subject to the constraints :

$$3x + y \geq 9$$

$$x + y \geq 7$$

$$x + 2y \geq 8$$

$$x, y \geq 0$$

Ans



Corner point	Value of $Z = 2x + y$
A(0, 9)	9
B(1, 6)	8
C(6, 1)	13
D(8, 0)	16

In the half-plane  $2x + y < 8$ , there is no point in common with the feasible region.

Hence, the minimum value of  $Z$  is 8, which is attained at  $x = 1, y = 6$ .

Correct graph

And shading

1½


1


½

29	<p>(a) A die with number 1 to 6 is biased such that <math>P(2) = \frac{3}{10}</math> and probability other numbers is equal. Find the mean of the number of times number appears on the dice, if the dice is thrown twice.</p> <p style="text-align: center;"><b>OR</b></p> <p>(b) Two dice are thrown. Defined are the following two events A and B :  <math>A = \{(x, y) : x + y = 9\}</math>, <math>B = \{(x, y) : x \neq 3\}</math>, where <math>(x, y)</math> denote a point in the sample space.  Check if events A and B are independent or mutually exclusive.</p>													
29(a) Ans	<p><math>P(2) = \frac{3}{10}</math>, <math>P(\text{any other number}) = 1 - \frac{3}{10} = \frac{7}{10}</math></p> <p>Let X represent the Random Variable “the number of 2’s”.</p> <p>Then <math>X = 0, 1, 2</math></p> <p>The probability distribution is</p> <table border="1" data-bbox="212 902 1131 1164"> <thead> <tr> <th>X</th> <th>P(X)</th> <th>XP(X)</th> </tr> </thead> <tbody> <tr> <td>0</td> <td><math>\frac{7}{10} \times \frac{7}{10} = \frac{49}{100}</math></td> <td>0</td> </tr> <tr> <td>1</td> <td><math>\frac{3}{10} \times \frac{7}{10} \times 2 = \frac{42}{100}</math></td> <td><math>\frac{42}{100}</math></td> </tr> <tr> <td>2</td> <td><math>\frac{3}{10} \times \frac{3}{10} = \frac{9}{100}</math></td> <td><math>\frac{18}{100}</math></td> </tr> </tbody> </table> <p>Mean = <math>\sum XP(X) = \frac{60}{100} = 0.6</math></p> <p style="text-align: center;"><b>OR</b></p>	X	P(X)	XP(X)	0	$\frac{7}{10} \times \frac{7}{10} = \frac{49}{100}$	0	1	$\frac{3}{10} \times \frac{7}{10} \times 2 = \frac{42}{100}$	$\frac{42}{100}$	2	$\frac{3}{10} \times \frac{3}{10} = \frac{9}{100}$	$\frac{18}{100}$	<p>½</p> <p>½</p> <p>1½</p> <p>½</p>
X	P(X)	XP(X)												
0	$\frac{7}{10} \times \frac{7}{10} = \frac{49}{100}$	0												
1	$\frac{3}{10} \times \frac{7}{10} \times 2 = \frac{42}{100}$	$\frac{42}{100}$												
2	$\frac{3}{10} \times \frac{3}{10} = \frac{9}{100}$	$\frac{18}{100}$												
29(b) Ans	<p><math>A = \{(3,6), (4,5), (5,4), (6,3)\}</math></p> <p><math>P(A) = \frac{4}{36} = \frac{1}{9}</math>, <math>P(B) = \frac{30}{36} = \frac{5}{6}</math></p> <p><math>P(A \cap B) = \frac{3}{36} = \frac{1}{12}</math></p> <p><math>P(A) \times P(B) = \frac{5}{54} \neq P(A \cap B)</math></p> <p>Therefore, A and B are not independent.</p> <p>A and B are not mutually exclusive as <math>A \cap B \neq \emptyset</math></p>	<p>1</p> <p>½</p> <p>1</p> <p>½</p>												

30	<p>(a) Solve the differential equation <math>2(y + 3) - xy \frac{dy}{dx} = 0</math>; given <math>y(1) = -2</math>.</p> <p style="text-align: center;"><b>OR</b></p> <p>(b) Solve the following differential equation :</p> $(1 + x^2) \frac{dy}{dx} + 2xy = 4x^2.$	
30(a) Ans	<p>Given differential equation can be written as</p> $\frac{y}{y+3} dy = \frac{2}{x} dx$ $\Rightarrow \int \left(1 - \frac{3}{y+3}\right) dy = 2 \int \frac{1}{x} dx$ $\Rightarrow y - 3 \log y + 3  = 2 \log x  + C$ <p><math>y = -2</math>, when <math>x = 1 \Rightarrow C = -2</math></p> <p>Hence, the required particular solution is</p> $\Rightarrow y - 3 \log y + 3  = 2 \log x  - 2$ <p style="text-align: center;"><b>OR</b></p>	1 1½ ½
30(b) Ans	<p>Given differential equation can be written as</p> $\frac{dy}{dx} + \frac{2x}{1+x^2} y = \frac{4x^2}{1+x^2}, \text{ which is linear in } y.$ $\text{I.F.} = e^{\int \frac{2x}{1+x^2} dx} = e^{\log(1+x^2)} = 1 + x^2$ <p>The solution is given by</p> $y(1 + x^2) = \int 4x^2 dx$ $\Rightarrow y(1 + x^2) = \frac{4}{3} x^3 + C$ <p>or <math>y = \frac{4x^3}{3(1+x^2)} + \frac{C}{1+x^2}</math>, which is the required general solution</p>	1 1 1
31.	<p>If <math>\int_a^b x^3 dx = 0</math> and <math>\int_a^b x^2 dx = \frac{2}{3}</math>, then find the values of a and b.</p>	
Ans		



	<p>S.D. = <math>\frac{ (\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) }{ \vec{b}_1 \times \vec{b}_2 } = \frac{12}{\sqrt{3}} = 4\sqrt{3}</math></p> <p style="text-align: center;">OR</p>	1
<p>32(b) Ans</p>	<div style="text-align: center;">  </div> <p>Let the image of A in the line be <math>A'(\alpha, \beta, \gamma)</math></p> <p>The point P, which is the point of intersection of the lines <math>l</math> and <math>AA'</math>, will have coordinates <math>(\lambda + 4, -\lambda + 2, -\lambda + 2)</math> for some <math>\lambda</math>.</p> <p>Drs of AP are <math>\langle \lambda + 2, -\lambda + 1, -\lambda \rangle</math></p> <p><math>AP \perp l</math></p> $(\lambda + 2) - (-\lambda + 1) - (-\lambda) = 0$ $\Rightarrow \lambda = -\frac{1}{3}$ <p>Therefore, the coordinates of P are <math>(\frac{11}{3}, \frac{7}{3}, \frac{7}{3})</math></p> <p>P is the mid-point of <math>AA'</math></p> $\Rightarrow \frac{2 + \alpha}{2} = \frac{11}{3}, \frac{1 + \beta}{2} = \frac{7}{3}, \frac{2 + \gamma}{2} = \frac{7}{3}$ $\Rightarrow \alpha = \frac{16}{3}, \beta = \frac{11}{3}, \gamma = \frac{8}{3}$ <p>The coordinates of the image are <math>(\frac{16}{3}, \frac{11}{3}, \frac{8}{3})</math></p> <p>The equation of <math>AA'</math> is</p> $\frac{x - 2}{\frac{10}{3}} = \frac{y - 1}{\frac{8}{3}} = \frac{z - 2}{\frac{2}{3}}$ <p>or,</p>	<p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> <p>1</p> <p><math>\frac{1}{2}</math></p> <p><math>1\frac{1}{2}</math></p> <p>1</p>

	$\frac{3(x-2)}{5} = \frac{3(y-1)}{4} = \frac{3(z-2)}{1}$	
33.	Find : $\int (\sqrt{\tan x} + \sqrt{\cot x}) dx.$	
Ans	$I = \int \frac{\sin x + \cos x}{\sqrt{\sin x \cos x}} dx$ $= \sqrt{2} \int \frac{\sin x + \cos x}{\sqrt{\sin 2x}} dx$ <p>Put <math>\sin x - \cos x = t \Rightarrow (\cos x + \sin x)dx = dt</math></p> <p>On squaring both sides, we get <math>1 - \sin 2x = t^2</math></p> $I = \sqrt{2} \int \frac{1}{\sqrt{1-t^2}} dt$ $= \sqrt{2} \sin^{-1} t + C$ $= \sqrt{2} \sin^{-1} (\sin x - \cos x) + C$	<p>2</p> <p>1½</p> <p>1½</p>
34.	Using integration, find the area of the region bounded by the line $y = 5x + 2$ , the $x$ -axis and the ordinates $x = -2$ and $x = 2$ .	
Ans	 <p>The required area</p>	<p>Correct Sketch and shading</p> <p>2</p>





**SECTION-E**

This section comprises 3 case study based questions of 4 marks each

36.

A school is organizing a debate competition with participants as speakers  $S = \{S_1, S_2, S_3, S_4\}$  and these are judged by judges  $J = \{J_1, J_2, J_3\}$ . Each speaker can be assigned one judge. Let  $R$  be a relation from set  $S$  to  $J$  defined as  $R = \{(x, y) : \text{speaker } x \text{ is judged by judge } y, x \in S, y \in J\}$ .



Based on the above, answer the following :

(i) How many relations can be there from  $S$  to  $J$ ? **1**

(ii) A student identifies a function from  $S$  to  $J$  as  $f = \{(S_1, J_1), (S_2, J_2), (S_3, J_2), (S_4, J_3)\}$  Check if it is bijective. **1**

(iii) (a) How many one-one functions can be there from set  $S$  to set  $J$ ? **2**

**OR**

(iii) (b) Another student considers a relation  $R_1 = \{(S_1, S_2), (S_2, S_4)\}$  in set  $S$ . Write minimum ordered pairs to be included in  $R_1$  so that  $R_1$  is reflexive but not symmetric. **2**

36 Ans  
(i)


The number of relations =  $2^{4 \times 3} = 2^{12}$

**1**

36 Ans  
(ii)

Since,  $S_2$  and  $S_3$  have been assigned the same judge  $J_2$ , the function is not one-one.  
Hence, it is not bijective.

**1**

36 (iii) (a)	There cannot exist any one-one function from S to J as $n(S) > n(J)$ . Hence, the number of one-one functions from S to J is 0.  OR	2
36 (iii) (b)	To make $R_1$ reflexive and not symmetric we need to add the following ordered pairs:  $(S_1, S_1), (S_2, S_2), (S_3, S_3), (S_4, S_4)$	2
37.	<p>Three persons viz. Amber, Bonzi and Comet are manufacturing cars which run on petrol and on battery as well. Their production share in the market is 60%, 30% and 10% respectively. Of their respective production capacities, 20%, 10% and 5% cars respectively are electric (or battery operated). Based on the above, answer the following :</p>  <p>(i) (a) What is the probability that a randomly selected car is an electric car ? 2</p> <p style="text-align: center;">OR</p> <p>(i) (b) What is the probability that a randomly selected car is a petrol car ? 2</p> <p>(ii) A car is selected at random and is found to be electric. What is the probability that it was manufactured by Comet ? 1</p> <p>(iii) A car is selected at random and is found to be electric. What is the probability that it was manufactured by Amber or Bonzi ? 1</p>	
37(i) (a) Ans	<p>Let A = Amber manufactures the car B = Bonzi manufactures the car C = Comet manufactures the car E = The selected car is electric</p> $P(A) = \frac{60}{100}, P(B) = \frac{30}{100}, P(C) = \frac{10}{100}$ $P(E) = P(A) \times P\left(\frac{E}{A}\right) + P(B) \times P\left(\frac{E}{B}\right) + P(C) \times P\left(\frac{E}{C}\right)$ $= \frac{60}{100} \times \frac{20}{100} + \frac{30}{100} \times \frac{10}{100} + \frac{10}{100} \times \frac{5}{100}$ $= \frac{155}{1000} \text{ or } \frac{31}{200}$ <p style="text-align: center;">OR</p>	<p><math>\frac{1}{2}</math></p> <p>1</p> <p><math>\frac{1}{2}</math></p>
37(i)(b) Ans	<p>Let A = Amber manufactures the car B = Bonzi manufactures the car</p>	

	<p>C = Comet manufactures the car</p> <p>E = The selected car is a petrol car</p> $P(A) = \frac{60}{100}, P(B) = \frac{30}{100}, P(C) = \frac{10}{100}$ $P(E) = P(A) \times P\left(\frac{E}{A}\right) + P(B) \times P\left(\frac{E}{B}\right) + P(C) \times P\left(\frac{E}{C}\right)$ $= \frac{60}{100} \times \frac{80}{100} + \frac{30}{100} \times \frac{90}{100} + \frac{10}{100} \times \frac{95}{100}$ $= \frac{845}{1000} \text{ or } \frac{169}{200}$	<p>1/2</p> <p>1</p> <p>1/2</p>
<p>37(ii) Ans</p>	$P\left(\frac{C}{E}\right) = \frac{P(C) \times P\left(\frac{E}{C}\right)}{P(E)}$ $= \frac{\frac{10}{100} \times \frac{5}{100}}{\frac{60}{100} \times \frac{20}{100} + \frac{30}{100} \times \frac{10}{100} + \frac{10}{100} \times \frac{5}{100}}$ $= \frac{50}{1550} = \frac{1}{31}$	<p>1</p>
<p>37(iii) Ans</p>	$P\left(\frac{A \text{ or } B}{E}\right) = 1 - P\left(\frac{C}{E}\right) = 1 - \frac{1}{31} = \frac{30}{31}$	<p>1</p>
<p>38.</p>	 <p>A small town is analyzing the pattern of a new street light installation. The lights are set up in such a way that the intensity of light at any point <math>x</math> metres from the start of the street can be modelled by <math>f(x) = e^x \sin x</math>, where <math>x</math> is in metres.</p> <p>Based on the above, answer the following :</p> <p>(i) Find the intervals on which the <math>f(x)</math> is increasing or decreasing, <math>x \in [0, \pi]</math>. <span style="float: right;">2</span></p> <p>(ii) Verify, whether each critical point when <math>x \in [0, \pi]</math> is a point of local maximum or local minimum or a point of inflexion. <span style="float: right;">2</span></p>	

