







Q.No	Value points	Mark
SECTION A		
1	(C)	1
2	(D)	1
3	(D)	1
4	(B)	1
5	(A)	1
6	(A)	1
7	(B)	1
8	(D)	1
9	(B)	1
10	(C)	1
11	(C)	1
12	(B)	1
13	(C)	1
14	(D)	1
15	(C)	1
16	(D)	1
SECTION B		
17	<ul style="list-style-type: none"> Amino-acids which cannot be synthesized in the body and must be obtained through diet. In zwitter ionic form, amino-acids react both with acids and bases./ Due to the presence of both carboxylic group and amino group. 	1 1
18	Order of the reaction =1 / First Rate =k[A]	1 1
OR		
18	Rate of the reaction will increase. Rate constant remains same.	1 1
19	Structural formula: $K_2[PtCl_6]$ IUPAC Name: Potassium hexachloridoplatinate(IV)	1 1
20	Galvanic cell which converts the energy of combustion of fuels directly into electrical energy. Advantages 1.High efficiency 2.Pollution free (or any other two correct advantages)	1 $\frac{1}{2} \times 2 = 1$
21	<p>a)</p> $\left(\begin{array}{c} R \\ \\ R-C-O \\ \\ R \end{array} \right)_3 Al$ <p>b)</p>	1 1
SECTION C		
22(a)	(a)No Sodium methoxide is a strong nucleophile as well as a strong base so elimination reaction of t-butyl bromide predominates over substitution. Methyl bromide and Sodium.t-butoxide / CH_3Br and $(CH_3)_3CONa$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}, \frac{1}{2}$


	1.Pentose sugar+ Nitrogenous base + Phosphate	1.Pentose sugar+ Nitrogenous base	1
	(c) (i) To preserve genetic information and Protein synthesis OR (c)(ii)Phosphodiester linkage Uracil		1 ½ + ½
30	(a)Chloroform and Acetone A-B interactions are stronger than A-A and B-B interaction. (b)(i) For any solution the partial vapour pressure of each volatile component is directly proportional to its mole fraction. OR (b)(ii) $p = p^0 x_1$; $p = K_H x$ When $p^0 = K_H$ $p \propto \chi$ for both. (c) The enthalpy of mixing of the pure components in the ideal solution is Zero/ $\Delta_{mix}H=0$. The Volume of mixing of the pure components in the ideal solution is Zero. $\Delta_{mix}V=0$ (or any other two suitable characteristics)	(or Any other correct example)	1 1 1 1 ½ + ½
SECTION E			
31	(a) (i) $k = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$ $k = \frac{2.303}{60} \log \frac{1.2 \times 10^{-2}}{0.2 \times 10^{-2}}$ $= \frac{2.303}{60} \log 6$ $= \frac{2.303}{60} \times 0.778$ $k = 2.98 \times 10^{-2} \text{ min}^{-1} / 0.0298 \text{ min}^{-1}$ (ii) (I) Order is determined experimentally. If one of the reactants is taken in excess.		1 1 1 1 1
OR			
31	(b)(i) $\log \frac{k_2}{k_1} = \frac{E_a}{2.303R} \left[\frac{1}{T_1} - \frac{1}{T_2} \right]$ $\log \frac{2k_1}{k_1} = \frac{E_a}{19.15} \left[\frac{1}{298} - \frac{1}{308} \right]$ $0.3 = \frac{E_a}{19.15} \left[\frac{10}{298 \times 308} \right]$ $E_a = \frac{0.3 \times 19.15 \times 298 \times 308}{10}$ $E_a = 52729 \text{ Jmol}^{-1} \text{ or } 52.729 \text{ kJmol}^{-1}$ (ii) (1). Rate= $k[\text{H}_2\text{O}_2] [\text{I}^-]$ (2) Overall order : 2/ Second Molecularity : 2 / Bimolecular		1 1 1 1 ½ ½

32	<p>(a)(i) (I)</p>  <p>(II) CH_3COCH_3</p> <p>(III)</p>  <p>(ii) (I) Benzoic acid with Sodium bicarbonate gives brisk effervescence. No reaction with Ethyl benzoate (ii) Propanal, when heated with ammoniacal solution of silver nitrate (Tollens' reagent) gives silver mirror. No reaction with propanone (or any other suitable chemical test)</p>	1 1 1 1 1
OR		
32	<p>(b)(i)(I)</p>  <p>(II) 1. $(\text{BH}_3)_2$, 2. $\text{H}_2\text{O}_2/\text{OH}^-$, 3. PCC</p> <p>(III)</p>  <p>(b)(ii) (I)</p>  <p>(II) $\text{C}_6\text{H}_5\text{CHO} \xrightarrow{\text{CH}_3\text{CHO, dil NaOH, } \Delta} \text{C}_6\text{H}_5\text{CH}=\text{CHCHO} \xrightarrow{\text{H}_2/\text{Ni}} \text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ (Or any other suitable method)</p>	1 1 1 1 1
33	<p>(a)(i) (I) CO being a strong field ligand, causes pairing of electrons therefore, there is no unpaired electron. Whereas Cl^- is a weak field ligand, does not cause pairing, therefore presence of unpaired electrons.</p> <p>(II) CO can form both sigma (σ) and pi (π) bond with central metal atom/Metal to ligand bonding creates synergic effect between CO and the Metal.</p> <p>(III) Mirror images are superimposable/ Presence of plane of symmetry.</p> <p>(ii) (I) $\Delta_0 > P$, causes pairing of electrons, therefore 1 unpaired electron (II) $\Delta_0 < P$, No pairing of electrons therefore 5 unpaired electrons</p>	1 1 1 1 1
OR		
33	<p>(b)(i) (I) Coordination Isomerism / $[\text{Cr}(\text{NH}_3)_6][\text{Co}(\text{CN})_6]$</p>	$\frac{1}{2}, \frac{1}{2}$

(II) Optical Isomerism /



(III) Geometrical isomerism /



(ii) Weak field ligands produce weak field and leads to small splitting of d-orbitals whereas strong field ligands produce strong field leading to large splitting of d-orbitals.

Strong field ligands cause pairing of electrons/a smaller number of unpaired electrons hence produces low spin complexes and weak field ligands causes no pairing of electrons/a greater number of unpaired electrons hence produces high spin complexes.

1/2, 1/2

1/2, 1/2

1

1