










Q.No	Value points	Mark										
SECTION A												
1	A	1										
2	D	1										
3	B	1										
4	C	1										
5	B	1										
6	B	1										
7	A	1										
8	C	1										
9	B	1										
10	D	1										
11	C	1										
12	B	1										
13	D	1										
14	A	1										
15	B	1										
16	A	1										
SECTION B												
17	a) Negative deviation Because A-B interactions are stronger than A-A and B-B interactions b) Because of same composition in liquid and vapour phase	$\frac{1}{2}$ $\frac{1}{2}$ 1										
18	Because molecularity of each elementary reaction in complex reaction may be different and hence meaningless for overall complex reaction whereas order of a complex reaction is experimentally determined by the slowest step in its mechanism and is therefore applicable for both.	1 1										
19	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;">DNA</th> <th style="width: 50%;">RNA</th> </tr> </thead> <tbody> <tr> <td>Double stranded</td> <td>Single stranded</td> </tr> <tr> <td>Sugar is deoxyribose</td> <td>Sugar is ribose</td> </tr> <tr> <td>Thymine base is present</td> <td>Uracil base is present</td> </tr> <tr> <td>It replicates</td> <td>It does not replicate</td> </tr> </tbody> </table> <p style="text-align: right;">(Any Two)</p>	DNA	RNA	Double stranded	Single stranded	Sugar is deoxyribose	Sugar is ribose	Thymine base is present	Uracil base is present	It replicates	It does not replicate	1+1
DNA	RNA											
Double stranded	Single stranded											
Sugar is deoxyribose	Sugar is ribose											
Thymine base is present	Uracil base is present											
It replicates	It does not replicate											
20	a) Dichloridobis(ethane-1,2-diamine)platinum(IV) ion 	1 1										
OR												
20	i) $[\text{Co}(\text{NH}_3)_5(\text{CO}_3)]\text{Cl}$ ii) Pentaamminenitrito-O-cobalt(III) chloride	1 1										
21	-Because C—X bond acquires a partial double bond character due to resonance/ sp^2 hybridized carbon of C-X bond leading to shorter bond length (Or any other suitable reason). -Nitro group withdraws the electron density from the benzene ring and thus facilitates the attack of the nucleophile on haloarene / $-\text{NO}_2$ group being electron withdrawing stabilises the intermediate carbanion.	1 1										

SECTION C		
22	(a) Lead storage battery Anode: $\text{Pb(s)} + \text{SO}_4^{2-}(\text{aq}) \rightarrow \text{PbSO}_4(\text{s}) + 2\text{e}^-$ Cathode: $\text{PbO}_2(\text{s}) + \text{SO}_4^{2-}(\text{aq}) + 4\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{PbSO}_4(\text{s}) + 2\text{H}_2\text{O}(\text{l})$	1 1 1
OR		
22	(b) Because at cathode the reaction with higher value of E° is preferred and therefore, the reduction of H_2O to H_2 gas is preferred whereas at anode water should get oxidised in preference to $\text{Cl}^-(\text{aq})$, however, on account of overpotential of oxygen, oxidation of Cl^- to Cl_2 gas is preferred. $\text{NaCl}(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightarrow \text{Na}^+(\text{aq}) + \text{OH}^-(\text{aq}) + \frac{1}{2}\text{H}_2(\text{g}) + \frac{1}{2}\text{Cl}_2(\text{g})$	1 1 1
23	$\Delta T_f = iK_f m$ $\Delta T_f = \frac{i \times K_f \times w_2 \times 1000}{M_2 \times w_1}$ $0.45 = \frac{i \times 5.12 \times 0.3 \times 1000}{60 \times 30}$ $i = 0.527$ $\alpha = \frac{i-1}{1/n-1}$ $\alpha = \frac{0.527-1}{1/2-1} \quad (n=2)$ $\alpha = 0.946 \text{ or } 94.6\%$ <p style="text-align: right;">(Or any other suitable method)</p>	$\frac{1}{2}$ 1 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
24	A= $\begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH}_2 - \text{I} \\ \\ \text{CH}_3 \end{array}$ B= $\begin{array}{c} \text{CH}_3 - \text{C} = \text{CH}_2 \\ \\ \text{CH}_3 \end{array}$ C= $\begin{array}{c} \text{I} \\ \\ \text{CH}_3 - \text{CH} - \text{CH}_3 \\ \\ \text{CH}_3 \end{array}$ D= $\begin{array}{c} \text{CH}_3\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_3 \\ \qquad \qquad \qquad \\ \text{CH}_3 \qquad \qquad \qquad \text{CH}_3 \end{array}$ $\begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH}_2 - \text{I} \\ \\ \text{CH}_3 \end{array} \xrightarrow{\text{KOH(alc)/}\Delta} \begin{array}{c} \text{CH}_3 - \text{C} = \text{CH}_2 \\ \\ \text{CH}_3 \end{array} + \text{KI} + \text{H}_2\text{O}$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 1

<p>25</p>	$2\text{N}_2\text{O}_5(\text{g}) \rightarrow 2\text{N}_2\text{O}_4(\text{g}) + \text{O}_2(\text{g})$ <p>Start $t = 0$ P_i atm 0 atm 0 atm At time t $(P_i - 2x)$ atm 2x atm x atm</p> <p>$P_t = P_i - 2x + 2x + x = P_i + x$ $x = P_t - P_i$ $p_A = P_i - 2x$ $= P_i - 2(P_t - P_i)$ $= 3P_i - 2P_t$</p> <p>$k = \frac{2.303}{t} \log \frac{P_i}{p_A}$</p> <p>Where $p_i = 0.5$ atm, $p_A = 3p_i - 2p_t$ $= (3 \times 0.5) - (2 \times 0.625)$ $= 0.25$ atm</p> <p>$k = \frac{2.303}{100\text{s}} \log \frac{0.5\text{ atm}}{0.25\text{ atm}}$</p> <p>$= \frac{2.303}{100\text{s}} \times 0.3010$ $= 6.93 \times 10^{-3} \text{ s}^{-1}$</p>	<p>1</p> <p>1</p> <p>1</p>
<p>26.</p>	<p>a)</p> $\begin{array}{c} \text{CHO} \\ \\ (\text{CHOH})_4 \\ \\ \text{CH}_2\text{OH} \end{array} \xrightarrow{\text{HCN}} \begin{array}{c} \text{CH} \begin{array}{l} \nearrow \text{CN} \\ \searrow \text{OH} \end{array} \\ \\ (\text{CHOH})_4 \\ \\ \text{CH}_2\text{OH} \end{array}$ <p>b)</p> $\begin{array}{c} \text{CHO} \\ \\ (\text{CHOH})_4 \\ \\ \text{CH}_2\text{OH} \end{array} \xrightarrow{\text{Br}_2 \text{ water}} \begin{array}{c} \text{COOH} \\ \\ (\text{CHOH})_4 \\ \\ \text{CH}_2\text{OH} \end{array}$ <p>c)</p> $\begin{array}{c} \text{CHO} \\ \\ (\text{CHOH})_4 \\ \\ \text{CH}_2\text{OH} \end{array} \xrightarrow{\text{Acetic anhydride}} \begin{array}{c} \text{CHO} \quad \text{O} \\ \quad \parallel \\ (\text{CH}-\text{O}-\text{C}-\text{CH}_3)_4 \\ \quad \parallel \\ \text{CH}_2-\text{O}-\text{C}-\text{CH}_3 \end{array}$	<p>1</p> <p>1</p> <p>1</p>
<p>27</p>	<p>a) $\text{CH}_3\text{I} +$ </p> <p>b) </p> <p>c) </p>	<p>1</p> <p>1</p> <p>1</p>

28	<p>a) Because the carboxyl group is deactivating and the catalyst aluminium chloride (Lewis acid) gets bonded to the carboxyl group.(forms salt)</p> <p>b) Because carbonyl carbon of HCHO is more electrophilic than CH₃CHO/ due to +I effect of methyl group/ steric effect of methyl group, CH₃CHO is less reactive.</p> <p>c) Because of greater electronegativity of sp² hybridised carbon to which carboxyl carbon is attached.</p>	1 1 1
SECTION D		
29	<p>a) Due to presence of one unpaired electron in t_{2g} which gets excited to e_g / Due to excitation energy t_{2g}¹ → e_g¹, it gives colour. (d-d transition) When heated, water is lost therefore crystal field splitting does not occur and it becomes colourless.</p> <p>b) The energy required to split the degenerate d-orbitals into two sets of orbitals (t_{2g} and e_g). /The difference of energy between the two sets of d-orbitals t_{2g} and e_g due to the presence of ligands in a definite geometry .</p> <p style="text-align: center;">OR</p> <p>b) (ii) Δ_o < P, weak field ligand Δ_o > P, strong field ligand</p> <p>c) Because the orbital splitting energies are not sufficiently large for forcing pairing / Due to low crystal field splitting energy.</p>	1 1 1 ½ + ½ 1
30	<p>a) (i)</p> $\begin{array}{ccc} \text{CH}_3\text{COOH} & \xrightarrow{\text{NH}_3, \text{Heat}} & \text{CH}_3\text{CONH}_2 \\ & & \downarrow \text{Br}_2/\text{NaOH} \\ & & \text{CH}_3\text{NH}_2 \end{array}$ <p>(ii)</p> $\text{CH}_3\text{-CH}_2\text{-C}\equiv\text{N} \xrightarrow{\text{H}_2/\text{Pt}} \text{CH}_3\text{-CH}_2\text{-CH}_2\text{-NH}_2$ <p style="text-align: right;">(or by any other method)</p> <p>b) Aniline undergoes resonance and as a result the electrons on the N-atom are less available for donation.</p> <p>c) (i) (CH₃)₃N < CH₃NH₂ < (CH₃)₂NH OR c) (ii) A = C₆H₅NH₂ ; B = =C₆H₅N₂⁺Cl</p>	1 1 1 1 ½ + ½
SECTION E		
31	<p>(a) (i) (I) (CH₃)₃C-CHO (II)</p> $\begin{array}{ccccccc} & & \text{CH}_3 & & \text{O} & & \\ & & & & & & \\ \text{H}_3\text{C} & - & \text{CH}_2 & - & \text{CH} & - & \text{C} & - & \text{H} \end{array}$ <p>(III) CH₃-CO-CH₂CH₂CH₃</p> <p>(ii)</p> $\text{>C=O} \xrightarrow[\text{HCl}]{\text{Zn-Hg}} \text{>CH}_2 + \text{H}_2\text{O}$ <p>(I)</p>	1 1 1 1

	<p>(II)</p> 	1
OR		
31	<p>(b)</p> <p>i)</p>  <p>ii) Because semicarbazide undergoes resonance involving only one of the two -NH₂ groups, which is attached directly to the carbonyl-carbon atom.</p> <p>iii)</p>  <p>iv)</p>  <p>v)</p> 	1 1 1 1 1
32	<p>(a) (i)</p> $E_{\text{Cell}} = (E^{\circ}_c - E^{\circ}_a) - \frac{0.059}{2} \log \frac{[\text{Zn}^{2+}]}{[\text{H}^+]^2}$ $= [(0) - (-0.76)] - \frac{0.059}{2} \log \frac{0.1}{(0.01)^2}$ $= 0.76 - 0.0295 \log 10^3$ $= 0.76 - 0.0885$ $= 0.6715 \text{ V} \quad \text{or } 0.67 \text{ V} \quad (\text{Deduct } \frac{1}{2} \text{ mark for no or incorrect unit})$ <p>(ii) The amounts of different substances liberated by the same quantity of electricity passing through the electrolytic solution are proportional to their chemical equivalent weights.</p> <p>6F</p>	1 1 1 1
OR		

32	<p>(b) (i)</p> $\Lambda_m = \frac{K}{C} \times 1000$ $\Lambda_m = \frac{2.48 \times 10^{-2}}{0.2} \times 1000$ $= 124 \text{ S cm}^2 \text{ mol}^{-1}$ $\Lambda_m^\circ = \lambda_{\text{K}^+}^\circ + \lambda_{\text{Cl}^-}^\circ$ $= 73.5 + 76.5$ $= 150 \text{ S cm}^2 \text{ mol}^{-1}$ $\alpha = \frac{\Lambda_m}{\Lambda_m^\circ}$ $= \frac{124}{150}$ $= 0.827$ <p>(ii) $E^\circ_{\text{cell}} = E^\circ_{\text{cathode}} - E^\circ_{\text{anode}}$ $= 0.34 - (-2.37)$ $= 2.71 \text{ V}$</p> <p>$\Delta_r G^\circ = -nFE^\circ_{\text{cell}}$ $= -2 \times 96500 \times 2.71$ $= -523030 \text{ J mol}^{-1} \text{ or } -523.03 \text{ kJ mol}^{-1}$</p> <p>(iii) Primary cell Maintains constant potential throughout its usage/ longer lifespan</p>	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>
33	<p>(a) (i) (I) Because Mn^{2+} is more stable than Mn^{3+} due to extra stable half-filled d^5 configuration. (II) Due to comparable energies of 5f, 6d and 7s orbitals (III) Due to the involvement of greater number of electrons from (n-1)d in addition to the ns electrons in the inter-atomic metallic bonding.</p> <p>(ii) (I) $5\text{SO}_3^{2-} + 2\text{MnO}_4^- + 6\text{H}^+ \longrightarrow 2\text{Mn}^{2+} + 3\text{H}_2\text{O} + 5\text{SO}_4^{2-}$ (II) $2\text{MnO}_4^- + \text{H}_2\text{O} + \Gamma \longrightarrow 2\text{MnO}_2 + 2\text{OH}^- + \text{IO}_3^-$</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
OR		
33	<p>b) (i) Mn, Zn, Ni, Cu (any two) (ii) K_2MnO_4, due to presence of one unpaired electron (iii) Similar radii of 4d and 5d series elements/ similar properties/ difficulty in separation of lanthanoids (or any other relevant consequence) (iv) It is prepared by fusion of MnO_2 with an alkali metal hydroxide and an oxidising agent / $2\text{MnO}_2 + 4\text{KOH} + \text{O}_2 \rightarrow 2\text{K}_2\text{MnO}_4 + 2\text{H}_2\text{O}$ (v) because of the ability of oxygen to form multiple bonds with metal</p>	<p>$\frac{1}{2}$, $\frac{1}{2}$</p> <p>$\frac{1}{2}$, $\frac{1}{2}$</p> <p>1</p> <p>1</p> <p>1</p>

