

LIMITS

1. Let α be a positive real number. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ and $g: (\alpha, \infty) \rightarrow \mathbb{R}$ be the functions defined by

$$f(x) = \sin\left(\frac{\pi x}{12}\right) \text{ and } g(x) = \frac{2 \log_e\left(\sqrt{x} - \sqrt{a}\right)}{\log_e\left(e^{\sqrt{x}} - e^{\sqrt{a}}\right)}.$$

Then the value of $\lim_{x \rightarrow a^+} f(g(x))$ is _____.

[JEE(Advanced) 2022]

2. If

$$\beta = \lim_{x \rightarrow 0} \frac{e^{x^3} - (1-x^3)^{\frac{1}{3}} + \left((1-x^2)^{\frac{1}{2}} - 1 \right) \sin x}{x \sin^2 x}$$

then the value of 6β is .

[JEE(Advanced) 2022]

3. Let e denote the base of the natural logarithm. The value of the real number a for which the right hand limit

$$\lim_{x \rightarrow 0^+} \frac{(1-x)^{\frac{1}{x}} - e^{-1}}{x^a}$$

is equal to a nonzero real number, is

[JEE(Advanced) 2020]

- #### 4. The value of the limit

$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{4\sqrt{2}(\sin 3x + \sin x)}{\left(2 \sin 2x \sin \frac{3x}{2} + \cos \frac{5x}{2}\right) - \left(\sqrt{2} + \sqrt{2} \cos 2x + \cos \frac{3x}{2}\right)}$$

is .

[JEE(Advanced) 2020]

5. Let $F : \mathbb{R} \rightarrow \mathbb{R}$ be a function. We say that f has

PROPERTY 1 if $\lim_{h \rightarrow 0} \frac{f(h) - f(0)}{\sqrt{|h|}}$ exists and is finite, and

PROPERTY 2 if $\lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h^2}$ exists and is finite.

Then which of the following options is/are correct?

[JEE(Advanced) 2019]

- (A) $f(x) = x|x|$ has PROPERTY 2 (B) $f(x) = x^{2/3}$ has PROPERTY 1
 (C) $f(x) = \sin x$ has PROPERTY 2 (D) $f(x) = |x|$ has PROPERTY 1

6. For any positive integer n , define $f_n : (0, \infty) \rightarrow \mathbb{R}$ as

$$f_n(x) = \sum_{j=1}^n \tan^{-1} \left(\frac{1}{1 + (x+j)(x+j-1)} \right) \text{ for all } x \in (0, \infty).$$

(Here, the inverse trigonometric function $\tan^{-1} x$ assume values in $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$)

Then, which of the following statement(s) is (are) TRUE?

[JEE(Advanced)-2018]

- (A) $\sum_{j=1}^5 \tan^2(f_j(0)) = 55$
 (B) $\sum_{j=1}^{10} (1 + f'_j(0)) \sec^2(f_j(0)) = 10$

(C) For any fixed positive integer n , $\lim_{x \rightarrow \infty} \tan(f_n(x)) = \frac{1}{n}$

(D) For any fixed positive integer n , $\lim_{x \rightarrow \infty} \sec^2(f_n(x)) = 1$

7. Let $f(x) = \frac{1-x(1+|1-x|)}{|1-x|} \cos\left(\frac{1}{1-x}\right)$ for $x \neq 1$. Then [JEE(Advanced) 2017]

(A) $\lim_{x \rightarrow 1^+} f(x)$ does not exist (B) $\lim_{x \rightarrow 1^-} f(x)$ does not exist
 (C) $\lim_{x \rightarrow 1^-} f(x) = 0$ (D) $\lim_{x \rightarrow 1^+} f(x) = 0$

8. Let $\alpha, \beta \in \mathbb{R}$ be such that $\lim_{x \rightarrow 0} \frac{x^2 \sin(\beta x)}{\alpha x - \sin x} = 1$. Then $6(\alpha + \beta)$ equals [JEE(Advanced) 2017]

9. Let $f(x) = \sin\left(\frac{\pi}{6} \sin\left(\frac{\pi}{2} \sin x\right)\right)$ for all $x \in \mathbb{R}$ and $g(x) = \frac{\pi}{2} \sin x$ for all $x \in \mathbb{R}$. Let $(fog)(x)$ denotes $f(g(x))$ and $(gof)(x)$ denotes $g(f(x))$. Then which of the following is (are) true? [JEE(Advanced) 2015]

(A) Range of f is $\left[-\frac{1}{2}, \frac{1}{2}\right]$ (B) Range of fog is $\left[-\frac{1}{2}, \frac{1}{2}\right]$
 (C) $\lim_{x \rightarrow 0} \frac{f(x)}{g(x)} = \frac{\pi}{6}$ (D) There is an $x \in \mathbb{R}$ such that $(gof)(x) = 1$

10. Let m and n be two positive integers greater than 1. If

$$\lim_{\alpha \rightarrow 0} \left(\frac{e^{\cos(\alpha^n)} - e}{\alpha^m} \right) = -\left(\frac{e}{2}\right)$$

then the value of $\frac{m}{n}$ is [JEE(Advanced) 2015]

11. The largest value of the non-negative integer a for which $\lim_{x \rightarrow 1} \left\{ \frac{-ax + \sin(x-1) + a}{x + \sin(x-1) - 1} \right\}^{\frac{1-x}{1-\sqrt{x}}} = \frac{1}{4}$ is [JEE(Advanced) 2014]

SOLUTIONS

1. Ans. (0.50)

Sol. $\lim_{x \rightarrow a^+} \frac{2\ln(\sqrt{x} - \sqrt{a})}{\ln(e^{\sqrt{x}} - e^{\sqrt{a}})} \quad \left(\begin{matrix} 0 \\ 0 \end{matrix} \text{ form} \right)$

∴ Using Lopital rule,

$$= 2 \lim_{x \rightarrow a^+} \frac{\left(\frac{1}{\sqrt{x} - \sqrt{a}} \right) \cdot \frac{1}{2\sqrt{x}}}{\left(\frac{1}{e^{\sqrt{x}} - e^{\sqrt{a}}} \right) \cdot e^{\sqrt{x}} \cdot \frac{1}{2\sqrt{x}}}$$

$$= \frac{2}{e^{\sqrt{a}}} \lim_{x \rightarrow a^+} \frac{\left(e^{\sqrt{x}} - e^{\sqrt{a}} \right)}{\left(\sqrt{x} - \sqrt{a} \right)} \quad \left(\begin{matrix} 0 \\ 0 \end{matrix} \right)$$

$$= \frac{2}{e^{\sqrt{a}}} \lim_{x \rightarrow a^+} \frac{\left(e^{\sqrt{x}} \cdot \frac{1}{2\sqrt{x}} - 0 \right)}{\left(\frac{1}{2\sqrt{x}} - 0 \right)} = 2$$

$$\text{so, } \lim_{x \rightarrow a^+} f(g(x)) = \lim_{x \rightarrow a^+} f(2)$$

$$= f(2) = \sin \frac{\pi}{6} = \frac{1}{2} = 0.50$$

2. Ans. (5)

Sol. $\beta = \lim_{x \rightarrow 0} \frac{e^{x^3} - (1-x^3)^{1/3}}{x \sin^2 x \cdot x^2} + \frac{\left((1-x^2)^{1/2} - 1 \right) \sin x}{x \frac{\sin^2 x}{x^2} \cdot x^2}$

use expansion

$$\beta = \lim_{x \rightarrow 0} \frac{\left(1+x^3 \right) - \left(1 - \frac{x^3}{3} \right)}{x^3} + \lim_{x \rightarrow 0} \frac{\left(\left(1 - \frac{x^2}{2} \right) - 1 \right) \sin x}{x^2}$$

$$\beta = \lim_{x \rightarrow 0} \frac{4x^3}{3x^3} + \lim_{x \rightarrow 0} \frac{-x^2}{2x^2}$$

$$\beta = \frac{4}{3} - \frac{1}{2} = \frac{5}{6}$$

$$6\beta = 5$$

3. Ans. (1.00)

Sol. $\lim_{x \rightarrow 0^+} \frac{e^{\left(\frac{\ln(1-x)}{x} \right)} - 1}{x^a}$

$$= \lim_{x \rightarrow 0^+} \frac{e^{\left(1 + \frac{\ln(1-x)}{x} \right)} - 1}{x^a}$$

$$\begin{aligned} &= \frac{1}{e} \lim_{x \rightarrow 0^+} \frac{1 + \frac{\ln(1-x)}{x}}{x^a} \\ &= \frac{1}{e} \lim_{x \rightarrow 0^+} \frac{\ln(1-x) + x}{x^{a+1}} \\ &= \frac{1}{e} \lim_{x \rightarrow 0^+} \frac{\left(-x - \frac{x^2}{2} - \frac{x^3}{3} - \dots \right) + x}{x^{a+1}} \end{aligned}$$

Thus, $a = 1$

4. Ans. (8)

Sol.

$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{4\sqrt{2} \cdot 2 \sin 2x \cos x}{2 \sin 2x \sin \frac{3x}{2} + \left(\cos \frac{5x}{2} - \cos \frac{3x}{2} \right) - \sqrt{2}(1 + \cos 2x)}$$

$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{16\sqrt{2} \sin x \cos^2 x}{2 \sin 2x \left(\sin \frac{3x}{2} - \sin \frac{x}{2} \right) - 2\sqrt{2} \cos^2 x}$$

$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{16\sqrt{2} \sin x \cos^2 x}{4 \sin x \cos x \left(2 \cos x \cdot \sin \frac{x}{2} \right) - 2\sqrt{2} \cos^2 x}$$

$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{16\sqrt{2} \sin x}{8 \sin x \cdot \sin \frac{x}{2} - 2\sqrt{2}} = 8$$

5. Ans. (B, D)

Sol. P-1 :

$$\lim_{h \rightarrow 0} \frac{f(h) - f(0)}{\sqrt{|h|}} = \text{exist and finite}$$

$$(B) f(x) = x^{2/3}, \lim_{h \rightarrow 0} \frac{h^{2/3} - 0}{\sqrt{|h|}} = \lim_{h \rightarrow 0} \frac{|h|^{2/3}}{\sqrt{|h|}} = 0$$

$$(D) f(x) = |x|, \lim_{h \rightarrow 0} \frac{|h| - 0}{\sqrt{|h|}} \Rightarrow \lim_{h \rightarrow 0} \sqrt{|h|} = 0$$

P-2 :

$$\lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h^2} = \text{exist and finite}$$

$$(A) f(x) = x|x|,$$

$$\lim_{h \rightarrow 0} \frac{h|h| - 0}{h^2} = \begin{cases} \text{RHL} = \lim_{h \rightarrow 0} \frac{h^2}{h^2} = 1 \\ \text{LHL} = \lim_{h \rightarrow 0} \frac{-h^2}{h^2} = -1 \end{cases}$$

$$(C) f(x) = \sin x, \lim_{h \rightarrow 0} \frac{\sin h - 0}{h^2} = \text{DNE}$$

6. Ans. (D)

$$\text{Sol. } f_n(x) = \sum_{j=1}^n \tan^{-1} \left(\frac{(x+j)-(x+j-1)}{1+(x+j)(x+j-1)} \right)$$

$$f_n(x) = \sum_{j=1}^n [\tan^{-1}(x+j) - \tan^{-1}(x+j-1)]$$

$$f_n(x) = \tan^{-1}(x+n) - \tan^{-1}x$$

$$f_n'(x) = \frac{1}{1+(x+n)^2} - \frac{1}{1+x^2}$$

$$\therefore \tan(f_n(x)) = \tan[\tan^{-1}(x+n) - \tan^{-1}x]$$

$$\tan(f_n(x)) = \frac{(x+n)-x}{1+x(x+n)}$$

$$\tan(f_n(x)) = \frac{n}{1+x^2+nx}$$

$$\therefore \sec^2(f_n(x)) = 1 + \tan^2(f_n(x))$$

$$\sec^2(f_n(x)) = 1 + \left(\frac{n}{1+x^2+nx} \right)^2$$

$\because 0$ is not in the domain of f_n

so, no meaning of $f_j'(0)$ and $f_j(0)$

\therefore option (A) and (B) are wrong

(C) For any fixed positive integer n ,

$$\lim_{x \rightarrow \infty} \tan(f_n(x)) = \lim_{x \rightarrow \infty} \frac{n}{1+x^2+nx} = 0$$

(D) For any fixed positive integer n ,

$$\begin{aligned} \lim_{x \rightarrow \infty} \sec^2(f_n(x)) &= \lim_{x \rightarrow \infty} \left(1 + \left(\frac{n}{1+x^2+nx} \right)^2 \right) \\ &= 1 \end{aligned}$$

7. Ans. (A,C)

$$\text{Sol. } f(x) = \begin{cases} (1-x) \cos \frac{1}{1-x}, & x < 1 \\ -(1+x) \cos \frac{1}{1-x}, & x > 1 \end{cases}$$

$$\lim_{x \rightarrow 1^+} f(x) = \text{d.n.e}, \lim_{x \rightarrow 1^-} f(x) = 0$$

8. Ans. (7)

$$\text{Sol. If } \alpha \neq 1, \text{ then } \lim_{x \rightarrow 0} \frac{x \sin \beta x}{\alpha x - \sin x} = 0$$

$$\therefore \alpha = 1 \Rightarrow \lim_{x \rightarrow 0} \frac{\beta x^3 \sin \beta x}{x^3 \left(\frac{x - \sin x}{x^3} \right)} = \frac{\beta}{1/6}$$

$$\Rightarrow 6\beta = 1 \Rightarrow \beta = \frac{1}{6}$$

$$6(\alpha + \beta) = 7$$

9. Ans. (A, B, C)

$$\text{Sol. (A) } f(x) = \sin \left(\frac{\pi}{6} \sin \left(\frac{\pi}{2} \sin x \right) \right), x \in \mathbb{R}$$

$$= \sin \left(\frac{\pi}{6} \sin \theta \right), \theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2} \right]$$

$$= \sin \alpha, \alpha \in \left[-\frac{\pi}{6}, \frac{\pi}{6} \right]$$

$$\text{(B) } f(g(x)) = f(t), t \in \left[-\frac{\pi}{2}, \frac{\pi}{2} \right]$$

$$\Rightarrow f(t) \in \left[-\frac{1}{2}, \frac{1}{2} \right]$$

$$\text{(C) } \lim_{x \rightarrow 0} \frac{\sin \left(\frac{\pi}{6} \sin \left(\frac{\pi}{2} \sin x \right) \right)}{\frac{\pi}{6} \sin \left(\frac{\pi}{2} \sin x \right)} \frac{\frac{\pi}{6} \sin \left(\frac{\pi}{2} \sin x \right)}{\frac{\pi}{2} \sin x}$$

$$= 1 \cdot \frac{\pi}{6}$$

$$\text{(D) } g(f(x)) = 1 \Rightarrow \sin f(x) = \frac{2}{\pi}$$

$$\text{but } f(x) \in \left[-\frac{1}{2}, \frac{1}{2} \right] \subset \left[-\frac{\pi}{6}, \frac{\pi}{6} \right]$$

$$\Rightarrow \sin f(x) \in \left[-\frac{1}{2}, \frac{1}{2} \right] \text{ so no solutions}$$

10. Ans. (2)

$$\text{Sol. } \lim_{\alpha \rightarrow 0} \frac{e(e^{\cos \alpha^n - 1} - 1)}{(\cos \alpha^n - 1)} \frac{(\cos \alpha^n - 1)}{(\alpha^n)^2} \alpha^{2n-m}$$

$$= -\frac{e}{2} \quad \therefore 2n = m \Rightarrow \frac{m}{n} = 2$$

11. Ans. (0)

$$\text{Sol. } \lim_{x \rightarrow 1} \left\{ \frac{\sin(x-1) + a(1-x)}{(x-1) + \sin(x-1)} \right\}^{\frac{(1+\sqrt{x})(1-\sqrt{x})}{1-\sqrt{x}}} = \frac{1}{4}$$

$$\lim_{x \rightarrow 1} \left\{ \frac{\sin(x-1) - a}{(x-1) + \frac{\sin(x-1)}{(x-1)}} \right\}^{1+\sqrt{x}} = \frac{1}{4} \Rightarrow \left(\frac{1-a}{2} \right)^2 = \frac{1}{4}$$

$$\Rightarrow (a-1)^2 = 1 \Rightarrow a = 2 \text{ or } 0$$

but for $a = 2$ base of above limit approaches $-\frac{1}{2}$ and exponent approaches to 2 and since base cannot be negative hence limit does not exist.