

### BINOMIAL THEOREM

1. Let  $a$  and  $b$  be two nonzero real numbers. If the coefficient of  $x^5$  in the expansion of  $\left(ax^2 + \frac{70}{27bx}\right)^4$  is equal to the coefficient of  $x^{-5}$  is equal to the coefficient of  $\left(ax - \frac{1}{bx^2}\right)^7$ , then the value of  $2b$  is

[JEE(Advanced) 2023]

2. For non-negative integers  $s$  and  $r$ , let  $\binom{s}{r} = \begin{cases} \frac{s!}{r!(s-r)!} & \text{if } r \leq s, \\ 0 & \text{if } r > s. \end{cases}$

For positive integers  $m$  and  $n$ , let  $g(m, n) = \sum_{p=0}^{m+n} \frac{f(m, n, p)}{\binom{n+p}{p}}$

where for any nonnegative integer  $p$ ,  $f(m, n, p) = \sum_{i=0}^p \binom{m}{i} \binom{n+i}{p} \binom{p+n}{p-i}$

Then which of the following statements is/are TRUE?

[JEE(Advanced) 2020]

- (A)  $g(m, n) = g(n, m)$  for all positive integers  $m, n$
- (B)  $g(m, n+1) = g(m+1, n)$  for all positive integers  $m, n$
- (C)  $g(2m, 2n) = 2g(m, n)$  for all positive integers  $m, n$
- (D)  $g(2m, 2n) = (g(m, n))^2$  for all positive integers  $m, n$

3. Suppose  $\det \begin{bmatrix} \sum_{k=0}^n k & \sum_{k=0}^n {}^n C_k k^2 \\ \sum_{k=0}^n {}^n C_k k & \sum_{k=0}^n {}^n C_k 3^k \end{bmatrix} = 0$ , holds for some positive integer  $n$ . Then  $\sum_{k=0}^n \frac{{}^n C_k}{k+1}$  equals

[JEE(Advanced) 2019]

4. Let  $X = ({}^{10} C_1)^2 + 2({}^{10} C_2)^2 + 3({}^{10} C_3)^2 + \dots + 10({}^{10} C_{10})^2$ , where  ${}^{10} C_r$ ,  $r \in \{1, 2, \dots, 10\}$  denote binomial coefficients. Then, the value of  $\frac{1}{1430} X$  is \_\_\_\_\_. [JEE(Advanced) 2018]

5. Let  $m$  be the smallest positive integer such that the coefficient of  $x^2$  in the expansion of  $(1+x)^2 + (1+x)^3 + \dots + (1+x)^{49} + (1+mx)^{50}$  is  $(3n+1)^{51} {}^5 C_3$  for some positive integer  $n$ . Then the value of  $n$  is [JEE(Advanced) 2016]

6. The coefficient of  $x^9$  in the expansion of  $(1+x)(1+x^2)(1+x^3)\dots(1+x^{100})$  is [JEE(Advanced) 2015]

7. Coefficient of  $x^{11}$  in the expansion of  $(1+x^2)^4(1+x^3)^7(1+x^4)^{12}$  is - [JEE(Advanced) 2014]

- (A) 1051
- (B) 1106
- (C) 1113
- (D) 1120

## SOLUTIONS

### 1. Ans. (3)

**Sol.**  $T_{r+1} = {}^4C_r (a \cdot x^2)^{4-r} \cdot \left(\frac{70}{27bx}\right)^r$

$$= {}^4C_r \cdot a^{4-r} \cdot \frac{70^r}{(27b)^r} \cdot x^{8-3r}$$

here  $8 - 3r = 5$

$$8 - 5 = 3r \Rightarrow r = 1$$

$$\therefore \text{coeff.} = 4 \cdot a^3 \cdot \frac{70}{27b}$$

$$T_{r+1} = {}^7C_r (ax)^{7-r} \left(\frac{-1}{bx^2}\right)^r$$

$$= {}^7C_r \cdot a^{7-r} \left(\frac{-1}{b}\right)^r \cdot x^{7-3r}$$

$$7 - 3r = -5 \Rightarrow 12 = 3r \Rightarrow r = 4$$

$$\text{coeff. : } {}^7C_4 \cdot a^3 \cdot \left(\frac{-1}{b}\right)^4 = \frac{35a^3}{b^4}$$

$$\text{now } \frac{35a^3}{b^4} = \frac{280a^3}{27b}$$

$$b^3 = \frac{35 \times 27}{280} = b = \frac{3}{2} \Rightarrow 2b = 3$$

### 2. Ans. (A, B, D)

**Sol.** Solving

$$\begin{aligned} f(m, n, p) &= \sum_{i=0}^p {}^mC_i \cdot {}^{n+i}C_p \cdot {}^{p+n}C_{p-i} \\ &= {}^mC_i \cdot {}^{n+i}C_p \cdot {}^{p+n}C_{p-i} \\ &= {}^mC_i \cdot \frac{(n+i)!}{p!(n-p+i)!} \times \frac{(n+p)!}{(p-i)!(n+i)!} \\ &= {}^mC_i \times \frac{(n+p)!}{p!} \times \frac{1}{(n-p+i)!(p-i)!} \\ &= {}^mC_i \times \frac{(n+p)!}{p!n!} \times \frac{n!}{(n-p+i)!(p-i)!} \end{aligned}$$

$${}^mC_i \cdot {}^{n+p}C_p \cdot {}^nC_{p-i} \quad \left\{ {}^mC_i \cdot {}^nC_{p-i} = {}^{m+n}C_p \right\}$$

$$f(m, n, p) = {}^{n+p}C_p \cdot {}^{m+n}C_p$$

$$\frac{f(m, n, p)}{{}^{n+p}C_p} = {}^{m+n}C_p$$

$$\text{Now, } g(m, n) = \sum_{p=0}^{m+n} \frac{f(m, n, p)}{{}^{n+p}C_p}$$

$$g(m, n) = \sum_{p=0}^{m+n} {}^{m+n}C_p$$

$$g(m, n) = 2^{m+n}$$

$$(A) g(m, n) = g(n, m)$$

$$(B) g(m, n+1) = 2^{m+n+1}$$

$$g(m+n, n) = 2^{m+1+n}$$

$$(D) g(2m, 2n) = 2^{2m+2n}$$

$$= (2^{m+n})^2$$

$$= (g(m, n))^2$$

### 3. Ans. (6.20)

**Sol.** Suppose

$$\left| \begin{array}{l} \frac{n(n+1)}{2} - n(n-1) \cdot 2^{n-2} + n \cdot 2^{n-1} \\ n \cdot 2^{n-1} \quad 4^n \end{array} \right| = 0$$

$$\frac{n(n+1)}{2} \cdot 4^n - n^2(n-1) \cdot 2^{2n-3} - n^2 \cdot 2^{2n-2} = 0$$

$$\frac{n(n+1)}{2} - \frac{n^2(n-1)}{8} - \frac{n^2}{4} = 0$$

$$n^2 - 3n - 4 = 0$$

$$n = 4$$

$$\text{Now } \sum_{k=0}^4 \frac{{}^4C_k}{k+1} = \sum_{k=0}^4 \frac{k+1}{5} \cdot {}^5C_{k+1} \frac{1}{k+1}$$

$$= \frac{1}{5} \cdot [{}^5C_1 + {}^5C_2 + {}^5C_3 + {}^5C_4 + {}^5C_5]$$

$$= \frac{1}{5} [2^5 - 1] = \frac{31}{5} = 6.20$$

### 4. Ans. (646)

**Sol.**  $X = \sum_{r=0}^n r \cdot ({}^nC_r)^2 ; n = 10$

$$X = n \cdot \sum_{r=0}^n {}^nC_r \cdot {}^{n-1}C_{r-1}$$

$$X = n \cdot \sum_{r=1}^n {}^nC_{n-r} \cdot {}^{n-1}C_{r-1}$$

$$X = n \cdot {}^{2n-1}C_{n-1} ; n = 10$$

$$X = 10 \cdot {}^{19}C_9$$

$$\frac{X}{1430} = \frac{1}{143} \cdot {}^{19}C_9$$

$$= 646$$

**5. Ans. (5)**

**Sol.** Coefficient of  $x^2$  in the expansion of

$$(1+x)^2 + (1+x)^3 + \dots + (1+x)^{49} + (1+mx)^{50}$$

$${}^2C_2 + {}^3C_2 + \dots + {}^{49}C_2 + {}^{50}C_2m^2 = (3n+1){}^{51}C_3$$

$${}^3C_3 + {}^3C_2 + \dots + {}^{49}C_2 + {}^{50}C_2m^2 = (3n+1){}^{51}C_3$$

$${}^{50}C_3 + {}^{50}C_2m^2 = (3n+1){}^{51}C_3$$

$$\frac{50 \cdot 49 \cdot 48}{6} + \frac{50 \cdot 49}{2} m^2 = (3n+1) \frac{51 \cdot 50 \cdot 49}{6}$$

$$m^2 = 51n + 1$$

must be a perfect square

$$\Rightarrow n = 5 \text{ and } m = 16$$

**6. Ans. (8)**

**Sol.** There are 8 products

$$1^{99}x^9, 1^{98}x^8x, 1^{98}x^2x^7, 1^{98}x^3x^6, 1^{98}x^4x^5$$

$$1^{97}x^2x^6, 1^{97}x^3x^5, 1^{97}x^2x^3x^4$$

which generate  $x^9$  so coeff. is 8

**7. Ans. (C)**

**Sol.** Coefficient of  $x^{11}$  in

$$(1+x^2)^4(1+x^3)^7(1+x^4)^{12}$$

$$= {}^4C_0. {}^7C_1. {}^{12}C_2 + {}^4C_1. {}^7C_3. {}^{12}C_0 + {}^4C_2. {}^7C_1. {}^{12}C_1 +$$

$${}^4C_4. {}^7C_1. {}^{12}C_0$$

$$= 462 + 140 + 504 + 7 = 1113$$