

PARABOLA

1. Let P be a point on the parabola $y^2 = 4ax$, where $a > 0$. The normal to the parabola at P meets the x-axis at a point Q. The area of the triangle PFQ, where F is the focus of the parabola, is 120. If the slope m of the normal and a are both positive integers, then the pair (a,m) is **[JEE(Advanced) 2023]**
 (A) (2, 3) (B) (1, 3) (C) (2, 4) (D) (3, 4)

2. Consider the parabola $y^2 = 4x$. Let S be the focus of the parabola. A pair of tangents drawn to the parabola from the point $P = (-2, 1)$ meet the parabola at P_1 and P_2 . Let Q_1 and Q_2 be points on the lines SP_1 and SP_2 respectively such that PQ_1 is perpendicular to SP_1 and PQ_2 is perpendicular to SP_2 . Then, which of the following is/are TRUE? **[JEE(Advanced) 2022]**
 (A) $SQ_1 = 2$ (B) $Q_1Q_2 = \frac{3\sqrt{10}}{5}$ (C) $PQ_1 = 3$ (D) $SQ_2 = 1$

3. Let E denote the parabola $y^2 = 8x$. Let $P = (-2, 4)$, and let Q and Q' be two distinct points on E such that the lines PQ and PQ' are tangents to E. Let F be the focus of E. Then which of the following statements is (are) TRUE? **[JEE(Advanced) 2021]**
 (A) The triangle PFQ is a right-angled triangle
 (B) The triangle QPQ' is a right-angled triangle
 (C) The distance between P and F is $5\sqrt{2}$
 (D) F lies on the line joining Q and Q'

Question Stem for Questions Nos. 4 and 5

Question Stem

Consider the region $R = \{(x, y) \in \mathbb{R} \times \mathbb{R} : x \geq 0 \text{ and } y^2 \leq 4 - x\}$. Let F be the family of all circles that are contained in R and have centers on the x-axis. Let C be the circle that has largest radius among the circles in F . Let (α, β) be a point where the circle C meets the curve $y^2 = 4 - x$.

4. The radius of the circle C is _____. **[JEE(Advanced) 2021]**
5. The value of α is _____. **[JEE(Advanced) 2021]**
6. If a chord, which is not a tangent, of the parabola $y^2 = 16x$ has the equation $2x + y = p$, and midpoint (h, k) , then which of the following is(are) possible value(s) of p, h and k? **[JEE(Advanced) 2017]**
 (A) $p = 5, h = 4, k = -3$ (B) $p = -1, h = 1, k = -3$
 (C) $p = -2, h = 2, k = -4$ (D) $p = 2, h = 3, k = -4$
7. Let P be the point on the parabola $y^2 = 4x$ which is at the shortest distance from the center S of the circle $x^2 + y^2 - 4x - 16y + 64 = 0$. Let Q be the point on the circle dividing the line segment SP internally. Then-
 (A) $SP = 2\sqrt{5}$
 (B) $SQ : QP = (\sqrt{5} + 1) : 2$
 (C) the x-intercept of the normal to the parabola at P is 6
 (D) the slope of the tangent to the circle at Q is $\frac{1}{2}$ **[JEE(Advanced) 2016]**

8. If the normals of the parabola $y^2 = 4x$ drawn at the end points of its latus rectum are tangents to the circle $(x - 3)^2 + (y + 2)^2 = r^2$, then the value of r^2 is **[JEE(Advanced) 2015]**
9. Let the curve C be the mirror image of the parabola $y^2 = 4x$ with respect to the line $x + y + 4 = 0$. If A and B are the points of intersection of C with the line $y = -5$, then the distance between A and B is **[JEE(Advanced) 2015]**
10. Let P and Q be distinct points on the parabola $y^2 = 2x$ such that a circle with PQ as diameter passes through the vertex O of the parabola. If P lies in the first quadrant and the area of the triangle ΔOPQ is $3\sqrt{2}$, then which of the following is(are) the coordinates of P ? **[JEE(Advanced) 2015]**
- (A) $(4, 2\sqrt{2})$ (B) $(9, 3\sqrt{2})$ (C) $(\frac{1}{4}, \frac{1}{\sqrt{2}})$ (D) $(1, \sqrt{2})$
11. The common tangents to the circle $x^2 + y^2 = 2$ and the parabola $y^2 = 8x$ touch the circle at the point P, Q and the parabola at the points R, S . Then the area of the quadrilateral $PQRS$ is - **[JEE(Advanced) 2014]**
- (A) 3 (B) 6 (C) 9 (D) 15

Paragraph For Questions 12 and 13

Let a, r, s, t be nonzero real numbers. Let $P(at^2, 2at)$, $Q, R(ar^2, 2ar)$ and $S(as^2, 2as)$ be distinct points on the parabola $y^2 = 4ax$. Suppose that PQ is the focal chord and lines QR and PK are parallel, where K is the point $(2a, 0)$.

12. The value of r is- **[JEE(Advanced) 2014]**
- (A) $-\frac{1}{t}$ (B) $\frac{t^2+1}{t}$ (C) $\frac{1}{t}$ (D) $\frac{t^2-1}{t}$
13. If $st = 1$, then the tangent at P and the normal at S to the parabola meet at a point whose ordinate is- **[JEE(Advanced) 2014]**
- (A) $\frac{(t^2+1)^2}{2t^3}$ (B) $\frac{a(t^2+1)^2}{2t^3}$ (C) $\frac{a(t^2+1)^2}{t^3}$ (D) $\frac{a(t^2+2)^2}{t^3}$

SOLUTIONS


1. **Ans. (A)**

Sol. Let point P ($at^2, 2at$)

normal at P is $y = -tx + 2at + at^3$

$y = 0, x = 2a + at^2$

$Q(2a + at^2, 0)$



$$\text{Area of } \Delta PFQ = \left| \frac{1}{2} (a + at^2)(2at) \right| = 120$$


$$\therefore m = -t$$

$$\therefore a^2 [1 + m^2] m = 120$$

(a, m) = (2, 3) will satisfy

2. **Ans. (B, C, D)**

Sol. Let equation of tangent with slope 'm' be



$$T : y = mx + \frac{1}{m}$$

T : passes through (-2, 1) so

$$1 = -2m + \frac{1}{m}$$

$$\Rightarrow m = -1 \text{ or } m = \frac{1}{2}$$

Points are given by $\left(\frac{a}{m^2}, \frac{2a}{m} \right)$

So, one point will be (1, -2) & (4, 4)

Let $P_1(4, 4)$ & $P_2(1, -2)$

$P_1S : 4x - 3y - 4 = 0$

$P_2S : x - 1 = 0$

$$PQ_1 = \left| \frac{4(-2) - 3(1) - 4}{5} \right| = 3$$


$SP = \sqrt{10}$; $PQ_2 = 3$; $SQ_1 = 1 = SQ_2$

$$\frac{1}{2} \left(\frac{Q_1Q_2}{2} \right) \times \sqrt{10} = \frac{1}{2} \times 3 \times 1$$

(comparing Areas)

$$\Rightarrow Q_1Q_2 = \frac{2 \times 3}{\sqrt{10}} = \frac{3\sqrt{10}}{5}$$

3. **Ans. (A, B, D)**



Sol.


Note that P lies on directrix so triangle PQQ' is right angled, hence QQ' passes through focus F.

$$PF = 4\sqrt{2}$$

Equation of QF is $y = x - 2$ & PF is $x + y = 2$

Hence A, B, D.

4. **Ans. (1.50)**



Sol.

Let the circle be

$$x^2 + y^2 + \lambda x = 0$$

For point of intersection of circle & parabola

$$y^2 = 4 - x.$$

$$x^2 + 4 - x + \lambda x = 0 \Rightarrow x^2 + x(\lambda - 1) + 4 = 0$$

$$\text{For tangency : } \Delta = 0 \Rightarrow (\lambda - 1)^2 - 16 = 0$$

$$\Rightarrow \lambda = 5 \text{ (rejected) or } \lambda = -3$$

$$\text{Circle : } x^2 + y^2 - 3x = 0$$

$$\text{Radius} = \frac{3}{2} = 1.5$$

5. Ans. (2.00)

Sol. For point of intersection :
 $x^2 - 4x + 4 = 0 \Rightarrow x = 2$ so $\alpha = 2$

6. Ans. (D)

Sol. Equation of chord with mid point (h, k) :

$$k \cdot y - 16 \left(\frac{x+h}{2} \right) = k^2 - 16h$$


$$\Rightarrow 8x - ky + k^2 - 8h = 0$$

Comparing with $2x + y - p = 0$, we get

$$k = -4; 2h - p = 4$$

only (D) satisfies above relation.

7. Ans. (A, C, D)



$$y^2 = 4x$$

point P lies on normal to parabola passing through centre of circle

$$y + tx = 2t + t^3 \quad \dots(i)$$

$$8 + 2t = 2t + t^3$$

$$t = 2$$

$$P(4, 4)$$

$$SP = \sqrt{(4-2)^2 + (4-8)^2}$$

$$SP = 2\sqrt{5}$$

$$SQ = 2$$

$$\Rightarrow PQ = 2\sqrt{5} - 2$$

$$\frac{SQ}{QP} = \frac{1}{\sqrt{5}-1} = \frac{\sqrt{5}+1}{4}$$

To find x intercept

put $y = 0$ in (i)


$$\Rightarrow x = 2 + t^2$$

$$x = 6$$

$$\therefore \text{Slope of common normal} = -t = -2$$

$$\therefore \text{Slope of tangent} = \frac{1}{2}$$

8. Ans. (2)



Sol.

The co-ordinates of latus rectum are (1,2) and (1,-2)

clearly slope of tangent is given by $\frac{dy}{dx} = \frac{2}{y}$

\therefore At $y = 2$ slope of normal = -1

and At $y = -2$ slope of normal = 1

\therefore Equation of normal at (1,2)

$$(y - 2) = -1(x - 1) \Rightarrow x + y = 3$$

Now, this line is tangent to circle

$$(x - 3)^2 + (y + 2)^2 = r^2$$

\therefore perpendicular distance from centre to line = Radius of circle

$$\therefore \frac{|3 - 2 - 3|}{\sqrt{2}} = r \Rightarrow r^2 = 2$$

9. Ans. (4)

Sol. Let there be a point $(t^2, 2t)$ on $y^2 = 4x$

Clearly its reflection in $x + y + 4 = 0$ is given by

$$\frac{x - t^2}{1} = \frac{y - 2t}{1} = \frac{-2(t^2 + 2t + 4)}{2}$$


$$\therefore x = -(2t + 4) \text{ \& } y = -(t^2 + 4)$$

$$\text{Now, } y = -5 \Rightarrow t = \pm 1$$

$$\therefore x = -6 \text{ or } x = -2$$

$$\therefore \text{Distance between A \& B} = 4$$

10. Ans. (A, D)



Sol.

$$\because \angle POQ = \frac{\pi}{2} \quad \Rightarrow \quad t_1 t_2 = -4$$

$$\therefore \begin{vmatrix} 1 & t_1^2 & t_1 & 1 \\ 2 & \frac{t_1^2}{2} & t_1 & 1 \\ 1 & t_2^2 & t_2 & 1 \\ 2 & \frac{t_2^2}{2} & t_2 & 1 \end{vmatrix} = 3\sqrt{2}$$

$$\Rightarrow \left| \frac{t_1^2 t_2 - t_1 t_2^2}{2} \right| = 6\sqrt{2}$$

$$\Rightarrow |t_1 - t_2| = 3\sqrt{2}$$


$$\Rightarrow t_1 + \frac{4}{t_1} = 3\sqrt{2} \quad (\because t_1 > 0)$$

We get $t_1 = 2\sqrt{2}, \sqrt{2}$

$P(4, 2\sqrt{2})$ or $(1, \sqrt{2})$

11. Ans. (D)

Sol.



$$y = mx + \frac{2}{m}$$

$$\left| \frac{0 - 0 + \frac{2}{m}}{\sqrt{1 + m^2}} \right| = \sqrt{2} \Rightarrow 2 = m^2(1 + m^2)$$

$$\Rightarrow m = \pm 1$$

$$TP : -x + y = 2$$

So $P(-1, 1)$ & $Q(-1, -1)$

& $R\left(\frac{2}{m}, \frac{4}{m}\right) \equiv R(2, 4)$ & $S(2, -4)$

$$\text{So } \Delta = \frac{1}{2} \cdot 10.3 = 15$$


So $P(-1, 1)$ & $Q(-1, -1)$

& $R\left(\frac{2}{m}, \frac{4}{m}\right) \equiv R(2, 4)$ & $S(2, -4)$

$$\text{So } \Delta = \frac{1}{2} \cdot 10.3 = 15$$

12. Ans. (D)

Sol.



\because PQ is a focal chord

\therefore co-ordinates of point Q are $\left(\frac{a}{t^2}, -\frac{2a}{t}\right)$

$$m_{QR} = \frac{2a\left(r + \frac{1}{t}\right)}{a\left(r^2 - \frac{1}{t^2}\right)} = \frac{2}{\left(r - \frac{1}{t}\right)}$$

$$m_{PK} = \frac{2at - 0}{a(t^2 - 2)} = \frac{2t}{t^2 - 2}$$

Given $m_{QR} = m_{PK}$

$$\Rightarrow \frac{2}{r - \frac{1}{t}} = \frac{2t}{t^2 - 2} \Rightarrow r = \frac{t^2 - 2}{t} + \frac{1}{t}$$

$$\Rightarrow r = t - \frac{2}{t} + \frac{1}{t} \Rightarrow r = \frac{t^2 - 1}{t}$$

13. Ans. (B)

Sol. Equation of tangent at point P is

$$ty = x + at^2 \quad \dots(i)$$

Equation of normal at point S is

$$\frac{1}{t}x + y = \frac{2a}{t} + \frac{a}{t^3}$$

$$\Rightarrow t^2x + t^3y = 2at^2 + a \quad \dots(ii)$$

Multiply equation (i) by t^2 and then subtract from equation (ii),

we get,

$$2t^3y = 2at^2 + at^4 + a$$

$$\Rightarrow 2t^3y = a(1 + t^4 + 2t^2)$$

$$\Rightarrow y = \frac{a(1 + t^2)^2}{2t^3}$$