

HYPERBOLA

1. Consider the hyperbola

$$\frac{x^2}{100} - \frac{y^2}{64} = 1$$

with foci at S and S₁, where S lies on the positive x-axis. Let P be a point on the hyperbola, in the first quadrant. Let $\angle SPS_1 = \alpha$, with $\alpha < \frac{\pi}{2}$. The straight line passing through the point S and having the same slope as that of the tangent at P to the hyperbola, intersects the straight line S₁P at P₁. Let δ be the distance of P from the straight line SP₁, and $\beta = S_1P$. Then the greatest integer less than or equal to $\frac{\beta\delta}{9} \sin \frac{\alpha}{2}$ is _____.

[JEE(Advanced) 2022]

2. Let a and b be positive real numbers such that $a > 1$ and $b < a$. Let P be a point in the first quadrant that lies on the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$. Suppose the tangent to the hyperbola at P passes through the point (1,0), and suppose the normal to the hyperbola at P cuts off equal intercepts on the coordinate axes. Let Δ denote the area of the triangle formed by the tangent at P, the normal at P and the x-axis. If e denotes the eccentricity of the hyperbola, then which of the following statements is/are TRUE ?

[JEE(Advanced) 2020]

- (A) $1 < e < \sqrt{2}$ (B) $\sqrt{2} < e < 2$ (C) $\Delta = a^4$ (D) $\Delta = b^4$

3. Let H : $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$, where $a > b > 0$, be a hyperbola in the xy-plane whose conjugate axis LM subtends an angle of 60° at one of its vertices N. Let the area of the triangle LMN be $4\sqrt{3}$.

[JEE(Advanced) 2018]

LIST-I

- P. The length of the conjugate axis of H is
 Q. The eccentricity of H is
 R. The distance between the foci of H is
 S. The length of the latus rectum of H is

LIST-II

1. 8
 2. $\frac{4}{\sqrt{3}}$
 3. $\frac{2}{\sqrt{3}}$
 4. 4

The correct option is :

- (A) P → 4; Q → 2, R → 1; S → 3
 (B) P → 4; Q → 3; R → 1; S → 2
 (C) P → 4; Q → 1, R → 3; S → 2
 (D) P → 3; Q → 4; R → 2; S → 1

4. If $2x - y + 1 = 0$ is tangent to the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{16} = 1$, then which of the following CANNOT be sides of a right angled triangle ?

[JEE(Advanced) 2017]

- (A) 2a, 4, 1 (B) 2a, 8, 1 (C) a, 4, 1 (D) a, 4, 2

MATCHING TYPE : (Q.5 to Q.7)

Column 1, 2 and 3 contain conics, equation of tangents to the conics and points of contact, respectively.

Column-1

Column-2

Column-3

(I) $x^2 + y^2 = a^2$

(i) $my = m^2x + a$

(P) $\left(\frac{a}{m^2}, \frac{2a}{m}\right)$

(II) $x^2 + a^2y^2 = a^2$

(ii) $y = mx + a\sqrt{m^2 + 1}$

(Q) $\left(\frac{-ma}{\sqrt{m^2 + 1}}, \frac{a}{\sqrt{m^2 + 1}}\right)$

(III) $y^2 = 4ax$

(iii) $y = mx + \sqrt{a^2m^2 - 1}$

(R) $\left(\frac{-a^2m}{\sqrt{a^2m^2 + 1}}, \frac{1}{\sqrt{a^2m^2 + 1}}\right)$

(IV) $x^2 - a^2y^2 = a^2$

(iv) $y = mx + \sqrt{a^2m^2 + 1}$

(S) $\left(\frac{-a^2m}{\sqrt{a^2m^2 - 1}}, \frac{-1}{\sqrt{a^2m^2 - 1}}\right)$

5. The tangent to a suitable conic (Column 1) at $\left(\sqrt{3}, \frac{1}{2}\right)$ is found to be $\sqrt{3}x + 2y = 4$, then which of the following options is the only **CORRECT** combination? **[JEE(Advanced) 2017]**

- (A) (II) (iii) (R) (B) (IV) (iv) (S) (C) (IV) (iii) (S) (D) (II) (iv) (R)

6. If a tangent to a suitable conic (Column 1) is found to be $y = x + 8$ and its point of contact is $(8,16)$, then which of the following options is the only **CORRECT** combination? **[JEE(Advanced) 2017]**

- (A) (III) (i) (P) (B) (III) (ii) (Q)
(C) (II) (iv) (R) (D) (I) (ii) (Q)

7. For $a = \sqrt{2}$, if a tangent is drawn to a suitable conic (Column 1) at the point of contact $(-1,1)$, then which of the following options is the only **CORRECT** combination for obtaining its equation? **[JEE(Advanced) 2017]**

- (A) (II) (ii) (Q) (B) (III) (i) (P)
(C) (I) (i) (P) (D) (I) (ii) (Q)


8. Consider the hyperbola $H : x^2 - y^2 = 1$ and a circle S with center $N(x_2, 0)$. Suppose that H and S touch each other at a point $P(x_1, y_1)$ with $x_1 > 1$ and $y_1 > 0$. The common tangent to H and S at P intersects the x -axis at point M . If (l, m) is the centroid of the triangle ΔPMN , then the correct expression(s) is(are) **[JEE(Advanced) 2015]**

- (A) $\frac{dl}{dx_1} = 1 - \frac{1}{3x_1^2}$ for $x_1 > 1$ (B) $\frac{dm}{dx_1} = \frac{x_1}{3(\sqrt{x_1^2 - 1})}$ for $x_1 > 1$
(C) $\frac{dl}{dx_1} = 1 + \frac{1}{3x_1^2}$ for $x_1 > 1$ (D) $\frac{dm}{dy_1} = \frac{1}{3}$ for $y_1 > 0$

SOLUTIONS

1. Ans. (7)

Sol.



$$S_1P - SP = 20$$

$$\beta - \frac{\delta}{\sin \frac{\alpha}{2}} = 20$$

$$\beta^2 + \frac{\delta^2}{\sin^2 \frac{\alpha}{2}} - 400 = \frac{2\beta\delta}{\sin \frac{\alpha}{2}}$$

$$\frac{1}{SP} = \frac{\sin \frac{\alpha}{2}}{\delta}$$

$$\cos \alpha = \frac{SP^2 + \beta^2 - 656}{2\beta \frac{\delta}{\sin \frac{\alpha}{2}}}$$

$$= \frac{\frac{2\beta\delta}{\sin \frac{\alpha}{2}} - 256}{2\beta S} = \cos \alpha$$

$$\frac{\lambda - 128}{\lambda} = \cos \alpha$$

$$\lambda(1 - \cos \alpha) = 128$$


$$\frac{\beta\delta}{\sin \frac{\alpha}{2}} \cdot 2 \sin^2 \frac{\alpha}{2} = 128$$

$$\frac{\beta\delta}{9} \sin \frac{\alpha}{2} = \frac{64}{9} \Rightarrow \left[\frac{\beta\delta}{9} \sin \frac{\alpha}{2} \right] = 7$$

where [.] denotes greatest integer function

2. Ans. (A, D)

Sol.



Since Normal at point P makes equal intercept on co-ordinate axes, therefore slope of Normal = -1

Hence slope of tangent = 1

Equation of tangent

$$y - 0 = 1(x - 1)$$

$$y = x - 1$$

Equation of tangent at (x_1, y_1)

$$\frac{xx_1}{a^2} - \frac{yy_1}{b^2} = 1$$

$$x - y = 1 \text{ (equation of Tangent)}$$

on comparing $x_1 = a^2, y_1 = b^2$

$$\text{Also } a^2 - b^2 = 1 \quad \dots(1)$$

Now equation of normal at $(x_1, y_1) = (a^2, b^2)$

$$y - b^2 = -1(x - a^2)$$

$$x + y = a^2 + b^2 \quad \dots(\text{Normal})$$

point of intersection with x-axis is $(a^2 + b^2)$

$$\text{Now } e = \sqrt{1 + \frac{b^2}{a^2}}$$

$$e = \sqrt{1 + \frac{b^2}{b^2 + 1}} \quad \left[\text{from (1)} \frac{b^2}{b^2 + 1} < 1 \right]$$

$$1 < e < \sqrt{2} \quad \text{option (A)}$$


$$\Delta = \frac{1}{2} \cdot AB \cdot PQ$$

$$\text{and } \Delta = \frac{1}{2} (a^2 + b^2 - 1) \cdot b^2$$

$$\Delta = \frac{1}{2} (2b^2) b^2 \quad (\text{from (1)} \quad a^2 - 1 = b^2)$$

$$\Delta = b^4 \quad \text{so option (D)}$$

3. **Ans. (B)**



$$\tan 30^\circ = \frac{b}{a}$$

$$\Rightarrow a = b\sqrt{3}$$

$$\text{Now area of } \triangle LMN = \frac{1}{2} \cdot 2b \cdot b\sqrt{3}$$

$$4\sqrt{3} = \sqrt{3}b^2$$

$$\Rightarrow b = 2 \text{ \& } a = 2\sqrt{3}$$

$$\Rightarrow e = \sqrt{1 + \frac{b^2}{a^2}} = \frac{2}{\sqrt{3}}$$

P. Length of conjugate axis = $2b = 4$

So P \rightarrow 4

Q. Eccentricity $e = \frac{2}{\sqrt{3}}$

So Q \rightarrow 3

R. Distance between foci = $2ae$

$$= 2(2\sqrt{3})\left(\frac{2}{\sqrt{3}}\right) = 8$$

So R \rightarrow 1

S. Length of latus rectum =

$$\frac{2b^2}{a} = \frac{2(2)^2}{2\sqrt{3}} = \frac{4}{\sqrt{3}}$$

So S \rightarrow 2

4. **Ans. (B, C, D)**

Sol. The line $y = mx + c$ is tangent to hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1, \text{ if } c^2 = a^2m^2 - b^2$$

$$\therefore (1)^2 = 4a^2 - 16 \Rightarrow a^2 = \frac{17}{4}$$

$$\Rightarrow a = \frac{\sqrt{17}}{2}$$

For option (A), sides are $\sqrt{17}, 4, 1$

(\Rightarrow Right angled triangle)

For option (B), sides are $\sqrt{17}, 8, 1$

(\Rightarrow Triangle is not possible)

For option (C), sides are $\frac{\sqrt{17}}{2}, 4, 1$

(\Rightarrow Triangle is not possible)

For option (D), sides are $\frac{\sqrt{17}}{2}, 4, 2$

(\Rightarrow Triangle exist but not right angled)

5. **Ans. (D)**

Sol. $P\left(\sqrt{3}, \frac{1}{2}\right)$; tangent $\sqrt{3}x + 2y = 4$

$$\Rightarrow (\sqrt{3})x + 4\left(\frac{1}{2}\right)y = 4 \text{ comparing with (II)}$$

$$\Rightarrow a = 2 \therefore y = mx + \sqrt{a^2m^2 + 1} \text{ is tangent}$$

$$\text{for } m = -\frac{\sqrt{3}}{2} \text{ i.e (ii)}$$

$$\therefore \text{ point of contact for } a = 2, m = -\frac{\sqrt{3}}{2} \text{ is R}$$

6. **Ans. (A)**

Sol. $y = x + 8$ is tangent $\Rightarrow m = 1$; $P(8, 16)$

Comparing tangent with (i) of column 2, $m = 1$ satisfied and $a = 8$ obtained which matches for point of contact (P) of column 3 and (III) of column 1.


7. **Ans. (D)**

Sol. For $a = \sqrt{2}$ and point $(-1, 1)$ only I of column-1 satisfies. Hence equation of tangent is $-x + y = 2$ or $y = x + 2 \Rightarrow m = 1$ which matches with (ii) of column-2 and also with Q of column-3

$$\text{Let } f(x) = x + \log_e x - x \log_e x, x \in (0, \infty).$$

8. Ans. (A, B, D)

Sol. Given H : $x^2 - y^2 = 1$



Now, equation of family of circle touching hyperbola at (x_1, y_1) is

$$(x - x_1)^2 + (y - y_1)^2 + \lambda(xx_1 - yy_1 - 1) = 0$$

Now, its centre is $(x_2, 0)$

$$\therefore 2y_1 + \lambda y_1 = 0 \quad \Rightarrow \quad \lambda = -2$$

$$\therefore x_2 = \frac{2x_1 - \lambda x_1}{2} = \frac{4x_1}{2} = 2x_1$$

$$\therefore P \equiv (x_1, \sqrt{x_1^2 - 1})$$

$$N \equiv (2x_1, 0)$$

$$\& \quad M \equiv \left(\frac{1}{x_1}, 0 \right)$$

$$\therefore \ell = \frac{3x_1 + \frac{1}{x_1}}{3} = x_1 + \frac{1}{3x_1}$$

$$\Rightarrow \frac{d\ell}{dx_1} = 1 - \frac{1}{3x_1^2} \quad x_1 > 1$$

$$m = \frac{\sqrt{x_1^2 - 1}}{3} \quad \Rightarrow \quad \frac{dm}{dx_1} = \frac{x_1}{3\sqrt{x_1^2 - 1}}$$

$$\text{Also } m = \frac{y_1}{3} \quad \Rightarrow \quad \frac{dm}{dy_1} = \frac{1}{3} \quad y_1 > 0$$