

PHYSICAL CHEMISTRY





REAL GAS

1. A gas has a compressibility factor of 0.5 and a molar volume of  $0.4 \text{ dm}^3 \text{ mol}^{-1}$  at a temperature of 800 K and pressure  $x \text{ atm}$ . If it shows ideal gas behaviour at the same temperature and pressure, the molar volume will be  $y \text{ dm}^3 \text{ mol}^{-1}$ . The value of  $x/y$  is \_\_\_\_\_. [JEE(Advanced) 2023]


[Use: Gas constant,  $R = 8 \times 10^{-2} \text{ L atm K}^{-1} \text{ mol}^{-1}$ ]

2. One mole of a monoatomic real gas satisfied the equation  $p(V-b) = RT$  where  $b$  is a constant. The relationship of interatomic potential  $V(r)$  and interatomic distance  $r$  for the gas is given by –

[JEE(Advanced) 2015]



3. For one mole of a van der Waals gas when  $b = 0$  and  $T = 300 \text{ K}$ , the  $PV$  vs.  $1/V$  plot is shown below. The value of the van der Waals constant  $a$  ( $\text{atm. liter}^2 \text{ mol}^{-2}$ ) is [IIT-JEE 2012]



- (A) 1.0                      (B) 4.5                      (C) 1.5                      (D) 3.0

SOLUTIONS

1. Ans. (100)

Sol. For gas :  $Z = 0.5$ ,  $V_m = 0.4 \text{ L/mol}$

$$T = 800 \text{ K}, P = X \text{ atm.}$$

$$\Rightarrow Z = \frac{PV_m}{RT}$$

$$\Rightarrow \frac{X(0.4)}{0.08 \times 800} = 0.5$$

$$\Rightarrow X = 80$$

For ideal gas,  $PV_m = RT$

$$\Rightarrow V_m = \frac{RT}{P} = \frac{0.08 \times 800}{80} = 0.8 \text{ L mol}^{-1} = y$$

$$\text{Then, } \frac{x}{y} = \frac{80}{0.8} = 100$$

2. Ans. (C)


Sol.  $P(V-b) = RT$

$$\therefore a = 0$$

Since only repulsive forces are present.

Repulsive forces contribute only at very close distance.

So potential energy increases abruptly.



3. Ans. (C)

Sol.  $\left(P + \frac{a}{V^2}\right)(V-0) = RT$  as  $(n = 1)$

$$PV = RT - \frac{a}{V}$$

On comparing slope from graph

$$-a = \frac{20.1 - 21.6}{3 - 2}$$

$$-a = \frac{-1.5}{1} = a = 1.5$$