






SOLUTIONS
Biology (Subject Code-044)
(PAPER CODE : 57/4/1)

Q.No.	EXPECTED ANSWER / VALUE POINTS	Marks	Total Marks
SECTION – A			
1.	(C) / 1300	1	1
2.	(A) / Secretory phase	1	1
3.	(C) / IUT	1	1
4.	(B) / form one linkage group	1	1
5.	(C) / (ii) and (iv)	1	1
6.	(B) / <i>Australopithecus</i>	1	1
7.	(D) / (i) and (iii)	1	1
8.	(B) / <i>Bacillus thuringiensis</i>	1	1
9.	(D) / Skull A is of an adult chimpanzee and skull B is of human	1	1
10.	(C) / dsRNA	1	1
11.	(D) / Competition	1	1
12.	(C) / Corn borer	1	1
13.	(A) / Both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of the Assertion (A)	1	1
14.	(B) / Both Assertion (A) and Reason (R) are true but Reason (R) is not the correct explanation of the Assertion(A)	1	1
15.	(A) / Both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of the Assertion (A)	1	1
16.	(A) / Both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of the Assertion (A)	1	1
SECTION – B			
17.	- Implants work by inhibiting ovulation / inhibiting implantation /changing the quality of cervical mucus to retard entry of sperms. - The effective periods are much longer than those of pills. (Any other correct advantage)	1 1	2

	(a)  (b) The RNA transcribed as 5 'G A U A C G U A C G U A 3'	$\frac{1}{2} \times 4$ 1	3
24.	- Stabilising – more individuals acquire the mean character value i.e variation is much reduced.  - Directional – more individuals acquire value other than the mean character value.  - Disruptive – more individuals acquire peripheral character value at both ends of the distribution curve two peaks are formed. 	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$	

//



1/2x6

3

25.

- (a) The cancer causing viruses called oncogenic viruses have genes called viral oncogenes which transform normal cell into cancerous cells.
- (b)
- When the cells of a benign tumour acquire the property of metastasis under certain conditions.
 - These cells damage the surrounding normal cells of the body / They compete with the normal cells and starve them of vital nutrients.

1

1

1


3

26.	(a) Proinsulin contains an extra stretch called the C-peptide in addition to chain A and B / C-peptide is not present in the (mature) insulin. (b) Steps to produce insulin through rDNA technology. - Eli Lilly an American Company prepared two DNA sequences corresponding to A and B chain of human insulin. - These chains A and B were then introduced in plasmid <i>E.coli</i> to produce multiple copies. - These chains were extracted of separately. - Chain A and B were combined by creating disulphide bonds to form human insulin.	1 ½x4	3
27.	(a) (i) • Curve (a) – The food and space are unlimited. • Curve (b)- The food and space are limited. (ii) Curve ‘a’ (iii) It represents the carrying capacity which means the maximum number of individuals that a given environment can support and beyond which further growth is not possible. OR (b) (i) Predation is the interspecific interaction in which a large animal called predator kills and consumes a smaller animals called prey. (ii) - It keeps the prey population under control. - It helps in maintaining species diversity in a community by reducing the intensity of competition among competing prey species. - It acts as conduit for energy transfer across trophic level. - Biological control methods adopted in agricultural pest control are based on the ability of the predator to regulate prey population. (any two) (iii) If a predator is too efficient and over exploits its prey then the prey might become extinct and following it the predator will also become extinct for lack of food.	½ ½ 1 1 1 ½ x2 1	3
28.	(a) Ex-situ conservation. (b) -Cryopreservation – (–196°C) sperms or eggs or animal tissues or gametes and embryos of threatened species can be stored . -In vitro using tissue culture methods plants can be propagated	½ 1+1	

	-Seed banks in which seeds of different genetic strains of commercially important plants can be kept. - Botanical Gardens in which threatened plants can be conserved in special settings. -Any other correct option with explanation. (Any two)		
	(c) Botanical Gardens , Wildlife safari parks (Any one)	$\frac{1}{2}$	3
	SECTION - D		
29.	(a) It is necessary for the entry of lactose into the cell . (b)(i) In the presence of lactose the repressor is inactivated by the interaction with the inducer that allow RNA polymerase access to promoter and transcription proceeds. OR (b)(ii) It is called negative regulation because the operon is switched off in the absence of inducer and transcription is prevented. (c) - In the absence of the inducer the repressor binds to the operator and prevents transcription of the structural gene. - When inducer or lactose is present it prevents the repressor from binding to the operator hence regulation is at the transcriptional level.	1 1 1 1	4
30.	(a) B – lymphocytes /B cells , and T – lymphocytes/ T Cells (b) (i) Physical barrier , physiological barriers , Cellular barriers , Cytokine barriers (any two) OR (b) (ii) Our immune system has the ability to distinguish between ‘self’ and ‘non-self’ cells or tissues.	$\frac{1}{2}+\frac{1}{2}$ $\frac{1}{2}+\frac{1}{2}$ 1	

	(c) <table> <tr> <th>Active Immunity</th> <th>Passive Immunity</th> </tr> <tr> <td>When antibodies are produced in our body in response to an antigen then it is called active immunity.</td> <td>When preformed antibodies are injected into the body for protection then it is called passive immunity.</td> </tr> <tr> <td>It is slow and takes time to give its full effective response.</td> <td>It gives quick response.</td> </tr> <tr> <td>It lasts for longer period.</td> <td>It last for shorter period.</td> </tr> <tr> <td>Memory cells are formed.</td> <td>Memory cells are not formed.</td> </tr> <tr> <td colspan="2">(Any two correct difference)</td> </tr> </table>	Active Immunity	Passive Immunity	When antibodies are produced in our body in response to an antigen then it is called active immunity.	When preformed antibodies are injected into the body for protection then it is called passive immunity.	It is slow and takes time to give its full effective response.	It gives quick response.	It lasts for longer period.	It last for shorter period.	Memory cells are formed.	Memory cells are not formed.	(Any two correct difference)		1+1	4
Active Immunity	Passive Immunity														
When antibodies are produced in our body in response to an antigen then it is called active immunity.	When preformed antibodies are injected into the body for protection then it is called passive immunity.														
It is slow and takes time to give its full effective response.	It gives quick response.														
It lasts for longer period.	It last for shorter period.														
Memory cells are formed.	Memory cells are not formed.														
(Any two correct difference)															
	SECTION - E														
31.	(a) - Gonadotropin releasing hormone/ GnRH , It stimulates the pituitary gland to release follicle stimulating hormone or FSH and Luteinising hormone or LH both act on ovary. - FSH/ Follicle stimulating hormone , stimulate follicular development and secretion of estrogen by the follicle cells. - LH/ Luteinising hormone , stimulates ovulation/ formation of corpus luteum from the ruptured graafian follicle. - Progesterone , Maintenance of endometrium necessary for implantation and other events of pregnancy. - Estrogen , responsible for the maturation of primary follicle into graafian follicle/ regeneration of endometrium lining. OR	½+½ ½+½ ½+½ ½+½ ½+½													

(b)(i)



$\frac{1}{2} \times 6$

($\frac{1}{2}$ mark for any 6 correct labelling)

(b)(ii)

•

Perisperm	Endosperm
In seeds residual persistent nucellus is called perisperm.	It is the product formed from the primary endosperm cell in the ovule.
The cells are diploid (2n).	Cells are triploid (3n).
(any one)	

1

• Examples :

Perisperm	Endosperm
Eg. Black pepper / beet/ any other example	eg. Castor / Wheat/ Maize/Barley/ Sunflower/ Coconut/any other example

$\frac{1}{2} + \frac{1}{2}$

5

32.	(a) - Types and location of gene: –α- Thalassemia is under the control of two genes HBA1 and HBA2 , both located on chromosome-16 –β- Thalassemia is the control of a single gene HBB , present on chromosomes -11 - Causes: It is due to mutation / deletion of one or more of the four alleles of the genes controlling the disease. - Symptoms: Anaemia/ extreme tiredness / pale skin/ Breathlessness/ any other correct symptom <table><tr><th>Thalassemia</th><th>Sickle cell anaemia</th></tr><tr><td>It is quantitative where less amount of haemoglobin is synthesised.</td><td>It is qualitative where defective haemoglobin is synthesised.</td></tr><tr><td>Both α and β chain are affected</td><td>Only β chain is affected (Any one)</td></tr></table>	Thalassemia	Sickle cell anaemia	It is quantitative where less amount of haemoglobin is synthesised.	It is qualitative where defective haemoglobin is synthesised.	Both α and β chain are affected	Only β chain is affected (Any one)	$\frac{1}{2}+\frac{1}{2}$ $\frac{1}{2}+\frac{1}{2}$ 1 1 1	
Thalassemia	Sickle cell anaemia								
It is quantitative where less amount of haemoglobin is synthesised.	It is qualitative where defective haemoglobin is synthesised.								
Both α and β chain are affected	Only β chain is affected (Any one)								

	OR (b) (i) Major goals of HGP: - Identify all the genes (20,000-25,000 in human DNA) - Determine the sequence of the three billion base pairs present in human DNA. - Store the information in database. - Improve the tools for data analysis. - Transfer the technologies to other sectors (like industries). - Address the ethical legal and social issues that may arise from this project. (any four) (ii) Methodologies : - Expressed Sequence Tags (ESTs) , It focuses on identifying all the genes that are expressed as RNAs. - Sequence Annotation , it aims at sequencing the whole set of genome that includes all coding sequences and non coding sequences and then assigning functions to different regions in the sequence. - (iii) • Yeast artificial chromosomes • It is used as vector.	$\frac{1}{2} \times 4$ $\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$	5
--	---	--	-----------------

33.	(a) (i) (I) Naming of restriction enzymes (EcoRI): -First letter 'E' of the name comes from genes (<i>Escherichia</i>) - The second two letters 'co' come from the species (<i>coli</i>) -The letter 'R' derived from the name of the strain - Roman number (1) indicate the order in which the enzymes were isolated from the strain of the bacteria. (II) The restriction enzyme EcoRI recognised the palindromic sequence 5' – G↓AATTC – 3' 3' – CTAA↑G – 5' (a) (ii) • These overhanging stretches are called 'Sticky ends' • They form hydrogen bond with their complementary cut counterpart that facilitates the action of enzyme DNA ligase. OR (b) (i) - When the alien DNA is inserted within the coding sequence of the enzyme β - galactosidase it results in inactivation of the enzyme which is referred as insertional inactivation. - The presence of chromogenic substrate gives blue coloured colonies if the plasmid in the bacteria does not have insert or is non recombinant. - Colonies which do not produce any colour are identified as recombinants as their gene for enzyme is inactivated due to an insert. (ii) <i>Agrobacterium tumefaciens</i>. (iii) When this <i>Agrobacterium</i> infects a plant cell it delivers 'T'-DNA or Ti tumor-inducing plasmid into the plant cell and transforms it into a tumor cell.	½ ½ ½ ½ 1 1 1 1 1 1 1 1 1	5
-----	--	--	---