

[SINGLE CORRECT CHOICE TYPE]

Q.1 A ray of light passes from a medium of refractive index μ into air. The angle of incidence is found to be half the angle of refraction. What is the angle of refraction? [3]

- (A) $\cos^{-1}(\mu/2)$ (B) $\sin^{-1}(\mu/2)$ (C) $2 \sin^{-1}(\mu/2)$ (D) $2 \cos^{-1}(\mu/2)$

Q.2 Consider the point P as the origin and line OP as the x-axis in the situation shown in the figure. Which of the following represents the coordinates of the image of the point object O. (Take $OP = 20$ cm) [3]

- (A) (20 cm, 0 cm) (B) (10 cm, [] cm)
 (C) (-10 cm, [] cm) (D) (10 cm, -[] cm)

Q.3 A thin convex lens is cut along its principal axis and one half of the lens is placed as shown in the figure. What are the x and y coordinates respectively, of the image w.r.t. the origin O of the given coordinate axis? The focal length of the lens is 20 cm. [3]





- (A) (60 cm, 12 mm) (B) (60 cm, 18 mm)
 (C) (12 cm, 2.4 mm) (D) (60 cm, -12 mm)

Q.4 An equi convex lens whose focal length in air is 20 cm is made up of a material whose refractive index is 1.5. If one of its surface is silvered and then dipped into a transparent liquid whose refractive index is 1.6. In the liquid it will behave like a [3]

- (A) converging lens (B) diverging lens (C) converging mirror (D) diverging mirror

Q.5 Refractive index of a transparent slab varies as $\mu = kx$ where x is the distance from origin. Time taken by the light to travel the slab of thickness t (as shown in the figure) [3]




Q.6 A ray moving along unit vector [] strikes a reflecting surface in xy plane. The unit vector along the reflected ray would be [3]

- (A) [] (B) [] (C) []

Q.7 The lens shown in the diagram has a refractive index higher than that of its surrounding medium. It will converge a paraxial beam incident on it [3]

- (A) from left only (B) from right only
 (C) either from left or right (D) None.



Q.8 A paraxial beam of light having diameter 4 mm is incident on a convex lens having focal length 10 cm. A screen is placed normal to the beam at a distance of 30cm on the other side. Radius of image formed on the screen is- [3]

- (A) 6 mm (B) 4 mm (C) 3mm (D) 2 mm

Q.9 A ray of light from air is incident at an angle of 70° into a medium having refractive index μ . The reflected and the refracted rays are found to suffer equal deviations in opposite direction. μ equals [3]

- (A) $\tan 70^\circ$ (B) $2\sin 70^\circ$ (C) $\operatorname{cosec} 70^\circ$ (D) None of these


Q.10 In the figure shown a ray of light hits the interface of the media I & II at angle of 30° . Refractive indices of the media are shown in the figure. The angle made by the ray with normal in medium three is [3]

(A) $\sin^{-1}\left(\frac{2}{3}\right)$

(B) $\sin^{-1}\left(\frac{\sqrt{2}}{3}\right)$


(C) ray will not come in medium III

(D) $i_3 = \sin^{-1}\left(\frac{2}{7}\right)$




Q.11 A thin prism with angle 4° and refractive index 1.5 is placed inside a thin transparent tube with water (refractive index, $4/3$) as shown. The deviation of light will be [3]

- (A) 2.67° upward (B) 0.67° downward
(C) 2.67° downward (D) 0.5° downward



Q.12 As shown in the figure particle at M vibrates simple harmonically about M with amplitude 0.1cm perpendicular to axis along a line. Find the amplitude of vibration of image [3]

- (A) 0.3 cm in same phase
(B) 0.1 cm in opposite phase
(C) 0.2 cm in same phase
(D) 0.2 cm in opposite phase




Q.13 Path of a ray entering from air to a medium of variable refractive index is shown in figure. Find the value of refractive index of the medium at P. [3]

(A) $\frac{\sqrt{3}}{2}$

(B) $\frac{1}{2}$


(C) $\sqrt{2}$

(D) 2



Q.14 Find the final deviation of the incident ray from a semi-circular mirror of radius 1m as shown in the figure. [3]

- (A) 270° (B) 180°
(C) 60° (D) 0°



Q.15 A equi-biconvex lens of radii of curvature 20 cm has refractive index 1.56 for blue colour and 1.48 for red colour. The approximate linear spread of focus for white light is [3]

- (A) 0.36 cm (B) 0.42 cm (C) 0.30 cm (D) 3 cm

Q.16 When an object is kept at a distance of 20 cm from a concave mirror, the image is formed at a distance of 10 cm. If the object is moved with a speed of 4 m/s along the principal axis, the speed with which the image moves is [3]


- (A) 10 cm/sec (B) 1 m/sec (C) 4 m/sec (D) 0.4 m/sec

- Q.17 Two small angled prisms of prism angle A and B respectively are combined to produce dispersion without deviation. Then the net dispersion produced is [3]
- (A) $(\mu_v - \mu_r)A + (\mu_{v'} - \mu_{r'})B$ (B) $(\mu_{v'} - \mu_{r'})A + (\mu_v - \mu_r)B$
 (C) $(\mu_v - \mu_r)A - (\mu_{v'} - \mu_{r'})B$ (D) $(\mu_{v'} - \mu_{r'})A + (\mu_v - \mu_r)B$

- Q.18 A beam of light AO is incident on a diverging lens as shown in the figure. The focal plane of the lens is also given. The possible direction of the refracted beam will be [3]
-
- (A*) (B) (C) (D) none of the above


- Q.19 An object is placed at distance of 10 cm in air from the front surface of a thick glass slab of thickness 6 cm whose back surface has been silvered and is placed in air. The image is formed at a distance of 14 cm from the silvered surface. The refractive index of the material of the thick mirror is [3]
- (A) 1.2 (B) 1.3 (C) 1.4 (D) 1.5
- Q.20 What should be the value of i (angle of incidence) so that the ray incident on the face AB of the equilateral prism will pass through BC perpendicularly [3]

- (A) $\sin^{-1}\left(\frac{1}{\sqrt{3}}\right)$ (B) $\cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$
 (C) $\sin^{-1}\left(\frac{2}{\sqrt{3}}\right)$ (D) $\cos^{-1}\left(\frac{2}{\sqrt{3}}\right)$



[MATRIX TYPE]


- Q.21 A spherical fish bowl of radius R is placed in front of a plane vertical mirror (M). The thickness of the wall of the fish bowl is very thin. The centre (C) of the spherical bowl is at a distance of 3R from the plane mirror. The bowl is filled with water and contains a fish (F). Fish (F) is at a distance of R from the centre of the spherical bowl as shown in the figure. Refractive index of water is $\frac{4}{3}$. Two surfaces are indicated in the bowl as first surface (1) and second surface (2) [6]




- | | |
|--|------------------------|
| Column-I | Column-II |
| Optical Event | Nature of image |
| (A) Refraction at first surface | (P) Virtual |
| (B) Refraction at second surface after reflection from mirror | (Q) Real |
| (C) Refraction at first surface after reflection from mirror and refraction from second surface. | (R) Magnified |
| | (S) Diminished |

[SUBJECTIVE TYPE]

- Q.22 A transparent cylinder of radius 1m has a mirrored surface on its right half as shown. A ray of light travelling in air is incident on the left side of the cylinder. The ray coming out from the cylinder is parallel to incident ray and at a distance $d = 1.00$ m from the incident ray. Find the refractive index of the material. [5]



- Q.23 A disc of radius r is rolling on a plane horizontal mirror with constant angular velocity ω as shown in the figure. Calculate velocity of image of point Q w.r.t. Q itself. P is contact point and line PQ makes an angle θ with vertical at given instant. [5]




[SINGLE CORRECT CHOICE TYPE]

Q.1 A ray of light passes from a medium of refractive index μ into air. The angle of incidence is found to be half the angle of refraction. What is the angle of refraction? [3]


- (A) $\cos^{-1}(\mu/2)$ (B) $\sin^{-1}(\mu/2)$ (C) $2 \sin^{-1}(\mu/2)$ (D*) $2\cos^{-1}(\mu/2)$

Q.2 Consider the point P as the origin and line OP as the x-axis in the situation shown in the figure. Which of the following represents the coordinates of the image of the point object O. (Take OP = 20 cm) [3]

- (A) (20 cm, 0 cm) (B) (10 cm, $10\sqrt{3}$ cm)
(C*) (-10 cm, $10\sqrt{3}$ cm) (D) (10 cm, $-10\sqrt{3}$ cm)



Q.3 A thin convex lens is cut along its principal axis and one half of the lens is placed as shown in the figure. What are the x and y coordinates respectively, of the image w.r.t. the origin O of the given coordinate axis? The focal length of the lens is 20 cm. [3]




- (A) (60 cm, 12 mm) (B*) (60 cm, 18 mm)
(C) (12 cm, 2.4 mm) (D) (60 cm, -12 mm)

Q.4 An equi convex lens whose focal length in air is 20 cm is made up of a material whose refractive index is 1.5. If one of its surface is silvered and then dipped into a transparent liquid whose refractive index is 1.6. In the liquid it will behave like a [3]

- (A) converging lens (B) diverging lens (C*) converging mirror (D) diverging mirror

Q.5 Refractive index of a transparent slab varies as $\mu = kx$ where x is the distance from origin. Time taken by the light to travel the slab of thickness t (as shown in the figure) [3]

- (A*) $\frac{t^2 k}{2c}$ (B) $\frac{t^2 k}{c}$
(C) $\frac{tk}{c}$ (D) $\frac{2tk}{2c}$




Q.6 A ray moving along unit vector $\frac{1}{2}(\hat{i} + \hat{j} + \sqrt{2}\hat{k})$ strikes a reflecting surface in xy plane. The unit vector along the reflected ray would be [3]

- (A) $-\frac{1}{2}(\hat{i} + \hat{j} + \sqrt{2}\hat{k})$ (B*) $\frac{1}{2}(\hat{i} + \hat{j} - \sqrt{2}\hat{k})$ (C) $-\frac{1}{2}(\hat{i} + \hat{j} - \sqrt{2}\hat{k})$ (D) None of these

Q.7 The lens shown in the diagram has a refractive index higher than that of its surrounding medium. It will converge a paraxial beam incident on it [3]

- (A) from left only (B) from right only
(C) either from left or right (D*) None.



Q.8 A paraxial beam of light having diameter 4 mm is incident on a convex lens having focal length 10 cm. A screen is placed normal to the beam at a distance of 30cm on the other side. Radius of image formed on the screen is- [3]

- (A) 6 mm (B*) 4 mm (C) 3mm (D) 2 mm

Q.9 A ray of light from air is incident at an angle of 70° into a medium having refractive index μ . The reflected and the refracted rays are found to suffer equal deviations in opposite direction. μ equals [3]

- (A) $\tan 70^\circ$ (B*) $2\sin 70^\circ$ (C) $\operatorname{cosec} 70^\circ$ (D) None of these


Q.10 In the figure shown a ray of light hits the interface of the media I & II at angle of 30° . Refractive indices of the media are shown in the figure. The angle made by the ray with normal in medium three is [3]

(A) $\sin^{-1}\left(\frac{2}{3}\right)$

(B) $\sin^{-1}\left(\frac{\sqrt{2}}{3}\right)$


(C) ray will not come in medium III

(D*) $i_3 = \sin^{-1}\left(\frac{2}{7}\right)$




Q.11 A thin prism with angle 4° and refractive index 1.5 is placed inside a thin transparent tube with water (refractive index, $4/3$) as shown. The deviation of light will be [3]

- (A) 2.67° upward (B*) 0.67° downward
(C) 2.67° downward (D) 0.5 downward



Q.12 As shown in the figure particle at M vibrates simple harmonically about M with amplitude 0.1cm perpendicular to axis along a line. Find the amplitude of vibration of image [3]

- (A) 0.3 cm in same phase
(B) 0.1 cm in opposite phase
(C) 0.2 cm in same phase
(D*) 0.2 cm in opposite phase




Q.13 Path of a ray entering from air to a medium of variable refractive index is shown in figure. Find the value of refractive index of the medium at P. [3]

(A) $\frac{\sqrt{3}}{2}$

(B) $\frac{1}{2}$


(C*) $\sqrt{2}$

(D) 2



Q.14 Find the final deviation of the incident ray from a semi-circular mirror of radius 1m as shown in the figure. [3]

- (A*) 270° (B) 180°
(C) 60° (D) 0°



Q.15 A equi-biconvex lens of radii of curvature 20 cm has refractive index 1.56 for blue colour and 1.48 for red colour. The approximate linear spread of focus for white light is [3]

- (A) 0.36 cm (B) 0.42 cm (C) 0.30 cm (D*) 3 cm





Q.16 When an object is kept at a distance of 20 cm from a concave mirror, the image is formed at a distance of 10 cm. If the object is moved with a speed of 4 m/s along the principal axis, the speed with which the image moves is [3]

- (A) 10 cm/sec (B*) 1 m/sec (C) 4 m/sec (D) 0.4 m/sec

Q.17 Two small angled prisms of prism angle A and B respectively are combined to produce dispersion without deviation. Then the net dispersion produced is [3]

- (A) $(\mu_v - \mu_r)A + (\mu_{v'} - \mu_{r'})B$ (B) $(\mu_{v'} - \mu_{r'})A + (\mu_v - \mu_r)B$
 (C*) $(\mu_v - \mu_r)A - (\mu_{v'} - \mu_{r'})B$ (D) $(\mu_{v'} - \mu_{r'})A + (\mu_v - \mu_r)B$

Q.18 A beam of light AO is incident on a diverging lens as shown in the figure. The focal plane of the lens is also given. The possible direction of the refracted beam will be [3]




(D) none of the above

Q.19 An object is placed at distance of 10 cm in air from the front surface of a thick glass slab of thickness 6 cm whose back surface has been silvered and is placed in air. The image is formed at a distance of 14 cm from the silvered surface. The refractive index of the material of the thick mirror is [3]

- (A*) 1.2 (B) 1.3 (C) 1.4 (D) 1.5

Q.20 What should be the value of i (angle of incidence) so that the ray incident on the face AB of the equilateral prism will pass through BC perpendicularly [3]


- (A*) $\sin^{-1}\left(\frac{1}{\sqrt{3}}\right)$ (B) $\cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$
 (C) $\sin^{-1}\left(\frac{2}{\sqrt{3}}\right)$ (D) $\cos^{-1}\left(\frac{2}{\sqrt{3}}\right)$



[MATRIX TYPE]

Q.21 A spherical fish bowl of radius R is placed in front of a plane vertical mirror (M). The thickness of the wall of the fish bowl is very thin. The centre (C) of the spherical bowl is at a distance of 3R from the plane mirror. The bowl is filled with water and contains a fish (F). Fish (F) is at a distance of R from the centre of the spherical bowl as shown in the figure. Refractive index of water is $\frac{4}{3}$. Two surfaces are indicated in the bowl as first surface

(1) and second surface (2) [6]



Column-I

Optical Event

- (A) Refraction at first surface
 (B) Refraction at second surface after reflection from mirror
 (C) Refraction at first surface after reflection from mirror and refraction from second

Column-II

Nature of image

- (P) Virtual
 (Q) Real
 (R) Magnified


surface.

(S) Diminished

[Ans. (A) P,R (B) Q,R (C) Q,S]


[SUBJECTIVE TYPE]

- Q.22 A transparent cylinder of radius 1m has a mirrored surface on its right half as shown. A ray of light travelling in air is incident on the left side of the cylinder. The ray coming out from the cylinder is parallel to incident ray and at a distance $d = 1.00$ m from the incident ray. Find the refractive index of the material. **[5]**



[Ans. $\mu = \frac{1}{2 \sin 15^\circ}$]

- Q.23 A disc of radius r is rolling on a plane horizontal mirror with constant angular velocity ω as shown in the figure. Calculate velocity of image of point Q w.r.t. Q itself. P is contact point and line PQ makes an angle θ with vertical at given instant. **[5]**



[Ans. $2r\omega \sin 2\theta$]