


# BIOTECHNOLOGY AND ITS APPLICATIONS

## BIOLOGY

### Single Correct Answer Type

- First hormone prepared by genetic engineering is:  
a) Insulin                      b) Oxytocin                      c) Adrenaline                      d) Somatotropin
- Retroviruses in animals including humans are able to change normal cells into**  
a) Germ cell                      b) Cancerous cells                      c) Cosmid                      d) Vector
- The restriction enzyme responsible for the cleavage of following sequence is  
5' – G – T – C – G – A – C – 3'  
3' – C – A – G – C – T – G – 5'  
a) *Alu* I                      b) *Bam* HI                      c) *Hind* II                      d) *Eco* RI
- pBR322 was the first artificial cloning vector developed in ...A... by ...B... and ...C... from *E. coli* plasmid. Here A, B and C can be  
a) A-1976, B-Boliver, C-Rodriquez                      b) A-1975, B-Tiselius, C-Rodriquez  
c) A-1977, B-Boliver, C-Rodriquez                      d) A-1978, B-HO Smith, C-KW Wileox
- Transfer of any gene into a completely different organism can be done through  
a) Genetic engineering      b) Tissue culture                      c) Transformation                      d) None of these
- An environmental agent that triggers transcription from an operon is a:  
a) Depressor                      b) Inducer                      c) Regulator                      d) Controlling element
- Recombinant DNA have integrated fragment of  
a) Antibiotic resistant gene                      b) Diseases resistant gene  
c) Allergy resistant gene                      d) All of these
- In plants, the tumour inducing plasmid (Ti) of *Agrobacterium tumefaciens* is used as a cloning vector. This statement is  
a) True                      b) False  
c) Sometimes (a) and sometimes (b)                      d) Neither (a) nor (b)
- If recombinant DNA carrying antibiotic resistance (*e. g.*, ampicillin) is transferred into *E. coli* cell, the host cell is transformed into ampicillin-resistant cells. The ampicillin resistant gene in this case is called a  
a) Vectors                      b) Plasmid                      c) Selectable marker                      d) Cloning sites
- Boviene spongiform encephalopathy disease is equal to:**  
a) Kala Azar                      b) Parkinson's disease  
c) Creutzfeldt-Jacob disease                      d) None of the above
- Known sequence of DNA that is used to find complementary DNA strand is:  
a) Vector                      b) Plasmid                      c) DNA probe                      d) Recombinant DNA
- Proteins are removed by treatment with  
a) Ribonuclease                      b) Chitinase                      c) Cellulase                      d) Protease
- Which of the following key factors, makes plasmid, the vector in genetic engineering?  
a) It is resistant to antibiotics                      b) It is resistant to restriction enzymes  
c) Its ability to carry a foreign gene                      d) Its ability to cause infection in the host
- I. *Ori* also controls the copy numbers of the linked DNA  
II. If a foreign DNA ligates at the *Bam* HI site of tetracycline resistance gene in the vector pBR322, the recombinant plasmid loses the tetracycline resistance due to insertion of foreign DNA  
Choose regarding the above statements  
a) I is true, II is false                      b) II is true, I is false                      c) Both are true                      d) Both are false
- When scientists make an animal superior by view of genotype, introducing some foreign genes in it, the phenomenon is called:

- a) Tissue culture                      b) Biotechnology                      c) Genetic engineering                      d) Immunisation
16. Many copies of a DNA molecule in a test tube are produced by:  
 a) Polymerase chain reaction (PCR)                      b) Molecular chain reaction (MCR)  
 c) Ephemeral chain reaction (ECR)                      d) All of them
17. Producing a 'giant mouse' in the laboratory was possible through:  
 a) Gene mutation                      b) Gene duplication                      c) Gene synthesis                      d) Gene manipulation
18. Downstream process includes  
 I. Separation of the product from the reactor  
 II. Purification of the product  
 III. Formation of the product with suitable preservatives  
 IV. Quality control testing and clinical trials in case of drugs  
 Which of the statements given above are correct?  
 a) I, II and III                      b) I, II and IV                      c) II, III and IV                      d) I, II, III and IV
19. More advancement in genetic engineering is due to  
 a) Restriction endonuclease                      b) Reverse transcription  
 c) Protease                      d) Zymase
20. Plasmid are suitable vectors for gene cloning because  
 a) These are small circular DNA molecules, which can integrate with host chromosomal DNA  
 b) These are small circular DNA molecules with their own replication origin site  
 c) These can shuttle between prokaryotic and eukaryotic cells  
 d) These often carry antibiotic resistance genes
21. Polymerase chain reaction is useful in  
 a) DNA synthesis                      b) DNA amplification  
 c) Protein synthesis                      d) Amino acid synthesis
22. Study the following diagram and identify *A*, *B* and *C*



- a) A- *Taq* polymerase, B-Denaturation at 94°C, C-Primer  
 b) A-Denaturation at 94°C, B- *Taq* polymerase, C-Primer  
 c) A-Primer, B-Denaturation at 94°C, C- *Taq* polymerase  
 d) A- *Taq* polymerase, B-Extension, C-Transformation
23. A bioreactor is  
 a) Hybridoma                      b) Culture containing radioactive isotopes  
 c) Culture for synthesis of new chemicals                      d) Fermentation tank
24. Which of the following techniques can be used to detect genetic disorders in human?  
 a) Polymerase Chain Reaction (PCR)                      b) Gel electrophoresis  
 c) Spectroscopy                      d) All of the above
25. Special sequence in the DNA recognized by restriction endonuclease is called  
 a) Restriction nucleotide sequence                      b) Palindromic nucleotide sequence  
 c) Recognition nucleotide sequence                      d) All of the above
26. Primers are  
 a) Small chemically synthesized oligonucleotides of about 10-18 nucleotides that are complementary to the region of template DNA  
 b) Chemically synthesized oligonucleotides of about 10-18 nucleotides that are not complementary to the region of template DNA  
 c) The double-stranded DNA that need to the amplified


- d) Specific sequences present on recombinant DNA
27. This method of finding a gene is used when researchers very little about the gene they are trying to find. This process results in a complete gene library : a collection of copies of DNA fragments that represent the entire genome of an organism. Identify the method  
 a) Cloning                      b) Shotgun cloning                      c) Gene synthesis                      d) Cloning
28. Consider the following statement about PCR  
 I. Polymerase Chain Reaction (PCR) is a technique of synthesizing multiple copies of the desired gene in *vitro*  
 II. This technique was developed by Kary Mullis in 1985  
 III. A single PCR amplification cycle involves three basic steps; denaturation, annealing and extension  
 Which of the statement given above are correct?  
 a) I and II                      b) I and III                      c) II and III                      d) I, II and III
29. A somatic plant cell has potential to develop into a full plant. This is called:  
 a) Totipotency                      b) Gene cloning                      c) Tissue culture                      d) Regeneration
30. *Ori* is a DNA sequence that is responsible for initiating replication. This statement is  
 a) True                      b) False  
 c) Sometimes (a) and sometimes (b)                      d) Neither (a) nor (b)
31. Plasmids are autonomously replicating circular extrachromosomal DNA. This statement is  
 a) True                      b) False  
 c) Sometimes (a) and sometimes (b)                      d) Neither (a) nor (b)
32. Genetic engineering is possible because:  
 a) The phenomenon of transduction in bacteria is well understood  
 b) We can see DNA by electron microscope  
 c) We can cut DNA at specific sites by endonucleases like DNA ase I  
 d) Restriction endonucleases purified form bacteria can be used in vitro
33. A single PCR amplification cycle involves  
 a) Denaturation                      b) Annealing                      c) Extension                      d) All of these
34. DNA fingerprinting is related to:  
 a) Molecular analysis of profiles of DNA samples  
 b) Analysis of DNA samples using imprinting devices  
 c) Techniques used for molecular analysis of different specimens of DNA  
 d) Techniques used in identification of fingerprints of different persons
35. The basic of DNA fingerprinting is:  
 a) The double helix                      b) Errors in base sequence  
 c) Polymorphism in sequence                      d) DNA replication
36. In genetic engineering, the terms vector is applied for:  
 a) Plasmid                      b) Sources of DNA                      c) Cell which receives                      d) Virus
37. Which of the following are used to gene cloning?  
 a) Nucleoids                      b) Chromosomes                      c) Mesosomes                      d) Plasmid
38. The process that preserves the distribution of DNA fragments in the gel while creating replica on the filter is one of the following  
 a) Directed sequencing of BAC counting                      b) Random shotgun sequencing  
 c) Electrophoresis                      d) Southern blotting
39. Two enzymes responsible for restricting the growth of bacteriophages in *E. coli* were isolated. One was methylase and other was restriction endonuclease. What is the significance of methylase?  
 a) Protection of host DNA from the action of restriction endonuclease by adding methyl group to one or two bases usually with in the sequence recognized by restriction enzyme  
 b) Able to ligate the two cohesive ends of DNA molecule  
 c) Able to remove the methyl group and hence, prevent the action of restriction endonuclease on host DNA  
 d) Able to cut the DNA of bacteriophage at specific sites
40. Single-stranded DNA molecules that can bind to and be used to detect other DNA molecules are called

- a) Primer                                      b) STRs                                      c) RFLPs                                      d) Probes

41. Which of the following enzyme is used in genetic engineering?

- a) Translocase                                      b) Topoisomerase  
c) DNase                                      d) Restriction endonuclease

42. The below diagram refer to PCR. Identify the steps A, B and C and select the correct option



- a) A-Denaturation of 94-96°C, B-Annealing of 40-60°C, C-Extension through *taq* polymerase at 72°C, D-Amplified  
b) A-Annealing of 94-96°C, B-Denaturation of 40-60°C, C-Extension through *taq* polymerase at 72°C, D-Amplified  
c) A-Extension through *taq* polymerase at 40-60°C, B-Amplified, C-Denaturation of 40-60°C, D-Annealing of 94-96°C  
d) A-Annealing, B-Extension through *taq* polymerase at 40-60°C, C-Denaturation of 94-96°C, D-Annealing of 40-60°C

43. The controlled use of biological agents, such as microorganism, plants or animal cell, for beneficial use is called

- a) Biochemistry                                      b) Molecular biology                                      c) Biotechnology                                      d) Microbiology

44. Humulin is a:

- a) Pig insulin                                      b) Human insulin                                      c) Viral insulin                                      d) Human clone

45. Find the incorrect statement:

- a) Gene therapy is a genetic engineering technique used to treat disease at molecular level by replacing defective genes with normal genes  
b) Calcitonin is a medically useful recombinant product in the treatment of infertility  
c) Bt toxin is a biodegradable insecticide obtained from *Bacillus thuringiensis*  
d) *Trichoderma* sp. is a biocontrol agent for fungal diseases of plants

46. Plasmids are extrachromosomal circular DNA molecules:

- a) Which have their own point of replication and can replicate independently  
b) Which have their own point of replication but cannot replicate independently  
c) Which do not have their own point of replication and cannot replicate independent of bacterial of bacterial chromosomal DNA  
d) None of the above

47. The genome map was produced under human genome project in:

- a) 1992                                      b) 1994                                      c) 1996                                      d) 2000


48. Term hybridoma implies:

- a) DNA-RNA hybrid                                      b) Recombination of DNA molecules  
c) Somatic hybridisation                                      d) Genetic fusion

49. Which of the following is a difficulty in getting prokaryotic cells to express eukaryotic genes?

- a) The signals that control gene expression are different and prokaryotic promoter regions must be added to the vector  
b) The genetic code differs between the two because prokaryotes substitute the base uracil for thymine

- c) Prokaryotic cells cannot transcribe introns because their genes do not have them  
 d) The ribosomes of prokaryotes are not large enough to handle long eukaryotic genes
50. In transgenics, the expression of transgene in the target tissue is known by:  
 a) Enhancer                                      b) Transgene                                      c) Promoter                                      d) Reporter
51. Identify A, B, C and D in the given diagram of *E. coli* cloning vector pBR322



- a) A-*Eco* RI, B-*Bam* HI, C-*Ori*, D-*amp*<sup>R</sup>                                      b) A- *amp*<sup>R</sup>, B- *Ori*, C-*Bam* HI, D-*Eco* RI  
 c) A-*Ori*, B-*Bam* HI, C-*Eco* RI, D-*amp*<sup>R</sup>                                      d) A-*Bam* HI, B-*Eco* RI, C- *amp*<sup>R</sup>, D-*Ori*
52. Consider the following statements  
 I. In microinjection method foreign DNA is directly injected into the nucleus of animal cell or plant cell by using micro needles or micro pipettes  
 II. Microinjection method is used in oocytes, eggs and embryo  
 III. Electroporation is the formation of temporary pores in the plasma membrane of host cell by using lysozyme or calcium chloride  
 IV. In chemical mediated gene transfer method certain chemicals such as CO<sub>2</sub> help foreign DNA to enter the host cell
- Which of the statements given above are correct?  
 a) I and II                                      b) I, II and III                                      c) II, III and IV                                      d) I, II, III and IV
53. The construction of the first recombinant DNA was done by using the native plasmid of:  
 a) *E. coli*                                      b) *Salmonella typhimurium*  
 c) *B. thuringiensis*                                      d) Yeast
54. Gene amplification using primers can be done by  
 a) Microinjection                                      b) ELISA  
 c) Polymerase chain reaction                                      d) Gene gun
55. Polyethylene glycol method is used for  
 a) Biodiesel production                                      b) Seedless fruit production  
 c) Energy production from sewage                                      d) Gene transfer without a vector
56. The enzymes, commonly used in genetic engineering are  
 a) Restriction endonuclease and polymerase                                      b) Endonuclease and ligase  
 c) Restriction endonuclease and ligase                                      d) Ligase and polymerase
57. Which one of the following techniques had helped to solve many mysteries involving murders, robberies and rapes?  
 a) Gene splicing                                      b) Computer technology  
 c) DNA fingerprinting                                      d) Gene cloning
58. Consider the following statements  
 I. Recombinant DNA technology popularly known as genetic engineering is a stream of biotechnology which deals with the manipulation of genetic material by man *in vitro*  
 II. pBR322 is the first artificial cloning vector developed in 1977 by Boliver and Rodriquez from *E. coli* plasmid  
 III. Restriction enzymes belongs to a class of enzymes called nucleases
- Which of the statements given above are correct?  
 a) I and II                                      b) I and III                                      c) II and III                                      d) I, II and III
59. What is C-DNA?

- a) Circular DNA  
 b) Cloned DNA  
 c) DNA produced from reverse transcription of RNA  
 d) Cytoplasmic DNA
60. PCR was developed by ...A... in ...B... and for this he received Nobel Prize for chemistry in ...C.... Here A, B and C can be recognized as  
 A                      B                      C  
 a) Kary Mullis 1990 1997                      b) Flemming 1985 1993  
 c) Kary Mullis 1985 1993                      d) Flemming 1990 1997
61. Cutting of a piece of DNA from a plasmid was done with the help of ...A... enzymes, popularly known as ...B.... Here A and B can be  
 a) A-Tu ligases; B-Molecular glu                      b) A-Restriction enzyme; B-Molecular scissors  
 c) A-Joining enzyme; B-Molecular glu                      d) A-DNA polymerases; B-Synthesising enzymes
62. In a genetic engineering experiment, restriction enzymes can be used for  
 a) Bacterial DNA only                      b) Viral DNA only  
 c) Any DNA fragment                      d) Eukaryotic DNA only
63. The components of a bioreactor are  
 I. an agitator system  
 II. an oxygen delivery system  
 III. foam control system  
 IV. temperature control system  
 V. pH control system  
 VI. sampling ports to with draw cultures periodically  
 Choose the correct option  
 a) I, II, III, IV and V                      b) II, IV, V and VI                      c) I, II, III, IV and VI                      d) All of these
64. The minimum length of cistron in base pairs which synthesizes a polypeptide of 50 amino acids is:  
 a) 50 bp                      b) 100 bp                      c) 150 bp                      d) 200 bp
65. I. DNA being a hydrophilic molecule cannot pass through cell membranes  
 II. The bacteria should be made competent to accept the DNA molecule  
 The correct option regarding the above statements is  
 a) I is true, but II is false                      b) II is true, but I is false  
 c) I and II are true                      d) I and III are false
66. In cloning plasmid pBR322  
 p stands for ...A...  
 B stands for ...B...  
 R stands for ...C...  
 Choose the correct option  
 a) A-plasmid, B-Boliver, C-Rodriquez                      b) A-plasmid, B-bacteria, C-Rodriquez  
 c) A-prophage, B-bacteriophage, C-Rodriquez                      d) A-prophage, B-Boliver, C-Rodriquez
67. Blood stains are found at the site of murder. If DNA profiling technique is to be used for identifying the criminal, which of the following is ideal for use?  
 a) Serum                      b) Erythrocytes                      c) Leucocytes                      d) Platelets
68. The Ti plasmid used in genetic engineering is obtained from:  
 a) *Bacillus thuringiensis*                      b) *Agrobacterium rhizogenes*  
 c) *Agrobacterium tumifaciens*                      d) *Escherichia coli*
69. Who got the Nobel prize in medicine for their discovery of 'G-proteins' and the role of these proteins in the cells:  
 a) Robert and Philip Sharp                      b) Gilman and Rodbell  
 c) Fischer and Krebs                      d) Ervin Nahar and Bert Sakmann
70. Which of the following is required to perform polymerase chain reaction?  
 I. DNA template

II. Primer

III. *Taq* polymerase and *vent* polymerase

Choose the correct option

a) I, II and III                      b) I and II                      c) II and III                      d) II and III

71. The basis for DNA fingerprinting is:
- Occurrence of restriction fragment length polymorphism (RFLP)
  - Phenotypic differences between individuals
  - Availability of cloned DNA
  - Knowledge of human karyotype
72. In genetic engineering, a DNA segment (gene) of interest, is transferred to the host cell through a vector. Consider the following four agents (I-IV) in this regard and select the correct option about which one or more of these can be used as a vector/vectors
- Bacterium
  - Plasmid
  - Plasmodium*
  - Bacteriophage
- a) I, II and IV                      b) I only                      c) I and III                      d) II and IV
73. Transfer of any gene into a completely different organism can be done through
- Genetic engineering
  - Tissue culture
  - Transformation
  - None of these
74. Thermostable enzymes '*taq*' and '*vent*' isolated from thermophilic bacteria are
- DNA polymerase
  - DNA ligases
  - Restriction endonucleases
  - RNA polymerases
75. Due to chloramphenicol resistance gene, one is able to select a transformed cell in the presence of chloramphenicol. The chloramphenicol resistance gene in this case is called
- Origin of replication
  - Selectable marker
  - Cloning sites
  - Insertional inactivation
76. GAATTC is the recognition site for the restriction endonuclease
- Eco*RI
  - Hind*II
  - Eco*RII
  - Bam*HI
77. Plasmid is
- An autonomously replicating circular extrachromosomal DNA
  - An autonomously replicating circular extrachromosomal RNA
  - An circular protein molecules
  - An autonomously replicating chromosomal DNA
78. The polymerase chain reaction is a technique that is used for
- In vivo* replication of DNA
  - In vivo* synthesis of *mRNA*
  - In vitro* synthesis of *mRNA*
  - In vitro* replication of specific DNA sequence using thermostable DNA polymerase
79. Yeast has become important in genetic engineering because it:
- Has plasmids that can be genetically engineered
  - Allows the study of eukaryotic gene regulation and expression
  - Grows readily and rapidly in the laboratory
  - All of the above
80. The genome of *Caenorhabditis elegans* consists of:
- 3 billion base pairs and 30,000 genes
  - 12 million base pairs and 6,000 genes
  - 4.7 million base pairs and 4,000 genes
  - 97 million base pairs and 18,000 genes
81. Two bacteria found to be very useful in genetic engineering experiments are:
- Nitrosomonas* and *Klebsiella*
  - Escherichia* and *Agrobacterium*
  - Nitrobacter* and *Azotobacter*
  - Rhizobium* and *Diplococcus*
82. Gel electrophoresis is used for:
- Isolation of DNA molecule





97. The total number of nitrogenous bases in human genome is estimated to be about:  
 a) 35 million                      b) 3.1 billion                      c) 3.5 million                      d) 35 thousand
98. Totipotency in cell is:  
 a) Flower in a culture medium  
 b) Development of fruit from a flower in a culture medium  
 c) Development of an organism from cell in culture medium  
 d) Development of all tissues of all kinds from a cell in a culture medium
99. Restriction enzymes was discovered by  
 a) Alexander Flemming                      b) Waksman  
 c) Berg                      d) Smith, Nathan and Arber
100. Identify the plasmid:  
 a) Alu I                      b) Hind III                      c) Eco RI                      d) pBR 322
101. Consider the following statements  
 I. Bioreactors are vessels of large volumes in which raw materials are biologically converted into specific products  
 II. One of the most commonly used bioreactors is of stirring type  
 III. Shake flasks are used for growing and mixing the desired materials on a small scale in the laboratory  
 IV. A large scale production of desired biotechnological product is done by using 'bioreactors'  
 a) I and II                      b) I and III                      c) I, II and III                      d) I, II, III and IV
102. The term 'Biotechnology' was given by  
 a) Craig Venter                      b) Robert Edward                      c) Karl Erkey                      d) Temin and Baltimore
103. A collection of organisms, usually viruses, bacteria or yeast, which have been transformed by the addition of extra genes from another species:  
 a) Gene replication                      b) Gene cloning                      c) Gene pool                      d) Gene library
104. Exonucleases cleaving nucleotides one at a time from the end of the polynucleotide chain are:  
 a) Specific for 5' end of RNA strand  
 b) Specific for 3' end of RNA strand  
 c) Specific for both 5' and 3' ends of nucleotide strands  
 d) Non-specific for 5' and 3' ends of nucleotide strand
105. The genetic recombinants obtained by insertion of plasmid into 1 phage genome is called:  
 a) Cosmid                      b) Phasmid                      c) Phagmid                      d) Foreign DNA
106. Which of the following statements is true?  
 a) In the historic cloning experiment of Dr. Wilmut, the transplanted nucleus was taken from an udder cell  
 b) Mammalian characters appeared first in dinosaurs  
 c) Heart of mammals is incapable of being in vitro  
 d) Pyramid of biomass is upright in pond ecosystem
107. Which of the following statement is not true?  
 I. DNA being a hydrophilic molecule cannot pass through cell membranes  
 II. *Agrobacterium tumefaciens* delivers a piece of DNA known as 'Z-DNA' in the Ti-plasmid which transforms normal plant cells into tumour cells to produce chemical against pathogens  
 III. Retrovirus, adenovirus, papillomavirus are also now used as cloning vectors in animal because of their ability to transform normal cells into cancerous cell.  
 IV. In genetic engineering, DNA from different sources are cut with the same restriction enzymes so that both DNA fragments have same kind of sticky ends  
 Choose the correct option  
 a) Only I                      b) Only II                      c) Only III                      d) Only IV
108. Which one of the following pairs is correctly matched?  
 a) RNA polymerase -RNA primer                      b) Restriction enzymes-Genetic Engineering  
 c) Central Dogma-codon                      d) Okazaki fragments-splicing
109. Bam HI, Eco RI, Sma H are the types of:  
 a) Restriction endooxidases                      b) Restriction endonucleases





- c) I, II and III  
d) None of the above
133. Process of formation of RNA from DNA is called  
a) Transduction                      b) Transcription                      c) Transformation                      d) Translation
134. Which of the following would not be used in preparing recombinant DNA?  
a) Plasmids                                      b) Phages  
c) Restriction enzymes                      d) DNA polymerase III
135. Which one of the following bacteria has found extensive use in genetic engineering work in plants?  
a) *Agrobacterium tumefaciens*                      b) *Clostridium septicum*  
c) *Xanthomonas citri*                      d) *Bacillus coagulens*
136. Which of the following components are used in gel electrophoresis?  
I. Ethidium bromide  
II. Restriction endonuclease  
III. Agarose  
IV. UV radiation  
Choose the correct option  
a) I and II                      b) I and III                      c) I, II and IV                      d) I, II, III and IV
137. What is the first step in Southern Blotting technique?  
a) Isolation of DNA from a nucleated cell such as the one from the scene of crime  
b) Denaturation of DNA on the gel for hybridization with specific probe  
c) Production of group of genetically identical cells  
d) Digestion of DNA by restriction enzyme
138. The most thoroughly studied of the known bacteria-plant interaction is the:  
a) Plant growth simulation by phosphate-solubilising bacteria  
b) Cyanobacterial symbiosis with some aquatic ferns  
c) Gall formation on certain angiosperms by *Agrobacterium*  
d) Nodulation of *Sesbania* stems by nitrogen fixing bacteria
139. Microorganisms can be grown in the bioreactor by  
a) Support growth system                      b) Agitated growth system  
c) Suspended growth system                      d) Both (a) and (b)
140. In Northern blotting RNAs are separated by gel electrophoresis and the RNA bands are transferred onto a membrane of:  
a) Diazobenzyl oxymethyl                      b) Diazobenzene  
c) Diazobromine                      d) None of the above
141. Which one of the following is commonly used in transfer of foreign DNA into crop plants?  
a) *Trichoderma harzianum*                      b) *Meloidogyne incognita*  
c) *Agrobacterium tumefaciens*                      d) *Penicillium expansum*
142. Which one among the following is just a cloning plasmid not an expression plasmid?  
a) pBAD-18-Cam                      b) pBCSK                      c) pUC 18                      d) pET
143. There ...A... are the DNA molecules that can carry a foreign ...B... segment into the host cell.  
Here A and B refers to  
A                      B  
a) Vector RNA                      b) Vector DNA  
c) Gene RNA                      d) Gene DNA
144. Probes, used in DNA fingerprinting are initially  
a) Single-stranded RNA                      b) Mini satellite  
c) 19 base long oligonucleotides                      d) All of the above
145. Application of PCR are  
I. detection of pathogens  
II. diagnosis of specific mutation  
III. DNA fingerprinting

Choose the correct option

- a) I and II                      b) I and III                      c) II and III                      d) I, II and III
146. A clone of sheep Dolly has been made by:  
a) Gene transfer                      b) Somatic cell cloning  
c) Nucleus transfer                      d) Germinal cell cloning
147. T<sub>1</sub>-plasmid used in genetic engineering is obtained from  
a) *Bacillus thuringiensis*                      b) *Agrobacterium rhizogenes*  
c) *Agrobacterium tumefaciens*                      d) *Pseudomonas syringae*
148. The role of DNA ligase in the construction of a recombinant DNA molecule is  
a) Formation of phosphodiester bond between two DNA fragments  
b) Formation of hydrogen bonds between sticky ends of DNA fragments  
c) Ligation of all purine and pyrimidine bases  
d) None of the above
149. Transgenic animals are produced by injecting foreign gene into the:  
a) Egg                      b) Nucleus of unfertilized egg  
c) Nucleus of fertilized egg                      d) Nucleus of sperm
150. Clonal cell lines can be obtained by:  
a) Autoradiography                      b) Tissue culture                      c) Centrifugation                      d) Cell fractionation
151. Electroporation procedure involves:  
a) Fast passage of food through sieve pores in phloem elements with the help of electric stimulation  
b) Opening of stomatal pores during night by artificial light  
c) Making transient pores in the cell membrane to introduce gene constructs  
d) Purification of saline water with the help of a membrane system
152. Which of the following is associated with genetic engineering?  
a) Plastids                      b) Plasmids                      c) Mutations                      d) Hybrid vigour
153. Biolistics (gene gun) is suitable for  
a) Disarming pathogen vectors                      b) Transformation of plants cells  
c) Construction recombinant DNA by joining with vectors                      d) DNA fingerprinting
154. Which of the following statements are correct for the enzyme *taq* polymerases?  
I. *Taq* polymerase is thermally unstable  
II. It requires primers for carrying out the process of polymerization  
III. *Taq* polymerase is isolated from thermophilic bacterium, *Thermus aquaticus*  
Choose the correct option  
a) I and II                      b) I and III                      c) II and III                      d) I, II and III
155. EFB stands for  
a) European Federation of Biotechnology                      b) Eurasian Federation of Biotechnology  
c) East Asia Federation of Biotechnology                      d) Ethiopian Federation of Biotechnology
156. The commonly used DNA fingerprinting technique in forensic studies is simply a method involving  
a) Southern blotting                      b) Northern blotting                      c) Eastern blotting                      d) Western blotting
157. *Cry* I endotoxins obtained from *Bacillus thuringiensis* are effective against  
a) Nematodes                      b) Bollworms                      c) Mosquitoes                      d) Flies
158. In the naming of restriction enzymes the first letter is derived from ...A... name and next two letters from the ...B... and fourth letter from ...C... of ...D... where the enzymes are extracted  
A to D in the statement can be  
A      B      C      D  
a) Genus species strain bacteria                      b) Species genus strain bacteria  
c) Genus species variety eukaryote                      d) Species genus variety eukaryote
159. Which of the following techniques is most commonly used to separate DNA molecules by size?  
a) Chromatography                      b) PCR                      c) RFLP                      d) Gel electrophoresis
160. Which one of the following scientists got the Nobel Prize for his invention polymerase chain reaction

- (PCR)?
- a) F. Sanger                      b) Paul Berg                      c) Alec Jeffreys                      d) Kary B. Mullis
161. Which is non-invasive technique of genetic counselling?
- a) Amniocentesis                      b) Chorionic biopsy  
c) Foetal blood sampling                      d) Ultrasonography
162. The colonies of recombinant bacteria appear white in contrast to blue colonies of non-recombinant bacteria because of:
- a) Insertional inactivation of alpha-galactosidase in non-recombinant bacteria  
b) Insertional inactivation of alpha-galactosidase in recombinant bacteria  
c) Inactivation of glycosidase enzyme in recombinant bacteria  
d) Non-recombinant bacteria containing beta-galactosidase
163. Which of the following steps are catalyzed by *taq* polymerase in a PCR reaction?
- a) Denaturation of template DNA                      b) Annealing of primers to template DNA  
c) Extension of primer end on the template DNA                      d) All of the above
164. I. In the process of recombinant DNA technology after several treatment the purified DNA is precipitated by adding chilled ethanol  
II. The bacterial/plant, animal cell is broken down by enzymes to release DNA, along with RNA, proteins, polysaccharides and lipids  
Choose the correct option for above statements
- a) I is true, but II is false                      b) I is false, but II is true  
c) I and II are true                      d) I and II are false
165. Which of the statements are correct about bioreactors?
- I. It provides all the optimal conditions for achieving the desired product by providing optimal growth conditions like temperature, pH, substrate, salt, vitamin and oxygen  
II. It is suited for large-scale production of microorganisms under aseptic conditions for a number of days  
Correct option is
- a) Only I                      b) Only II                      c) I and II                      d) None of the above
166. *Taq* polymerase enzyme used in PCR is isolated from
- a) *Thermus aquaticus*                      b) *Thermococcus litoralis*  
c) *Salmonella typhimurium*                      d) None of the above
167. The first hormone artificially produced by culturing bacteria is:
- a) Insulin                      b) Thyroxine                      c) Testosterone                      d) Adrenaline
168. A gene is made up of:
- a) DNA                      b) RNA                      c) Either DNA or RNA                      d) Amino acids
169. The first restriction endonuclease type II ...A..., was isolated by Smith, Wilcox and Kelley from ...B... bacterium. It was formed to cut DNA molecules at a particular point by recognizing a specific sequence of six base pairs, known as the ...C.... Here A, B and C can be
- |                   |                                  |                      |
|-------------------|----------------------------------|----------------------|
| A                 | B                                | C                    |
| a) <i>Eco</i> RI  | <i>Escherichia</i> RY 13         | Restriction sequence |
| b) <i>Eco</i> RII | <i>E. coli</i> R 245             | Recognition sequence |
| c) <i>Hind</i> II | <i>Haemophilus influenza</i>     | Recognition sequence |
| d) <i>Bam</i> HI  | <i>Bacillus amyoliquefaciens</i> | Restriction sequence |
170. In gel electrophoresis, the separated DNA fragments are visualized after staining the DNA with ...A... followed by exposure to ...B...  
Here A and B refers to
- |                      |                    |
|----------------------|--------------------|
| <b>A</b>             | <b>B</b>           |
| a) B-galactosidase   | Infrared radiation |
| b) Ethidium bromide  | UV radiation       |
| c) Ethidium nitrate  | $\gamma$ -rays     |
| d) Ethidium chloride | Radiowave          |
171. In DNA fingerprinting:
- a) A positive identification can be made

- b) Multiple restriction enzyme digests/generate unique fragments
- c) The polymerase chain reaction amplifies fewer DNA
- d) The variability of repeated sequences between two restriction sites is evaluated

172. Cosmid is:

- a) Extragenetic material in Mycoplasma
- b) Circular DNA in bacteria
- c) Extra DNA in bacteria
- d) Fragment of DNA inserted in bacteria for forming copies

173. Following enzymes/chemical/technique are used in the process of gel electrophoresis

- I. sample DNA is cut into fragments
- II. restriction endonucleases
- III. agarose gel
- IV. ethidium bromide
- V. UV-radiation
- VI. elution

Mark the correct sequence of their use

- a) I, II, III, VI, V and IV
- b) I, II, III, VI, V and IV
- c) IV, V, VI, I, II and III
- d) I, II, IV, V, VI and III

174. Improvement of genotype of an organism by addition of some foreign genes is:

- a) Genetic diversity
- b) Gene handling
- c) Tissue culture
- d) Genetic engineering

175. Which one is a true statement regarding DNA polymerase used in polymerase chain reaction?

- a) DNA polymerase is responsible for DNA synthesis
- b) It is isolated from Protozoa
- c) It serves as a selectable marker
- d) It is used to ligate introduced DNA in recipient plant cell

176. Most sensitive technique to detect malignant cell in non-hodgkin's lymphoma is

- a) Polymerase chain reaction
- b) Gene therapy
- c) Stem cell therapy
- d) None of the above

177. Gene therapy involves:

- a) Introducing of a normal genes in cell
- b) Eliminating defective and useless genes
- c) Treating of defective genes with radiations
- d) Replacement of defective genes by normal ones

178. Human Genome project was the thought of:

- a) Jean Dausset
- b) Watson
- c) Crick
- d) None of the above

179. Which conserved motifs are found in *E. coli* genes?

- a) TATA box
- b) CAAT box
- c) Pribnow box
- d) All of these

180. Given below is a sample of a portion of DNA strand giving the base sequence on the opposite strands. What is so special shown in it?

5' \_\_\_\_\_ GAATTC \_\_\_\_\_ 3'  
3' \_\_\_\_\_ CTTAAG \_\_\_\_\_ 5'

- a) Replication completed
- b) Deletion mutation
- c) Start codon at the 5' end
- d) Palindromic sequence of base pairs

181. The DNA used as a carrier for transferring a fragment of foreign DNA into a suitable host is called

- a) Cloning vector
- b) Vehicle DNA
- c) Gene carrier
- d) All of these

182. The nuclease enzyme, which begins its attack from free end of a polynucleotide, is?

- a) Exonuclease
- b) Kinase
- c) Polymerase
- d) Endonuclease

183. Genetically engineered bacterium used in production of:


- a) Thyroxine
- b) Human insulin
- c) Epinephrine
- d) Cortisol

184. In Southern blotting..... is separated by gel electrophoresis:

- a) DNA
- b) m-RNA
- c) t-RNA
- d) Protein

185. Taq polymerase enzyme is found in:

- a) *Thermus aquateucus*    b) *E. coli*    c) *Pseudomonas*    d) *Agrobacterium*
186. The process used for separation of protein in polyacrylamide gel is called:  
 a) Southern blotting    b) Northern blotting    c) Western blotting    d) Eastern blotting
187. Which of the following methods(s) is used to introduce foreign DNA into host cells?  
 a) Gene gun method    b) Gel electrophoresis    c) Elution    d) Extension
188. The figure shown three steps (A, B, C) of Polymerase Chain Reaction PCR. Select the option giving correct identification together with what represents?



- a) B-denaturation at a temperature of about 98°C separating the two DNA strands  
 b) A-denaturation at a temperature of about 50°C  
 c) C-extension in the presence of heat stable DNA polymerase  
 d) A-annealing with three sets of primers
189. DNA fingerprinting method is very useful for:  
 a) DNA tests for identity and relationships    b) Forensic studies  
 c) Polymorphism    d) All of the above
190. Restriction endonucleases are used as:  
 a) Molecular build up at nucleotides  
 b) Molecular degradation to DNA breakup  
 c) Molecular knives for cutting DNA at specific sites  
 d) Molecular cement to combine DNA sites
191. After completion of the biosynthetic stage in the bioreactors, the product undergoes. Separation and purification processes, collectively termed as  
 a) Transformation    b) Transduction  
 c) Downstream processing    d) Upstream processing
192. Which of the following should be chosen for best yield if one has to produce a recombinant protein or enzyme on a large scale, using microbial plants/animal/human cell?  
 a) Stirred-tank bioreactor    b) Electrophoresis  
 c) Laboratory flask of largest capacity    d) All of the above
193. Go through the figure and select the option for C and D. Here A and B are taken as vector/plasmid DNA and foreign DNA respectively



**Restriction enzyme recognizing palindrome C**    **Enzyme joining the sticky ends D**

- a) *Eco* RI    DNA ligase    b) DNA ligase    *Eco* RI  
 c) Exonuclease    DNA ligase    d) DNA ligase    Exonuclease
194. Which of the following is known as molecular scissors of DNA?  
 a) Ligase    b) Polymerases  
 c) Restriction endonucleases    d) Transcriptase
195. A kind of biotechnology involving manipulation of DNA is  
 a) DNA replication    b) Genetic engineering    c) Denaturation    d) Renaturation


196. Harris and J.F. Watkins in 1965 first time reported the fusion of following cell lines to form hybrids:

- a) Mouse and man    b) Mouse and hamster





211. Stirred-tank bioreactors have advantages over shake flasks because they
- Provide high temperature and pH
  - Provide better aeration and mixing properties
  - Do not allow the entry of CO<sub>2</sub>
  - Are easy to operate
212. During 'gene cloning' which is called a gene taxi?
- Vaccine
  - Plasmid
  - Bacteria
  - Protozoa
213. TATAATG sequence near the RNA start point of prokaryotic promoter is:
- Nicks
  - DNA marker
  - Pallindrome
  - Pribnow box
214. I. Copy number is defined as the number of copies of plasmid present in a cell  
 II. It varies from 15-100 copies per cell  
 Choose regarding the above statements
- I is true, II is false
  - II is true, I is false
  - Both are true
  - Both are false
215. Which one of the following hydrolyses internal phosphodiester bonds in a polynucleotide chain?
- Lipase
  - Protease
  - Exonuclease
  - Endonuclease
216. What does Bt stand for the popular crop Bt cotton?
- Best
  - Best type
  - Biotechnology
  - Bacillus thuringiensis
217. Which of the following statement is incorrect?
- Cosmid contains gene coding for viral protein
  - Cosmid replicates like plasmids
  - Cosmid has antibiotic resistant marker gene
  - Cos site has 12 bases helping to join complete genome to make it circular
218. An attenuated virus:
- Is a virus that is non-pathogenic
  - In an elongated viral particle
  - Can transfer recombinant DNA to other viruses
  - Will not produce an immune response
219. Which of the following has popularized the PCR (polymerase chain reaction)?
- Easy availability of DNA template
  - Availability of synthetic primers
  - Availability of cheap deoxyribonucleotides
  - Availability of 'Thermostable' DNA polymerase
220. Choose the correct statement with reference to 'Dolly':
- She was created by taking nucleus from unfertilized eggs and cytoplasm from unfertilized eggs
  - She was created by taking nucleus from under udder cells and cytoplasm from unfertilized eggs
  - She was created by taking cytoplasm from udder cell and nucleus from unfertilized eggs
  - She was created by taking cytoplasm from udder cell and nucleus from fertilized eggs
221. The first recombinant DNA was constructed by
- Stanley Cohen
  - Herbert Boyer
  - Both (a) and (b)
  - Temin and Baltimore
222. Study the given diagram and identify the enzymes A and B involves in steps I and II



Step I

a) *Eco*RI

Step II

DNA ligase

b) *Alu*I

DNA ligase

- c) *Hind*II                                      DNA polymerase                                      d) Restriction endonuclease DNA polymerase
223. Which one of the following is a correct statement
- "Bt" in "Bt-cotton" indicates that it is a genetically modified organism produced through biotechnology
  - Somatic hybridization involves fusion of two complete plant cells carrying desired genes
  - The anticoagulant hirudin is being produced from transgenic *Brassica napus* seeds
  - "Flavr Savr" variety of tomato has enhanced the production of ethylene which improves its taste
224. The transgenic animals are those which have:
- Foreign RNA in all its cell
  - Foreign DNA in all its cells
  - Foreign DNA in some of its cells
  - Both 'A' and 'C'
225. Which of the following is not correctly matched for the organism and its cell wall degrading enzyme?
- Plant cells-Cellulase
  - Algae-Methylase
  - Fungi-Chitinase
  - Bacteria-Lysozyme
226. Petroleum-lysing bacteria are being engineering for the removal of oil spills. What is the most realistic danger of these bacteria to the environment?
- Mutations leading to the production of a strain pathogenic to humans
  - Extinction of natural microbes due to the competitive advantage of the "petro-bacterium"
  - Destruction of natural oil deposits
  - Poisoning of the food chain
227. c-DNA probes are copied from the messenger RNA molecules with the help of:
- Restriction enzymes
  - Reverse transcriptase
  - DNA polymerase
  - Adenosine deaminase
228. Mishandling of genetic engineering may cause:
- Genetic erosion
  - Green revolution
  - Silver revolution
  - White revolution
229. Gene for cloning may be chemically synthesized:
- When the exact sequence of nucleotides is known
  - Through the use of restriction enzymes and gel electrophoresis to separate restriction fragments
  - By the Sanger method
  - By making complementary DNA from genes without introns
230. Source of *taq* polymerase used in PCR is a
- Thermophilic fungus
  - Mesophilic fungus
  - Thermophilic bacterium
  - Halophilic bacterium
231. Genetic engineering has been successfully used for producing:
- Transgenic models for studying new treatment for certain cardiac diseases
  - Transgenic Cow-Rosie which produces high fat milk for making ghee
  - Animals like bulls for farm work as they have super power
  - Transgenic mice for testing safety of polio vaccine before sue in humans
232. Which of the following is used as a best genetic vector in plants?
- Bacillus thuringiensis*
  - Agrobacterium fumefaciens*
  - Pseudomonas putida*
  - None of the above
233. Plants in comparison to animals are more rapidly manipulated by genetic engineering. Select out the most probable reason for this
- Totipotency shown by plant cells
  - Single somatic cell can regenerate a whole plant body
  - Genetic engineering is supplemented with plant tissue culture techniques
  - All of the above
234. Which of the following pairs is correctly matched?
- Central dogma-Codon
  - Okazaki fragments-Splicing
  - RNA polymerase-RNA primer
  - Restriction enzymes-Genetic engineering
235. Recombinant DNA technology is related with:
- Stanley Cohen and Harbert Boyer
  - Bateson and Punnet
  - Huxley and Harvey
  - Schleiden and Schwann
236. Western blotting technique was developed by:

- a) Alwin                                      b) Edwin                                      c) Towbin                                      d) Thomas

237. In recombinant DNA technique, the term vector refers to a  
 a) Donor DNA, it is identified and picked up through electrophoresis  
 b) Plasmid, transfers DNA into living cell  
 c) Collection of entire genome in form of plasmid  
 d) Enzyme, cuts the DNA at specific sites

238. Complete transduction is:  
 a) Transfer of whole genome with the help of virus  
 b) Picking up of one or more genes by a phage and transfer it to second host  
 c) Integration of gene brought by viral particle into genome of new host  
 d) Both B and C


239. The function of polymerase chain reaction (PCR) is:  
 a) Translation                                      b) Transduction                                      c) DNA amplification                                      d) None of these

240. The steps involved in the Southern blot test are as follows

- I. X-ray film  
 II. Electrophoresis  
 III. Digestion with restriction enzyme  
 IV. Ethidium bromide  
 V. Radioactive probe

Choose the option having correct sequential order of these events

- a) III, II, IV, V and I                                      b) III, IV, II, V and I                                      c) III, II, V, IV and I                                      d) II, IV, III, V and I
241. The given figure is the diagrammatic representation of the *E. coli* vector pBR322. Which one of the given options correctly identifies its certain component(s)?



- a) *Ori*-original restriction enzymes                                      b) *Rop*-reduced osmotic pressure  
 c) *Hind*III, *Eco*RI-selectable markers                                      d) *amp*<sup>R</sup>, *tet*<sup>R</sup>-antibiotic resistance genes

242. The restriction enzyme(s) used in recombinant DNA technology that make staggered cuts in DNA leaving sticky ends is/are

- a) *Eco*RI                                      b) *Hind*II                                      c) *Bam*HI                                      d) All of the above

243. RNA processing is:

- a) An event that occurs after RNA transcribed                                      b) The rejection of old, worn-out RNA  
 c) An event that occurs before RNA is transcribed                                      d) Both (A) and (C)

244. Find out the wrong statements

- a) Mobile genetic elements, transposons were visualized by Barbara McClintock  
 b) Udder cell and somatic cell is used to produce the cloned sheep by nuclear transplantation method  
 c) In pedigree analysis, a person immediately affected by and action is called propositus  
 d) DNA ligases are used to cleave a DNA molecule

245. Widely used tool in genetic engineering of crop plants is:


- a) Protoplast fusion                                      b) Transposon  
 c) Microinjection                                      d) Agrobacterium mediation

246. DNA fingerprinting method is very useful for:

- a) DNA tests for identity and relationships                                      b) Forensic studies  
 c) Polymorphism                                      d) All of the above



260. Identify and match the labelled items *A, B, C, D, E, F* and *G* in the diagram below from the list I-VII given with components



- I. DNA polymerase
- II. plasmid
- III. plasmid with 'sticky ends'
- IV. DNA ligase
- V. restriction endonuclease
- VI. recombinant DNA
- VII. reverse transcriptase

The correct components are

A B C D E F G

- a) VII I II V III IV VI
- c) VII V III I II IV VI

- b) VII VI V IV III II I
- d) I II IV VI III V VII

261. A technology which has found immense use in solving cases of disputed parentage is:

- a) DNA fingerprinting
- b) Polymerase chain reaction
- c) Recombinant DNA technology
- d) Monoclonal antibody production

262. The most important feature in a plasmid to be used as a vector is

- a) Origin of replication
- b) Presence of a selectable marker
- c) Presence of sites for restriction endonuclease
- d) Its size

263. DNA gyrase, the enzyme that participates in the process of DNA replication is a type of

- a) DNA ligase
- b) DNA polymerase
- c) DNA topoisomerase
- d) Reverse transcriptase

264. Abnormal gene is replaced by normal gene through:

- a) Gene therapy
- b) Medicines
- c) Cloning
- d) Radiation

265. The key tools required for the recombinant DNA technology are

- I. restriction enzymes
- II. Polymerase enzymes
- III. host organism ligases
- IV. Vectors
- V. host organisms

Select the correct option

- a) I, II and III
- b) I, III, IV and V
- c) I, II, III and V
- d) I, II, III, IV and V

266. A tumour inducing plasmid widely used in the production of transgenic plants in that of:

- a) *Escherichia coli*
- b) *Bacillus thuringiensis*
- c) *Staphylococcus aureus*
- d) *Agrobacterium tumefaciens*


267. Which one of the following palindromic base sequences in DNA can be easily cut at about the middle by some particular restriction enzyme?

- a) 5' \_\_\_\_\_ GATATG \_\_\_\_\_ 3'  
3' \_\_\_\_\_ CTAATA \_\_\_\_\_ 5'
- b) 5' \_\_\_\_\_ GAATTC \_\_\_\_\_ 3'  
3' \_\_\_\_\_ CTTAAG \_\_\_\_\_ 5'



- d) By treating the DNA with restriction endonucleases
281. Transgenic organisms are produced by:
- Deleting sex chromosomes
  - Inducing gene mutations
  - Introducing foreign genes
  - Arresting spindle fibre formation
282. Manipulation of gene and genetic material by man is a fast emerging branch of science which started with the formation of recombinant DNA molecules. This branch of science is named as
- Recombinant DNA technology
  - Genetic engineering
  - DNA manipulation biotechnology
  - All of the above
283. Ligases catalyse the formation of bonds between
- C = C
  - P = O
  - C - C
  - H - H
284. The characteristics of a molecular probe are
- very long molecule
  - double-stranded
  - DNA or RNA
  - complementary to a part of desired gene
- The correct pair is
- I and II
  - II and III
  - III and IV
  - IV and I
285. VNTR analysis involves
- Analyzing specific loci for two base repeating units usually less than 100 bp in size
  - Analyzing specific loci for 2-4 bp repeating units
  - PCR amplification of specific genes
  - Cutting DNA with restriction enzyme and analyzing the banding pattern of fragments
286. Manipulation of DNA in genetic engineering became possible due to the discovery of
- Restriction endonuclease
  - DNA ligase
  - Transcriptase
  - Primase

287. Study the given figure carefully and select the correct statements regarding this



- It represents typical agarose gel electrophoresis which showing differential migration of DNA fragments
  - Lane 1 contains undigested DNA fragments
  - Lanes 2 to 4 contains digested DNA fragment
  - Smallest DNA bands are present at (A) position and largest DNA bands are present at (B) position
- I, II and III
  - I, II and IV
  - II and III
  - III and IV
288. Matching sequence of DNA between two evidences, one of the criminal with the suspect is known as:
- DNA fingerprinting
  - DNA amplification
  - Gene mapping
  - DNA resolution
289. Alec Jeffreys developed the DNA fingerprinting technique. The probe he used was
- Ribozyme
  - Sex chromosomes
  - SNP
  - VNTR
290. In addition to *taq* polymerase enzyme which other thermostable DNA polymerases have been isolated to be used in Polymerase Chain Reaction (PCR)?
- Vent* polymerase
  - Pfu* polymerase
  - Both (a) and (b)
  - None of these
291. PCR proceeds in three distinct steps governed by temperature. They are in order of
- Denaturation, synthesis (polymerization), annealing
  - Annealing, synthesis (polymerization), denaturation
  - Synthesis (polymerization), annealing, denaturation
  - Denaturation, annealing, synthesis (polymerization)
292. One of the following is transgenic of organisms:
- Holly sheep and Flavr savr tomato
  - Holly sheep and Cotton Bt





- a) Diplococcus sp. and Pseudomonas sp.  
 b) Crown gall bacterium and Caenorhabditis elegans  
 c) Escherichia coli and Agrobacterium tumefaciens  
 d) Vibrio cholerae and a tailed bacteriophage
303. Minisatellite or Variable Number Tandem Repeat (VNTR) are used in  
 a) Gene therapy                      b) Gene mapping                      c) DNA fingerprinting                      d) Restriction enzymes
304. Having become an expert on gel electrophoresis, you are asked to examine a gel for a colleague. Where would you find the smallest segment of DNA?  
 a) Near the positive electrode, farthest away from the wells  
 b) Near the negative electrode, close to the wells  
 c) Near the top, near the negative pole  
 d) Near the middle they tend to slow-down after the first few minutes
305. Improvement of genotype of an organism by addition of some foreign genes is:  
 a) Genetic diversity                      b) Gene handling                      c) Tissue culture                      d) Genetic engineering
306. The structure involved in genetic engineering is  
 a) Codon                      b) Anticodon                      c) Vector                      d) Plasmid
307. In agarose gel electrophoresis, DNA molecules are separated on the basis of their  
 a) Charge only                      b) Size only                      c) Charge to size ratio                      d) All of these
308. In gel electrophoresis, the sample DNA is cut into fragments by  
 a) Restriction endonucleases                      b) Exonuclease  
 c) Endonuclease                      d) Anhydro L-galactose
309. Molecular scissors, which cut DNA at specific site:  
 a) Ligase                      b) Cellulase  
 c) Pectinase                      d) Polymerase
310. PCR stands for:  
 a) Polymerase Cyclic Reaction                      b) Polymerase Chain Reaction  
 c) Polyethyl Cytosine Reaction                      d) Polymerization Chain Reaction
311. In case of polymerase chain reaction, temperature, required for the steps  
 A. Denaturation  
 B. Annealing  
 C. Extension  
 a) A-94°C, B-40°C, C-72°C                      b) A-40°C, B-72°C, C-94°C  
 c) A-72°C, B-94°C, C-40°C                      d) A-94°C, B-72°C, C-40°C
312. DNA can be introduced into any cell by:  
 a) Injection  
 b) Being complexed with calcium salts  
 c) Being placed along with the cell into a gene gun  
 d) Gel electrophoresis
313. An improved variety of transgenic basmati rice:  
 a) Gives high yield and is rich in Vitamin A  
 b) Is completely resistant to all insect pests and diseases of paddy  
 c) Gives high yield but has no characteristic aroma  
 d) Does not require chemical fertilizers and growth hormones
314. Which of the following organelles is associated with genetic engineering?  
 a) Plastids                      b) Plasmids                      c) Chloroplast                      d) Mitochondria
315. Human genome contains about:  
 a) 10,000 nucleotides                      b) 10,000 genes                      c) 6 billion nucleotides                      d) 6 billion genes
316. An artificial process of infecting cells with naked viral DNA is:  
 a) Translation                      b) Transduction                      c) Transfection                      d) Transgenic
317. Match the correct one:  
 a) RNA Polymerase-RNA primer                      b) Respiration-Lysosome

- c) Restriction enzyme-genetic engineering                      d) Central dogma-DNA structure


318. For transformation, microparticles coated with DNA are to be bombarded with gene gun are made up of:

- a) Platinum or Zinc                      b) Silicon or Platinum                      c) Gold or Tungsten                      d) Silver or Platinum

319. You are attempting to introduce a gene that imparts larval moth resistance to bean plants. Which of the following vectors are you most likely to use?

- a) Phage DNA                      b) Bacterial plasmid                      c) Ti plasmid                      d) Yeast plasmid

320. Simple stirred-tank bioreactor is given below. Identify A,B,C,D and E



	A	B	C	D	E
a)	Motor	Foam braker	Sterile air	Steam for sterilization	Acid/ Base of pH control

c)	Acid/ Base of pH control	Motor	Foam braker	Sterile air	Steam for sterilization
----	--------------------------	-------	-------------	-------------	-------------------------

b)	Foam braker	Sterile air	Steam for sterilization	Acid/ Base of pH control	
d)	Sterile air	Steam for sterilization	Foam braker	Motor	Acid/ Base of pH control

321. Protein engineering is used to study the proteins to compare the catalytic properties of:

- a) Normal and mutated form of enzyme                      b) Normal form of enzyme  
c) Mutated form of enzyme                      d) Normal and mutated form of proteins

322. Genes that are involved in turning on or off the transcription of a set of structural genes are called:

- a) Polymorphic genes                      b) Operator genes                      c) Redundant genes                      d) Regulatory genes

323. The experimental manipulation of DNA of different species, producing recombination DNA is known as

- a) Gel electrophoresis                      b) Transformation  
c) Genetic engineering                      d) Replication technology

324. Plasmid is used as carrier because:

- a) It has both ends with replicating points  
b) It has no free ends  
c) It is circular DNA with a capacity of binding with eukaryotic DNA  
d) All of the above

325. Which of the following statement is correct in the context of observing DNA separated by agarose gel electrophoresis?

- a) DNA can be seen in visible light  
b) DNA can be seen without staining in visible light  
c) Ethidium bromide stained DNA can be seen in visible light  
d) Ethidium bromide stained DNA can be seen under exposure to UV light

326. Nitrogen fixing genes are called:

- a) 'Nif' genes                      b) Plasmid genes                      c) Leg genes                      d) Cos genes


327. The genetically-modified (GM) brinjal in India has been developed for:

- a) Enhancing shelf life                      b) Enhancing mineral content  
c) Drought-resistance                      d) Insect-resistance

328. Variable number of tandem repeats (VTNRs) in the DNA molecule are highly useful in:

- a) Monoclonal antibody production                      b) DNA fingerprinting

- c) Recombinant DNA technology
329. Protoplasts of two different species are fused in:
- a) Clona propagation
- c) Micropropagation
330. Identify the correct match for the given diagram
- d) Stem cell culture
- b) Organography
- d) Somatic hybridization



**Apparatus                      function**

- a) Gene gun - Vectorless direct gene transfer
- b) Electrophoresis - Differential migration of DNA fragments
- c) Bioreactor - Raw materials are biologically converted into specific products
- d) Respirometer - Finding out rate of respiration
331. DNA fingerprinting technique was first developed by:
- a) Jeffreys, Wilson and Thein
- b) Schleiden and Schwann
- c) Edward and Steptoe
- d) Boysen and Jensen
332. Using recombinant technology, genes from a donor cell can be transplanted into a bacterium for DNA replication and protein synthesis. The kinds of cells that can be used as a donor in this technology are
- a) Bacteria
- b) Either yeast or bacteria
- c) Eukaryotic cells
- d) Any kind of cell
333. Transformation is defined as the procedure by which a piece of ...A... is introduced into a ...B... host. Here A and B refers to
- A                      B
- a) RNA      Virus
- b) DNA      Bacteria
- c) RNA      Bacteria
- d) DNA      Virus

# BIOTECHNOLOGY AND ITS APPLICATIONS

## BIOLOGY

### : ANSWER KEY :

1)	a	2)	b	3)	c	4)	c	173)	b	174)	a	175)	a	176)	a
5)	a	6)	b	7)	a	8)	a	177)	d	178)	a	179)	c	180)	d
9)	c	10)	c	11)	c	12)	d	181)	d	182)	a	183)	b	184)	a
13)	c	14)	c	15)	c	16)	a	185)	a	186)	c	187)	a	188)	c
17)	d	18)	d	19)	a	20)	b	189)	d	190)	c	191)	c	192)	a
21)	b	22)	a	23)	d	24)	a	193)	a	194)	c	195)	b	196)	a
25)	b	26)	a	27)	b	28)	d	197)	d	198)	b	199)	a	200)	c
29)	a	30)	a	31)	a	32)	d	201)	d	202)	d	203)	a	204)	a
33)	d	34)	a	35)	c	36)	a	205)	d	206)	c	207)	d	208)	c
37)	d	38)	d	39)	a	40)	d	209)	d	210)	a	211)	b	212)	b
41)	d	42)	a	43)	c	44)	b	213)	d	214)	c	215)	d	216)	d
45)	b	46)	a	47)	b	48)	c	217)	a	218)	a	219)	d	220)	b
49)	a	50)	d	51)	a	52)	a	221)	c	222)	a	223)	c	224)	b
53)	b	54)	c	55)	d	56)	c	225)	b	226)	c	227)	d	228)	a
57)	c	58)	d	59)	b	60)	c	229)	a	230)	c	231)	d	232)	b
61)	b	62)	c	63)	d	64)	c	233)	d	234)	b	235)	a	236)	c
65)	c	66)	a	67)	c	68)	c	237)	b	238)	c	239)	c	240)	a
69)	b	70)	a	71)	a	72)	d	241)	d	242)	d	243)	a	244)	d
73)	a	74)	a	75)	b	76)	a	245)	d	246)	d	247)	a	248)	a
77)	a	78)	d	79)	d	80)	d	249)	b	250)	c	251)	b	252)	b
81)	b	82)	c	83)	a	84)	d	253)	b	254)	b	255)	d	256)	d
85)	a	86)	b	87)	a	88)	b	257)	b	258)	c	259)	b	260)	a
89)	a	90)	d	91)	d	92)	a	261)	a	262)	a	263)	c	264)	a
93)	b	94)	d	95)	a	96)	a	265)	d	266)	d	267)	b	268)	d
97)	b	98)	c	99)	d	100)	d	269)	a	270)	d	271)	b	272)	a
101)	d	102)	c	103)	d	104)	c	273)	c	274)	c	275)	b	276)	d
105)	b	106)	a	107)	b	108)	b	277)	a	278)	a	279)	d	280)	b
109)	b	110)	b	111)	b	112)	b	281)	c	282)	d	283)	b	284)	c
113)	d	114)	d	115)	c	116)	c	285)	d	286)	a	287)	a	288)	a
117)	a	118)	c	119)	c	120)	a	289)	d	290)	c	291)	d	292)	d
121)	a	122)	c	123)	b	124)	d	293)	d	294)	d	295)	c	296)	a
125)	a	126)	a	127)	a	128)	b	297)	a	298)	a	299)	c	300)	c
129)	b	130)	a	131)	d	132)	d	301)	c	302)	c	303)	c	304)	a
133)	b	134)	d	135)	a	136)	d	305)	d	306)	d	307)	b	308)	a
137)	d	138)	c	139)	d	140)	a	309)	c	310)	b	311)	a	312)	b
141)	a	142)	c	143)	b	144)	b	313)	a	314)	b	315)	c	316)	c
145)	d	146)	c	147)	c	148)	a	317)	c	318)	c	319)	c	320)	a
149)	c	150)	b	151)	c	152)	b	321)	a	322)	b	323)	c	324)	c
153)	b	154)	c	155)	a	156)	a	325)	d	326)	a	327)	d	328)	b
157)	b	158)	a	159)	d	160)	d	329)	d	330)	c	331)	a	332)	d
161)	d	162)	c	163)	c	164)	c	333)	b						
165)	c	166)	a	167)	a	168)	c								
169)	c	170)	b	171)	d	172)	d								

# BIOTECHNOLOGY AND ITS APPLICATIONS

## BIOLOGY

### : HINTS AND SOLUTIONS :

- 2 **(b)**  
Retroviruses in animals including humans are able to change normal cells into cancerous cell
- 4 **(c)**  
pBR322 vector was the first artificial cloning vector constructed in 1977 by Boliver and Rodriquez. It is widely used in gene cloning experiments in pBR322  
**p** – Denotes that it is plasmid  
**BR** – stands for Boliver and Rodriquez who constructed this plasmid  
322 is a number given to distinguish this plasmid from others developed in the same laboratory
- 5 **(a)**  
Genetic engineering is defined as the modification of genetic information of living organism by direct manipulation of their DNA. Thus, a gene of known function (economic importance) can be transferred from its normal location into a cell *via* a suitable mobile genetic element called vector such as plasmid, phage, etc.
- 7 **(a)**  
Recombinant DNA having integrated fragment of antibiotic resistant gene
- 8 **(a)**  
True. In plants, the tumour inducing plasmid ( $T_i$ ) of *Agrobacterium tumefaciens* is used as a cloning vector
- 9 **(c)**  
Gene encoding resistance to antibiotics like ampicillin, chloramphenicol, tetracycline or Kanamycin, are useful selectable markers for *E.coli*. The normal *E.coli* cells do not carry resistance against any of these antibiotics
- 12 **(d)**  
Proteins are removed by treatment with protease
- 13 **(c)**  
Plasmids, cosmids or bacteriophages can be used as vector in genetic engineering. Plasmids are most widely used circular, extrachromosomal DNA segments seen in the bacterial cells. They carry a foreign gene or desired gene to the host. The size of plasmids ranges from  $1 \times 10^6$  to  $200 \times 10^6$  daltons
- 14 **(c)**  
Both are true, *Ori* also controls the copy numbers of the linked DNA  
If a foreign DNA ligates at the *Bam* HI site tetracycline resistance gene in the vector pBR322, the recombinant plasmid loses the tetracycline
- 18 **(d)**  
After the formation of the product in the bioreactors, it undergoes through some processes before a finished product to be ready for marketing. *The processes include* (i) separation and (ii) purification of product which are collectively called the downstream processing  
The product is subjected to quality control testing and kept in suitable preservatives. If drugs are to be manufactured such formulation has to undergo through clinical trials. A proper quality control testing for each product is also needed. The downstream processing and quality control test are different from product to product
- 19 **(a)**  
Endonucleases are enzymes that produce internal cuts called cleavage DNA molecule. A class of endonucleases cleavage DNA only within or near those sites which have specific base sequences, such endonucleases are known as restriction endonucleases and sites recognized by them are called recognition sites. Restriction endonucleases have major role in genetic engineering
- 20 **(b)**  
Plasmid is an extrachromosomal genetic of DNA that is capable of replicating independently of host chromosome. It forms the basis of many cloning vectors used in genetic engineering
- 21 **(b)**  
PCR was discovered by Kary Mullis. In Polymerase Chain Reaction (PCR), a segment of DNA is amplified. *Taq* DNA polymerase enzyme is used PCR, this enzyme is temperature resistant
- 22 **(a)**  
A- *Taq* polymerase, B-Denaturation (air), C-Prime

23 **(d)**  
Bioreactors (fermenters) are considered as vessel in which raw material are biologically converted into specific products by microbes, plant and animal cells and/or their enzymes

24 **(a)**  
By using PCR phenylketonuria, muscular dystrophy, sickle-cell anaemia, hepatitis, chlamydia and tuberculosis can be diagnosed

26 **(a)**  
Primers are small chemically synthesized oligonucleotides of about 10-18 nucleotides long that are complementary to the sequences present at the 3' ends of the target DNA segment

27 **(b)**  
Shotgun cloning involves cutting the DNA of the entire genome into pieces with restriction enzyme, inserting these pieces or fragments into bacteria or yeast with plasmids or viruses and allowing the organism to reproduce making copies or clones of the DNA fragments

28 **(d)**  
The Polymerase Chain Reaction or PCR, as it is commonly called, was originally invented by Kary Mullis in 1985. Kary Mullis shared the Nobel Prize with Michael Smith in Chemistry in 1993. PCR is best defined as the DNA replication *in vitro*. A single PCR amplification cycle involves three basis steps; denaturation, annealing and extension (polymerization)

30 **(a)**  
True, *Ori* is a DNA sequence that is responsible for initiating replication. Any piece of DNA, which linked to this sequence can replicated with in the host cells

31 **(a)**  
True. Plasmids are autonomously replicating circular extra-chromosomal DNA

33 **(d)**  
*PCR is carried out in the following three steps*  
Denaturation, Annealing and Extension

37 **(d)**  
Plasmid which is extra chromosomal DNA molecule and help in gene cloning

38 **(d)**  
A restriction fragment containing a specific gene of interest can be identified by gel electrophoresis followed by transferring of DNA to a membrane as a solid support matrix using a procedure called a Southern blot

39 **(a)**

Protection of host DNA from the action of restriction endonuclease by adding methyl group to one or two bases usually with in the sequence recognized by restriction enzyme

40 **(d)**  
Single stranded DNA molecules that can bind to and be used to detect other DNA molecule are called probes

42 **(a)**  
**Principle of PCR** The purpose of a PCR (Polymerase Chain Reaction) is to make a huge number of copies of a gene. This is necessary to have enough starting template for sequencing There are three major steps in a PCR, which are repeated for 30 or 40 cycles. This is done on an automated cyclers, which can heat and cool the tubes with the reaction mixture in a very short time

(i) **Denaturation at 95°C** During the denaturation, the double-strand melts open to single-stranded DNA, all enzymatic reactions stop (for example : the extension from a previous cycle)

(ii) **Annealing at 54°C** The primers are jiggling around, caused by the Brownian motion. Ionic bonds are constantly formed and broken between the single-stranded primer and the single-stranded template. The more stable bounds last a little bit longer (primers that fit exactly) and on that little piece of double-stranded DNA (template and primer), the polymerase can attach and starts copying the template. Once there are a few bases built in, the ionic bond is so strong between the template and the primer, that it does not break anymore

(iii) **Extension at 72°C** This is the ideal working temperature for the polymerase. The primers, where there are a few bases built in, a already have a stronger ionic attraction to the template than the forces breaking these attractions.

Primers, that are on positions with no exact match, get loose again (because of the higher temperature) and don't give an extension of the fragment


The bases (complementary to the template) are coupled to the primer on the 3' side (the polymerase adds *d*NTPs from 5' to 3', reading the template from 3' to 5' side, bases are added complementary to the template)

43 **(c)**  
The controlled use of biological agents, such as microorganism, plants or animal cell, for

beneficial use is called biotechnology. This technology involves cutting and pasting of desired DNA fragments into the specified hosts for the benefits of human beings

51 (a)

*Eco*RI.



52 (a)

**Microinjection** DNA is directly injected into plant protoplasts or cells (specifically into the nucleus or cytoplasm) using fine tipped (0.5-1.0 micrometer diameter) glass needle or micropipette. This method of gene transfer is used to introduce DNA into large cells such as oocytes, eggs, and the cells of early embryo

**Electroporation** It involves a pulse of high voltage applied to protoplasts/cells/tissues to make transient (temporary) pores in the plasma membranes which facilitates the uptake of foreign DNA

The cells are placed in a solution containing DNA and subjected to electrical shock to cause holes in the membranes. The foreign DNA fragments enter through the holes into the cytoplasm and then to nucleus

**Chemical Mediated Gene Transfer** Chemicals like Polyethylene Glycol (PEG) and sulphate induce DNA uptake into plant protoplasts. Calcium phosphate is also used to transfer DNA into cultured cells

55 (d)

Polyethylene glycol method is used for gene transfer without a vector. It is a chemical method for direct gene transfer to protoplast

56 (c)

Restriction endonucleases and ligase are commonly used enzymes in genetic engineering

57 (c)

DNA fingerprinting is a modern technique that compares sets of DNA by locating identical sequences of nucleotides. It is often used to solve many mysteries involving murders,

robberies and rapes

58 (d)

Genetic engineering is a branch of biotechnology, which deals with the manipulation of genetic material by man. The technique of genetic engineering includes

(i) formation of 'recombinant DNA'

(ii) use of gene cloning

(iii) gene transfer

1. pBR 322 was the first artificial cloning vector constructed in 1977 by Boliver and Rodriguer. It is widely used in gene cloning experiments

2. Restriction enzymes belongs to a class of enzymes called nucleases

60 (c)

A - Key Mullis

B - 1985

C - 1993

61 (b)

Cutting of piece of DNA from a plasmid was done with the help of restriction enzyme, popularly known as molecular scissors

62 (c)

Different kinds of specific enzymes are used in genetic engineering, e.g., cleaving enzymes → These enzymes are used to break DNA molecules  
*They are of three types*

(i) Exonucleases

(ii) Endonucleases

(iii) Restriction endonucleases

63 (d)

*Components of a bioreactors*

An agitator system

An oxygen delivery system

Foam control system

Temperature control system

pH control system

sampling ports to withdraw culture periodically

65 (c)

Both are true

66 (a)

A-plasmid, B-Boliver, C-Rodriquez.

pBR322 vector was the first artificial cloning vector constructed in 1977 by Boliver and Rodriquez. It is widely used in gene cloning experiments in pBR322

**p** - Denotes that it is plasmid

**BR** - stands for Boliver and Rodriquez who



- constructed this plasmid  
322 is a number given to distinguish this plasmid from others developed in the same laboratory
- 67 **(c)**  
DNA fingerprinting is a technique to identify a person on the basis of person's DNA specificity. The technique is based upon the fact that the DNA constitution of an individual carries some specific sequence of nucleotides, which do not carry any information for protein synthesis  
From the given options, leucocytes are to be used for identifying the criminal because they are nucleotide, whereas erythrocytes are enucleated
- 70 **(a)**  
*The basic requirements of a PCR reaction are the following*  
**DNA Template** Any source that contains one or more target DNA molecules to be amplified can be taken as template  
**Two Nucleotide Primers** Primers, which are oligonucleotides, that hybridise to the target DNA region, one to each strand of the double helix  
**Enzyme** *Taq* polymerase and *vent* polymerase
- 72 **(d)**  
Circular plasmid DNA which is used as a vector, can be cleaved at one site with the help of enzyme to give a linear DNA molecule. A foreign DNA segment can now be inserted, by joining the ends of broken circular DNA to the two ends of foreign DNA, thus regenerating a bigger circular DNA molecule that can now be separated by gel electrophoresis on the basis of its size  
Bacteriophages provide another source of cloning vectors. Since, usually, a phage has a linear DNA molecule, a single break will generate two fragments, which are later joined together with foreign DNA to generate a chimeric phage particle
- 73 **(a)**  
Genetic engineering is defined as the modification of genetic information of living organisms by direct manipulation of their DNA  
Thus, a gene of known function (or economic importance) can be transferred from its normal location into a cell *via* a suitable mobile genetic element called vector such as plasmid phage, etc.
- 74 **(a)**  
Thermostable enzymes '*Taq* and *Vent*' isolated from thermophilic bacteria are DNA polymerase  
*Taq* polymerase, isolated from a *Thermophilic bacterium, Thermus aquaticus* and *vent* polymerase, isolated from a thermophilic bacterium *Thermococcus litoralis*
- 75 **(b)**  
Due to chlorophenicol resistance gene, one is able to select a transformed cell in the presence of chloramphenicol. The chloramphenicol resistance gene in this case is called selectable marker
- 76 **(a)**  
The restriction endonuclease *Eco* RI is obtained from *Esherichia coli* RY 13. The recognition sequence for this is GAATTC, CTTAAG
- 77 **(a)**  
Autonomously replicating circular extrachromosomal DNA.  
Manipulation of gene and genetic material by man is a fast emerging branch of science, which started with the formation of recombinant DNA molecule. This branch of science is named as recombinant DNA technology, genetic engineering and DNA manipulation technology, genetic engineering and DNA manipulation technology. This technology involves cutting and pasting of desired DNA fragments into the specified hosts for the benefits of human beings
- 78 **(d)**  
The polymerase chain reaction is a technique that is used for *in vitro* replication of specific DNA sequence using thermostable DNA polymerase. The polymerase chain reaction or PCR, was originally invented by Kary Mullis in 1985. Kary Mullis shared the Nobel Prize with Michael Smith in chemistry in 1993
- 86 **(b)**  
The Polymerase Chain Reaction (PCR) is a technique by which small samples of DNA can be quickly amplified. The repeated amplification is achieved by the use of thermostable DNA polymerase (*i.e.*, *taq* polymerase isolated from a bacterium, *Thermus aquaticus*) which remain active during the high temperature induced denaturation of double-stranded DNA
- 88 **(b)**  
Exonucleases remove nucleotides from the terminal ends (either 5' or 3') of DNA in one strand of duplex
- 90 **(d)**  
PCR is a technique of synthesizing multiple copies of the desired gene or (DNA) *in vitro*. *The basic requirement of PCR* are DNA template, two nucleotide primers and enzyme (DNA polymerase)
- 91 **(d)**

- Agrobacterium tumefaciens* (soil inhabiting plant bacterium) is a pathogen of several dicot plants. It delivers a piece of DNA known as 'T-DNA' in the Ti-plasmid which transforms normal plant cells into tumour cells to produce chemicals against pathogens
- 92 **(a)**  
Restriction endonuclease recognize a specific DNA base sequence (recognition sequence, recognition site, restriction sequence or restriction site having palindromic sequence) and cleaves both the strands of DNA at or near that site. The enzyme cuts the DNA, generating restriction fragments with overhanging ends or blunt ends
- 95 **(a)**  
*Agrobacterium tumefaciens* (updated scientific name *Rhizobium radiobacter*) is the casual agent of crown gall disease (the formation on tumour) in over 140 species of dicot. It is a rod-shaped, Gram negative, soil bacterium (Smith, *et. al* 1907). Symptoms are caused by the insertion of a small segment of DNA, known as T-DNA (transfer DNA) into the plant cell, which is incorporated at a semi-random location into the plant genome
- 96 **(a)**  
True, the polymerase chain reaction is a reaction in which amplification of specific DNA sequences is carried out in *vitro*
- 99 **(d)**  
Restriction enzyme are known as molecular knives or molecular scissors and are used to cut DNA at specific sites of DNA. These were first discovered by Smith, Nathan and Arber
- 101 **(d)**  
Small volume cultures are usually employed in laboratories for research and production of less quantities of products. *e.g.*, in shake flasks. However, large scale production of the products is carried out in 'bioreactor'  
Bioreactors are large vessels (having a volume of 100 to 1000 L) which are used for biological conversion of raw materials into specific products. The most commonly used bioreactors are of stirring type
- 102 **(c)**  
The term 'Biotechnology' was given in 1917 by a Hungarian Engineer, Karl Erkey, to describe a process or large scale production of pigs
- 107 **(b)**  
*Agrobacterium tumefaciens* delivers a piece of DNA known as 'T-DNA' in the Ti-plasmid which transforms normal plant cells into tumour cells to produce chemical against pathogens
- 110 **(b)**  
Kary Mullis  
Gene encoding resistance to antibiotics like ampicillin, chloramphenicol, tetracycline or Kanamycin, are useful selectable markers for *E.coli*. The normal *E.coli* cells do not carry resistance against any of these antibiotics
- 114 **(d)**  
Ti-plasmid is found in *Agrobacterium tumefaciens*, which produces crown gall (tomour) in a large number of dicot species. *A. tumefaciens* is a Gram negative soil bacterium that infects a wide range of plants and causes crown galls
- 115 **(c)**  
The science of recombinant technology took birth when Cohen and Boyer (1972) were able to introduce a piece of antibiotic resistance gene containing foreign DNA into plasmid of *Salmonella typhimurium*. This modified plasmid was them inserted into *E. coli* to get clones of recombinant DNA. Thus, Cohen and Boyer discovered recombinant technology
- 116 **(c)**  
In recombinant DNA technology, a desired segment of DNA or a gene is made to combine with the DNA of an organism where it will multiply and produce it copies. Plasmids and viruses are the most commonly used cloning vectors in recombinant DNA technology
- 119 **(c)**  
Selectable marker helps to select the host cells which contain the vector and eliminate the non-transformants. Genes encoding resistance to antibiotics like ampicillin, chloramphenicol, tetracycline or kanamycin are useful selectable markers of *E.coli*. The normal *E.coli* cells do not carry resistance against any of these antibiotics
- 122 **(c)**  
Herbert Boyer discovered that restriction enzymes have the capability of cutting DNA strands in a particular fashion, which left what has become known as sticky ends on the strands
- 123 **(b)**  
A Southern blot.  
A restriction fragment containing a specific gene of interest can be identified by gel electrophoresis followed by transferring of DNA to a membrane as a solid support matrix using a procedure called a



- (particularly cancer of mouth) near wounds
- 154 **(c)**  
The final step in PCR is extension (polymerization), where in *Taq* DNA polymerase synthesizes the DNA region between the primers using deoxynucleotide triphosphates and  $Mg^{2+}$ . It means the primers are extended towards each other so that the DNA segment lying between the two primer is copied. The optimum temperature for this polymerization step is  $72^{\circ}C$   
*Taq* polymerase is thermostable enzyme, isolated from Thermophilic bacterium, *Thermus aquaticus*
- 155 **(a)**  
EFB – European Federation of Biotechnology  
A definition of biotechnology which covers both traditional views and modern molecular biotechnology has been given by European Federation of Biotechnology. According to EFB “Biotechnology is the integrated use of biochemistry, microbiology and engineering sciences in order to achieve technological application of the capabilities of microorganisms, cultured tissues/cells and part there of”
- 156 **(a)**  
A technique developed by EM Southern in 1975 for detection of a specific DNA sequences (gene or other) in a large, complex sample of DNA (*e. g.*, cellular DNA). It is also used to determine the molecular weight of a restriction fragment and to measure relative amounts in different sample  
**Uses** Southern blots are used in gene discovery and mapping, evolution and development studies, diagnostics and forensics  
In regards to genetically modified organisms, Southern blotting is used as a definitive test to ensure that a particular section of DNA of known genetic sequence has been successfully incorporated into the genome of the host organism
- 157 **(b)**  
*CryI* endotoxins obtained from *Bacillus thuringiensis* are effective against bollworm larvae
- 158 **(a)**  
In the naming of restriction enzymes the first letter is derived from genus name and next two letters from the species name of the prokaryotic cell from where the enzymes are extracted
- 159 **(d)**  
A molecule of DNA can be cut into fragments by

the enzyme restriction endonucleases. These fragments of DNA can be separated by a technique of gel electrophoresis. It is a technique used for the separation of substances of different ionic properties

- 163 **(c)**  
During extension, the enzymes *Taq* polymerase synthesizes the DNA segment between the primers. The two primers extend towards each other in order to copy the DNA segment typing between the two primers  
This step requires presence of deoxynucleoside triphosphate (*d*NTPs) and  $Mg^{2+}$  and occurs at  $72^{\circ}C$
- 164 **(c)**  
Both are true in the process for the isolation of DNA, after several treatments the purified DNA is precipitated by adding chilled ethanol. The bacterial/plant, animal cell is broken down by enzymes to release DNA, along with RNA, proteins, polysaccharide and lipids
- 165 **(c)**  
Bioreactors are vessels of large volumes (100-1000 litres) in which raw materials are biologically converted into specific products. It provides all the optimal conditions for achieving the desired product by providing optimal growth conditions like temperature, pH, substrate, salts vitamins and oxygen. Stirred-tank bioreactors are commonly used bioreactors. There are cylindrical with curved base to facilitate proper mixing of the contents. The stirrer mixes the contents and makes oxygen available throughout the bioreactor
- 166 **(a)**  
*Thermus aquaticus*.  
DNA polymerase which is stable at high temperature ( $>90^{\circ}C$ ) is required to carry out the synthesis of new DNA. The DNA polymerase like *Taq* polymerase is generally used in PCR reactions which is isolated from a bacterium *Thermus aquaticus*
- 169 **(c)**  
The first restriction endonuclease type II was isolated by Smith, Wilcox and Kelley from *Haemophilus influenza* bacterium. It was formed to cut DNA molecules at a particular point of recognizing a specific sequence of six base pairs, known as the recognition sequence
- 170 **(b)**  
In gel electrophoresis, the separated DNA

- fragments are visualized after staining the DNA with ethidium bromide followed by exposure to UV radiation
- 173 **(b)**  
In gel electrophoresis a molecule of DNA can be cut into fragments by the enzyme restriction endonuclease. DNA fragments move towards the anode according to their molecular size through the agarose gel  
The separated DNA fragments can be observed only after staining them with a solution of ethidium bromide. The bright orange coloured bands of DNA can be seen only under UV light. These bands of DNA fragments are cut out from the gel and extracted by using convenient technique. This step is called elution
- 175 **(a)**  
DNA polymerase which is stable at high temperature ( $>90^{\circ}\text{C}$ ) is required to carry out the synthesis of new DNA. The DNA polymerase like *Taq* polymerase is generally used in PCR reactions which is isolated from a bacterium *Thermus aquaticus*
- 176 **(a)**  
Most sensitive technique to detect malignant cell in non-hodgkins lymphoma is polymerase chain reaction. In recent times, PCR is being used in the detection of HIV (Virus of AIDS)
- 179 **(c)**  
The Pribnow box (also known as the Pribnow – Schaller box) is the sequence TATAAT of six nucleotides that is an essential part of a promoter site on DNA for transcription to occur in bacteria
- 187 **(a)**  
Gene gun method was first developed by Prof. Stanford and coworkers at Cornell University, USA in 1987. This method is used to introduce foreign DNA into host cell
- 188 **(c)**  
During extension, the enzyme DNA polymerase synthesizes the DNA segment between the primers. DNA polymerase is a heat stable enzyme
- 191 **(c)**  
After the formation of the product in the bioreactors, it undergoes through some processes before a finished product to be ready for marketing. The processes include (i) separation and (ii) purification of products, which are collectively called the downstream processing
- 192 **(a)**  
The stirred-tank bioreactor is well suited for large-scale production of protein of enzyme by using microbial plant/animal/human cells
- 193 **(a)**  
A-DNA is vector/plasmid DNA and B-is foreign DNA.  
C-The restriction enzyme that recognizes this palindrome-*Eco* RI  
D-The enzyme that can link these two DNA fragment-DNA ligase
- 194 **(c)**  
Restriction endonuclease was isolated for the first time by W Arber in 1962 in bacteria. They are called molecular scissors or biological scissors. In 1978 Arber, Smith and Nathan were awarded the Nobel Prize for the discovery of restriction endonuclease
- 195 **(b)**  
In genetic engineering *r*DNA technology is applied to several biotechnological processes for obtaining particular biochemical improvement of genetic make up of an organism and fighting genetic defects
- 197 **(d)**  
Primer and DNA polymerase.  
PCR is a technique of synthesizing multiple copies of the desired gene or (DNA) *in vitro*. The basic requirement of PCR are DNA template, two nucleotide primers and enzyme (DNA polymerase)
- 198 **(b)**  
An antibiotics resistance gene in a vector usually helps in the selection of transformed cell
- 200 **(c)**  
Bioreactors are considered as vessels in which raw materials are biologically converted into specific products by microbes, plant and animal cells and or their enzymes. Small volume cultures can not give large quantities of the products. Large scale production (100-1000 L) of the products is carried out in bioreactors. A bioreactor provides the optimal conditions for obtaining the desired product by providing optimum growth conditions such as temperature, pH, substrate, vitamins, oxygen and salts. In the sparged stirred tank bioreactor, sterile air bubbles are sparged. The surface area for oxygen transfer is increased
- 203 **(a)**  
*Vent* polymerase enzyme used in PCR is isolated from *Thermococcus litoralis*
- 211 **(b)**

- A stirred-tank bioreactor is more advantageous, than shake flasks. It has an agitator system to mix the contents properly, an oxygen delivery system to make availability of oxygen, a foam control system, a temperature control system, a pH control system and a sampling port to withdraw the small volumes of the culture periodically
- 212 **(b)**  
During gene cloning plasmid is called gene taxi. Molecular biologists add desired gene desired gene to plasmids, then insert the new plasmid with the added gene into a living bacterium
- 214 **(c)**  
Both are true. Copy number is defined as the number of copies of vectors present in a cell. It varies from 1-100 copies per cell
- 219 **(d)**  
Availability of thermostable DNA polymerase. DNA polymerase which is stable at high temperature (>90°C) is required to carry out the synthesis of new DNA. The DNA polymerase like *Taq* polymerase is generally used in PCR reactions which is isolated from a bacterium *Thermus aquaticus*
- 221 **(c)**  
Stanley Cohen and Herbert Boyer generated first recombinant DNA molecule by combining a gene from a bacterium with plasmid of *Escherichia coli*
- 230 **(c)**  
Thermophilic bacterium.  
Thermostable enzymes '*Taq* and *Vent*' isolated from thermophilic bacteria are DNA polymerase *Taq* polymerase, isolated from a *Thermophilic bacterium*, *Thermus aquaticus* and *vent* polymerase, isolated from a thermophilic bacterium *Thermococcus litoralis*
- 232 **(b)**  
*Agrobacterium tumefaciens* is used as a best genetic vector in plants
- 233 **(d)**  
*Plants in comparison to animals are more rapidly manipulated by genetic engineering reasons are*  
(i) Totipotency (having the ability to differentiate into all cell types) shown by plant cells  
(ii) Single somatic cell can regenerate a whole plant body  
(iii) Genetic engineering is supplemented with plant tissue culture techniques
- 237 **(b)**  
Vector is a plasmid or virus DNA used to introduce genes into a host cell, where the genes may be amplified (gene cloning) or otherwise manipulated
- 240 **(a)**  
Digestion with restriction enzyme  
↓  
Electrophoresis  
↓  
Ethidium bromide  
↓  
Radioactive probe  
↓  
X-ray film
- 241 **(d)**  
*amp<sup>R</sup>* (amplification resistance gene) and *tet<sup>R</sup>* (tetracycline resistance gene) are antibiotic resistance genes
- 244 **(d)**  
Restriction endonucleases cleave DNA molecules only at specific nucleotide sequence called restriction sites. DNA ligase enzymes is used to joins bits of DNA
- 248 **(a)**  
Mobile genetic element is broadly any genetic element capable of moving itself, with or without duplication, from one site in a genome to another. Mobile genetic elements include plasmids, viruses, transposable genetic elements (transposons), short interspread elements, pathogenicity islands and so on. The term 'transposon' was introduced **RW Hedges** and **AE Jacob** in 1974, 'controlling elements' or jumping genes, discovered by **Barbara McClintock** (1950) in maize
- 251 **(b)**  
Special sequence in the DNA recognized by restriction endonuclease is called palindromic nucleotide sequence.  
Restriction endonuclease recognizes palindromic sequences in DNA and cuts them  
The palindromes in DNA are base pair sequences that are the same when read forward (left to right) or backward (right to left) from a central axis of symmetry  
For example  
(i) 5' - G A A T T C - 3'  
3' - C T T A A G - 5'  
(ii) 5' - G G A T C C - 3'  
3' - C C T A G G - 5'
- 253 **(b)**  
Identification of DNA with desirable gene

↓

Introduction of DNA into host to form recombinant DNA

↓

Maintenance of DNA in host and gene cloning

↓

Gene transfer

254 (b)

*Recombinant DNA technology involved the following steps*

- (i) Isolation of DNA
- (ii) Fragmentation of DNA by restriction endonucleases
- (iii) Isolation of a desired DNA fragment
- (iv) Amplification of the gene of interest
- (v) Ligation of the DNA fragment into a vector
- (vi) Insertion of recombinant DNA into the host
- (vii) Culturing the host cells on a suitable medium at a large scale
- (viii) Extraction of the desired gene product
- (ix) Downstream processing of the products as finished product, ready for marketing

258 (c)

- A - Competency
- B - Calcium
- C - microinjection method

262 (a)


The most important feature in a plasmid to be used as a vector is origin of replication (*Ori*). Origin of replication is a specific sequence of DNA bases which is responsible for initiating replication. A prokaryotic DNA has a single origin of replication while eukaryotic DNA may have more than one origin of replication

263 (c)

DNA gyrase, the enzyme that participates in the process of DNA replication is a type of DNA topoisomerase

265 (d)

Three types of 'biological tool' are used in the formation of recombinant DNA



- (ii) Cloning vectors (vehicle vectors)
- (iii) Complementary host (for transformation with recombinant DNA)

268 (d)

In recent times PCR is being used in the detection of HIV (Virus of AIDS). By using PCR phenylketonuria, muscular dystrophy, sickle-cell anaemia, hepatitis, chlamydia and tuberculosis also can be diagnosed

269 (a)

Agarose is extracted from sea weeds. It is a polysaccharide. In gel electrophoresis, DNA fragments separate according to size through the pores of agarose gel

270 (d)

DNA polymerase remains active at high temperature. Usually *Taq* DNA polymerase, isolated from a thermophilic bacterium *Thermus aquaticus*, is used in most of the cases

272 (a)

*The science of biotechnology is based mainly on two core technologies*

- (i) **Genetic engineering**, which is the manipulation of genes by man. It includes techniques to alter the nature of genetic material (DNA and RNA), to introduce these into host organisms and thus, change the phenotype of the host organism
- (ii) **Biochemical engineering**, *i.e.*, processes that help the growth of desired microbe/eukaryotic cell in large quantities in a sterile medium for the manufacture and multiplication of biotechnological product

273 (c)

Each bioreactor has a cylindrical stirred-tank to facilitate the mixing of contents. The stirrer provides facility of mixing the contents as well as availability of oxygen throughout the process

275 (b)

*Taq* DNA polymerase is a thermostable enzyme, isolated from a *Thermophilic bacterium, Thermus*

*aquaticus*

278 (a) A sparged stirred-tank bioreactor being stirred at 200 RPM

280 (b) The Polymerase Chain Reaction (PCR) is a technique by which small samples DNA can be quickly amplified. Starting with only one gene sized pieces of DNA, this technique is used to make literally billions of copies in only a few hours

283 (b) Ligase catalyse the formation of bonds between P = O

284 (c) A probe is radioactively labelled ( $P^{32}$ ) nucleic acid (20-40 nucleotide long) with a short sequence complementary to at least one part of the desired DNA gene

285 (d) VNTRs were an important sources of RFLP genetic markers used in linkage analysis of genomes. VNTRs have become essential to forensic crime investigations, *via* DNA fingerprinting

286 (a) Isolation of restriction endonucleases by **Nathans** and **Smith** (1970) made it possible to cut DNA at specific sites. Restriction enzyme can cut both strains of DNA when foreign nucleotides are introduced in the cell. They cleave DNA to generate a nick with a 5' phosphoryl and 3' hydroxyl terminus

287 (a) Largest DNA bands will be at (A) and smallest DNA bands will be at (B) because in this DNA is move according to their size in agarose small DNA fragment will have small resistant so this fragment move to long distance as compared to large DNA fragment

289 (d) The technique of fingerprinting was initially developed by Alec Jeffreys. He used a satellite DNA as probe that shows very high degree of polymorphism. It was called as Variable Number of Tandem Repeats (VNTRs)

290 (c) *Vent* polymerase and *pfu* polymerase both

291 (d) A single PCR amplification cycle involves three

basic steps; denaturation, annealing and extension (polymerization)

Denaturation – Melting of target DNA


Annealing – Join

Extension – Polymerisation

295 (c) A-Bacteria, B-Virus, C-Cosmid

296 (a) The DNA fragments are seen as orange coloured bands. The separated bands of DNA are cut out and extracted from the gel piece. This step is called elution

298 (a) Simple stirred-tank bioreactor, sparged stirred-tank.



A-Simple stirred-tank bioreactor for continuous culture.

B-Sparged stirred-tank bioreactor through, which sterile (free from any germs) air bubbles are sparged

**Bioreactor** (fermenters) Bioreactors are considered as vessels in which raw materials are biologically converted into specific products by microbes, plant and animal cells and/or their enzymes. Small volume cultures cannot give large quantities of the products. Large scale production (100-1000 L) of the products in carried out in **bioreactors**. A bioreactor provides the optimal condition for obtaining the desired product by providing optimum growth conditions such as temperature, Ph, substrate, vitamins, oxygen and salts

**Types of Bioreactors** The most commonly used



bioreactors are of **stirring type**. Stirring type bioreactors are (i) **Simple stirred tank bioreactors** and (ii) **Sparged stirred-tank bioreactor** as shown in figure. In the sparged stirred-tank bioreactor, sterile air bubbles are sparged. The surface area for oxygen transfer is increased

300 (c)

DNA being a hydrophilic molecule can not pass through cell membranes. Therefore, the bacteria should be made competent to accept the DNA molecule

In this case the cell is treated with specific concentration of a divalent cation such as calcium to increase pore size in cell wall


The cells are incubated with recombinant DNA on ice, followed by placing them briefly at 42°C and then putting it back on ice. This is called heat shock treatment. The bacteria now takes up the recombinant DNA

303 (c)

DNA fingerprinting technique is very useful in solving disputed parentage cases and forensic cases. DNA fingerprinting are obtained from RFLP and VNTR (satellite DNA) analysis of blood, hair or other material found the place of crime

304 (a)

A molecule of DNA can be cut into fragments by the enzyme restriction endonucleases. These fragments of DNA can be separated by a technique of gel electrophoresis. In this process the smallest segment of DNA travel towards anode (+ve electrode), farthest away from the wells



306 (d)

The structure involved in genetic engineering is plasmid. Plasmids were discovered by William Hays and Joshua Lederberg (1952). These are extrachromosomal, self-replicating usually circular, double-stranded DNA molecules found naturally in many bacteria and also in some yeast

307 (b)

After the cutting of DNA by restriction enzymes fragments of DNA are formed. Separation of DNA fragments according to their size or length is done by a technique called gel electrophoresis developed by **A Tiselius** in 1937

308 (a)

In gel electrophoresis, the sample DNA is cut into fragments by restriction endonucleases

311 (a)

A-Denaturation - 94°C

B-Annealing - 40° – 60°C

C-Extension - 72°C


323 (c)

Genetic engineering

325 (d)

The separated DNA fragments can be seen only after staining the DNA with a compound known as ethidium bromide (E + Br) followed by exposure to UV radiation as bright orange coloured bands

330 (c)



A-Simple stirred-tank bioreactor for continuous culture.

B-Sparged stirred-tank bioreactor through, which sterile (free from any germs) air bubbles are sparged

**Bioreactor** (fermenters) Bioreactors are considered as vessels in which raw materials are biologically converted into specific products by microbes, plant and animal cells and/or their enzymes. Small volume cultures cannot give large quantities of the products. Large scale production (100-1000 L) of the products in carried out in **bioreactors**. A bioreactor provides the optimal condition for obtaining the desired product by providing optimum growth conditions such as temperature, Ph, substrate, vitamins, oxygen and salts

**Types of Bioreactors** The most commonly used

bioreactors are of **stirring type**. Stirring type bioreactors are (i) **Simple stirred tank bioreactors** and (ii) **Sparged stirred-tank bioreactor** as shown in figure. In the sparged stirred-tank bioreactor, sterile air bubbles are sparged. The surface area for oxygen transfer is increased

332 **(d)**

A variety of cell types are used as a donor in

recombinant DNA technology

333 **(b)**

A-DNA; B-Bacteria