

PHYSICS

Q. 1 A calorimeter contain 20 gm of water at 20 °C. The water equivalent of Calorimeter is 10 gm. The amount of heat required to raise the temperature of water from 20 °C to 60 °C (Sp. heat of water = 1cal/gm-°C)

- (A) 1000 cal (B) 800 cal
(C) 400 cal (D) 1200 cal **[D]**

Sol. $Q = 30 \times 1 \times 40$
 $= 1200 \text{ cal.}$

Q. 2 If two sphere having same radius but the ratio of their specific heat is 2 : 1 and the ratio of their density is 1 : 4 then the ratio of their heat capacity is -

- (A) 2 : 1 (B) 1 : 2
(C) 1 : 1 (D) 1 : 4 **[B]**

Sol. $\frac{C_1}{C_2} = \frac{\rho_1 V_1 S_1}{\rho_2 V_2 S_2}$
 $= \frac{2}{1} \times \frac{1}{4} = \frac{1}{2}$

Q. 3 Ice at 0°C is added to 200gm of water initially at 70°C in a vacuum flask. When 50 gm of ice has been added and has all melted, the temperature of flask and contents is 40°C, When a further 80 gm of ice is added and has all melted, the temperature of whole becomes 10°C. Neglecting heat lost to surroundings the latent heat of fusion of ice is :

- (A) 80 cal/gm (B) 90 cal/gm
(C) 70 cal/gm (D) 540 cal/gm **[B]**

Sol. According to principle of calorimetry,
 $ML_F + Ms \Delta T = (ms\Delta T)_{\text{water}} + (ms\Delta T)_{\text{flask}}$
 $50L_f + 50 \times 1 \times (40 - 0)$
 $= 200 \times 1 \times (70 - 40) + W (70 - 40)$
or $50L_f + 200 = (200 + W) 30$
or $5L_f = 400 + 3W$ (i)
Now the system contains (200 + 50) gm of water at 40°C, so when further 80 gm of ice is added.
 $80L_f + 80 \times 1 \times (10 - 0)$
 $= 250 \times 1 \times (40 - 10) + W (40 - 10)$
or $80L_f + 800 = (250 + W) 30$
or $80L_f = 670 + 3W$ (ii)
Solving equation (i) and (ii),
 $L_f = 90 \text{ cal/gm}$ and $W = \frac{50}{3} \text{ gm}$

Q.4 Heat is absorbed by a body but its temperature does not change. Which of the following statements explains the phenomenon ?

- (A) Only kinetic energy of vibration increases
(B) Only potential energy of intermolecular force field increases
(C) No increase in internal energy takes place
(D) Increase in kinetic energy is balanced by decrease in potential energy

Sol. [B]
Only potential energy will increase as
 $U = K.E + P.E$

Q.5 If specific heat of a substance is infinite, it means-

- (A) Heat is given out
(B) Heat is taken in
(C) No change in temperature takes place whether heat is taken in or given out
(D) All of the above

Sol. [C]
 $Q = m.c.\Delta\theta \Rightarrow c = \frac{Q}{m.\Delta\theta}$; when $\Delta\theta = 0 \Rightarrow c = \infty$

Q.6 2 gm ice at 0°C is mixed with 5 gm steam at 100°C in a calorimeter of negligible heat capacity. At equilibrium the calorimeter will contain -

- (A) $\frac{13}{3}$ gm steam and water at 100°C
(B) $\frac{1}{3}$ gm ice and water at 0°C
(C) only water at temperature 67°C
(D) none of these

Sol. [A]
Heat given by steam = Heat taken by ice.

Q.7 A copper block of mass 2 kg is heated to a temperature of 500°C and then placed in a large block of ice at 0°C. What is the maximum amount of ice that can melt ? The specific heat of copper is 400 J kg⁻¹ °C⁻¹ and latent heat of fusion of ice is

- $3.5 \times 10^5 \text{ J kg}^{-1}$:
(A) $\frac{4}{3}$ kg (B) $\frac{6}{5}$ kg

(C) $\frac{8}{7}$ kg (D) $\frac{10}{9}$ kg [C]

Sol. $2 \times 400 \times 500 = m \times 3.5 \times 10^5$
 $4 = m \times 3.36 \Rightarrow m = \frac{4}{3.36} \text{ kg} = \frac{40}{35} = \frac{8}{7} \text{ kg}$

Q.8 Evaporation of perspiration is an important mechanism for temperature control in warm-blooded animals. What mass of water must evaporate from the surface of an 80 kg human body to cool it 1°C? The specific heat capacity of the human body is approximately 1 cal g⁻¹(°C)⁻¹, and the latent heat of vaporisation of water at body temperature (37°C) is 577 cal g⁻¹.

- (A) 139 g (B) 128 g
 (C) 110 g (D) 109 g

Sol. [A]
 $80 \times 1000 \times 1 = m \times 577$
 $m = \frac{8000}{577} = 139 \text{ gm}$

Q.9 A stream of photons impinging normally on a completely absorbing screen in vacuum exerts a pressure P. If I is the irradiance then -

- (A) $P = \frac{2I}{c}$ (B) $P = Ic$ (C) $P = \frac{I}{c}$ (D) $P = 2Ic$

Sol.[C] Pressure = $\frac{F}{A} = \frac{P \times n}{At} = \frac{h}{\lambda} \times \frac{n}{At}$

Irradiance = Intensity = $I = \frac{nhc}{\lambda At}$

Q.10 5 g of steam at 100°C is passed into 6g of ice at 0°C. If the latent heats of steam and ice are 540 cal/g and 80 cal/g, then the final temperature is :

- (A) 0°C (B) 50°C
 (C) 30°C (D) 100°C [D]

Sol. $6 < 3$
 So, temp of mixture is 100°C

Q.11 The temperatures of equal masses of three different liquids A, B and C are 12°C, 19°C and 28°C respectively. The temperature when A and B are mixed is 16°C. When B and C are mixed it is 23°C. The temperature when A and C are mixed will be -

- (A) 15°C (B) 18.2°C
 (C) 20.25°C (D) 24.5°C

Sol. [C] $m_{SA} \times 4 = m_{SB} \times 5 \Rightarrow 4s_A = 3s_B$ (i)

$m_{SB} \times 4 = m_{SC} \times 5 \Rightarrow 4s_B = 5s_C$ (ii)

By (i) & (ii)

$16s_A = 15s_C$ (iii)

$m_{SA}(\theta - 12) = m_{SC}(28 - \theta)$ (iv)

By (i) & (iv)

$\theta = 20.25^\circ\text{C}$

Q.12 The temperature of a body on Kelvin scale is found to be x K. When it is measured by a Fahrenheit thermometer, it is found to be x°F. Then x is :

- (A) 301.25 (B) 574.25
 (C) 313 (D) 40 [B]

Sol. $\frac{x-273}{373-273} = \frac{f-32}{212-32}$

$\Rightarrow x-273 = \frac{5}{9}(x-32)$

$\Rightarrow 9x-9 \times 273 = 5x-160 \Rightarrow x = 574.25$

Q.13 A substance of mass m kg requires a power input of P watts to remain in the molten state at its melting point. When the power is turned off, the sample completely solidifies in time t sec. What is the latent heat of fusion of the substance?


- (A) $\frac{Pm}{t}$ (B) $\frac{Pt}{m}$ (C) $\frac{m}{Pt}$ (D) $\frac{t}{Pm}$

Sol.[B] Heat lost in t sec = mL or heat lost per sec = $\frac{mL}{t}$.

This must be the heat supplied for keeping the substance in molten state per sec.

$\therefore \frac{mL}{t} = P$ or $L = \frac{Pt}{m}$

Q.14 Which of the curves in figure represents the relation between Celsius and Fahrenheit temperature-



- (A) 1 (B) 2 (C) 3 (D) 4

Sol. [A] $\frac{C}{5} = \frac{F-32}{9} \Rightarrow C = \left(\frac{5}{9}\right)F - \frac{20}{3}$. Hence graph

between °C and °F will be a straight line with positive slope and negative intercept.

Q.15 Two liquids A and B are at 32°C and 24°C. When mixed in equal masses the temperature of the mixture is found to be 28°C. Their specific heats are in the ratio of -

- (A) 3 : 2 (B) 2 : 3
(C) 1 : 1 (D) 4 : 3

Sol. [B]

$$\frac{H_1}{H_2} = \frac{R_1^3}{R_2^3} \times \frac{\rho_1}{\rho_2} \times \frac{s_1}{s_2}$$

Q.16 Boiling water is changing into steam. At this stage the specific heat of water is-

- (A) < 1 (B) ∞ (C) 1 (D) 0

Sol. [B]

$$C = \frac{Q}{m \Delta \theta}; \text{ as } \Delta \theta = 0, \text{ hence } c \text{ becomes } \infty.]$$

Q.17 A 1 kg cube of ice of volume 1000 ml at temperature 0°C is placed in a cylinder of cross sectional area 200 cm². If ice melts only on top surface by absorbing heat from environment at a rate 80 cal per second. The time when ice lifts from base of cylinder is -

- (given density of water 1000 kg/m³)
(A) 4 min. 10 sec. (B) 8 min. 20 sec.
(C) will never lift (D) more than 10 min.

Sol. [B] Ice will float when completely immersed.

Q.18 In an energy recycling process, X g of steam at 100°C becomes water at 100°C which converts Y g of ice at 0°C into water at 100°C. The ratio of X/Y will be -

- (A) 1/3 (B) 2/3 (C) 3 (D) 2

Sol. [A]

$$\begin{aligned} \text{Sp. heat of vaporization} &= 22.68 \times 10^5 \text{ J/kg} \\ &= X \times 10^{-3} \times 22.68 \times 10^5 \\ &= Y \times 10^{-3} \times 3.36 \times 10^5 + Y \times 10^{-3} \times 4200 \times 100 \end{aligned}$$

$$\therefore \frac{X}{Y} = \frac{1}{3}$$

Q.19 If a thermometer reads freezing point of water as 20°C and boiling point as 150°C, how much thermometer read when the actual temperature is 60°C-

- (A) 98°C (B) 110°C (C) 40°C (D) 60°C

Sol. [A] Temperature on any scale can be converted into

$$\text{other scale by } \frac{x - \text{LFP}}{\text{UFP} - \text{LFP}} = \text{constant for all scales}$$

$$\frac{x - 20}{150 - 20} = \frac{60}{100} \Rightarrow x = 98^\circ\text{C}$$

Q.20 Liquids A and B are at 30°C and 20°C. When mixed in equal masses, the temperature of the mixture is found to be 26°C. Their specific heats are of ratio -

- (A) 3 : 2 (B) 1 : 1 (C) 2 : 3 (D) 4 : 3

Sol. [A] $m s_A (30 - 26) = m s_B (26 - 20)$

$$4s_A = 6s_B \Rightarrow \frac{s_A}{s_B} = \frac{3}{2}$$

Q.21 A body A of mass 0.5 kg and specific heat 0.85 is at a temperature of 60°C. Another body B of mass 0.3 kg and specific heat 0.9 is at a temperature of 90°C. When they are connected to a conducting rod, heat will flow from -

- (A) A to B
(B) B to A
(C) heat can't flow
(D) first from A to B, then B to A

Sol. [B] Heat flows from a body at higher temperature to a body at lower temperature. So heat will flow from B to A.

Q.22 10 gm of ice at 0°C is mixed with 10 gm steam at 100°C in a container of negligible heat capacity. Amount of steam in the mixture after some time will be ($S_w = 1 \text{ cal/gm}^\circ\text{C}$, $L_v = 540 \text{ cal/gm}$, $L_f = 80 \text{ cal/gm}$) -

- (A) 0 gm (B) $\frac{20}{3}$ gm
(C) $\frac{10}{3}$ gm (D) None of these [B]

Sol. Amount of steam required to convert all the ice in water at 100°C is $\frac{10}{3}$ gm.

Q.23 The ratio of densities of two substance is 2 : 3 and their specific heats are in the ratio 3 : 4. The ratio of their thermal capacities for unit volume is -

- (A) 1 : 1 (B) 1 : 2 (C) 2 : 1 (D) 8 : 9

Sol. [B] $\frac{c_1}{c_2} = \frac{m_1 S_1}{m_2 S_2} = \frac{\rho_1 v S_1}{\rho_2 v S_2} = \frac{2}{3} \times \frac{3}{4} = \frac{1}{2}$

Q.24 The absolute zero temperature in Fahrenheit is -

- (A) - 273°F (B) -32°F
(C) -460°F (D) -132°F

Sol. [C] $\frac{F-32}{9} = \frac{K-273}{5} \Rightarrow \frac{F-32}{9} = \frac{0-273}{5}$
 $\Rightarrow F = -459.4^{\circ}\text{F} = -460^{\circ}\text{F}$

- Q.25** Latent heat of ice is 80 cal g^{-1} and $J = 4.2 \text{ J cal}^{-1}$. A boy can melt 60 gram ice in 1 minute by chewing. His power is –
 (A) 4800 W (B) 336 W
 (C) 1.33 W (D) 0.75 W [B]

- Q.26** Heat required to convert one gram of ice at 0°C into steam at 100°C is –
 (A) 100 cal (B) 0.01 kcal
 (C) 716 cal (D) 1 kcal [C]

- Q.27** 80 gm of water at 30°C is poured on a large block of ice at 0°C . The mass of ice that melts is –
 (A) 160 gm (B) 80 gm
 (C) 40 gm (D) 30 gm

Sol. [D]
 Heat released by water
 $\Delta Q = 80 \times 1 \times 30 = 2400 \text{ cal} \dots\dots(i)$
 Mass of Ice melt
 $2400 = m \times 80 \quad [\Delta Q = mL]$
 $\therefore m = \frac{2400}{80} = 30 \text{ gm}$

- Q.28** One gm of ice at 0°C is added to 5gm of water at 10°C . If the latent heat is 80 cal/gm , the final temperature of the mixture is –
 (A) 5°C (B) 0°C
 (C) -5°C (D) none of these [B]

- Q.29** One gm of ice is mixed with one gm of steam. After thermal equilibrium is reached, the temperature of mixture is –
 (A) 100°C (B) 55°C
 (C) 75°C (D) 0°C [A]

- Q.30** Which one of the following would raise the temperature of 20 gm of water at 30°C most when mixed with :
 (A) 20 gm of water at 40°C
 (B) 40 gm of water at 35°
 (C) 10 gm of water at 50°C
 (D) 4 gm of water at 80°C [D]

Sol. $T = \frac{m_1s_1T_1 + m_2s_2T_2}{m_1s_1 + m_2s_2}$,

T is maximum, If we take 4 gm of water at 80°C

- Q.31** When 300 J of heat is added to 25 gm of sample of a material its temperature rises from 25°C to 45°C . The thermal capacity of the sample and specific heat of the material are respectively given by –
 (A) $15 \text{ J}^{\circ}\text{C}$, $600 \text{ J/kg}^{\circ}\text{C}$
 (B) $600 \text{ J}^{\circ}\text{C}$, $15 \text{ J}^{\circ}\text{C-kg}$
 (C) $150 \text{ J}^{\circ}\text{C}$, $60 \text{ J/kg}^{\circ}\text{C}$
 (D) none of these [A]

- Q.32** 70 calories are required to raise the temperature of 2 moles of an ideal gas at constant pressure from 30°C to 35°C . The amount of heat required (in calories) to raise the temperature of the same gas through the same range (30°C to 35°C) at constant volume is –
 (A) 30 (B) 50
 (C) 70 (D) 90 [B]

- Q.33** The heat required to convert one kg of ice to water is –
 (A) latent heat of vaporisation (B) specific heat
 (C) molar specific heat (D) latent heat of fusion [D]

- Q.34** A liquid of mass m and specific heat C is heated to a temperature 2T. Another liquid of mass (m/2) and specific heat 2C is heated to a temperature 2T. If these two liquids are mixed, the resulting temperature of the mixture is –
 (A) $(2/3)T$ (B) $(8/5)T$
 (C) $(3/5)T$ (D) $(3/2)T$ [D]

- Q.35** 10 gm of ice at -20°C is added to 10gm of water at 50°C . $S_{\text{water}} = 1 \text{ cal/gm}^{\circ}\text{C}$, $S_{\text{ice}} = 0.5 \text{ cal/gm}^{\circ}\text{C}$. $L_{\text{f,ice}} = 80 \text{ cal/gm}$. The resulting temperature is –
 (A) -20°C (B) 15°C
 (C) 0°C (D) 50°C [C]

Sol. Some ice will left in the mixture.

- Q.36** Liquids A and B are at 30°C and 20°C . When mixed in equal masses, the temperature of the

mixture is found to be 26°C. Their specific heats are in the ratio of -

- (A) 3 : 2 (B) 1 : 1
(C) 2 : 3 (D) 4 : 3 [A]

Q.37 The densities of two materials X and Y are in the ratio 1 : 3. Their specific heats are in the ratio 3 : 1. If we take same volumes of the two substances, the ratio of their thermal capacities will be :

- (A) 1 : 1 (B) 1 : 3
(C) 1 : 6 (D) 1 : 9 [A]

Sol. $\frac{C_1}{C_2} = \frac{P_1 V s_1}{P_2 V s_2} = \frac{1}{3} \times \frac{3}{1} = 1 : 1$

Q.38 Two tanks A and B contains water at 30°C and 80°C respectively, calculate the amount of water that must be taken from each tank to prepare 40kg of water at 50°C -

- (A) 24 kg, 16 kg (B) 16 kg, 24 kg
(C) 20 kg, 20 kg (D) 30 kg, 10 kg [A]

Sol. $m_1 \times 1 (50 - 30) = m_2(80 - 50) \dots(i)$
 $m_1 + m_2 = 40 \text{ kg} \dots(ii)$

Q.39 Two liquids are at temperature 20°C and 40°C. When same mass of both of them is mixed, the temperature of the mixture is 32°C. What is the ratio of the their specific heats ?

- (A) 1/3 (B) 2/3
(C) 1/5 (D) 2/5 [B]

Sol. $m s_A(32 - 20) = m s_B(40 - 32)$
 $12 s_A = 8 s_B$
 $\frac{s_A}{s_B} = \frac{2}{3}$

Q.40 The minimum mass of ice at 0°C required to just condense m grams of steam at 100°C is: (latent heat of steam and ice are 540 cal/g and 80 cal/g respectively)

- (A) 5.4 m (B) 7 m
(C) 3 m (D) 6.75 m [C]

Sol. Let m_1 is mass of ice required
then, $m_1 \times 80 + m_1 \times 1 \times 100 = 540 m$
 $180 m_1 = 540 m \Rightarrow m_1 = 3m$

Q.41 Two bodies of specific heats S_1 and S_2 having same heat capacities are combined to form a single composite body. Specific heat of composite body is -

- (A) $S_1 + S_2$ (B) $\frac{S_1 + S_2}{2}$
(C) $\frac{2S_1 S_2}{S_1 + S_2}$ (D) $\frac{1}{S_1} + \frac{1}{S_2}$ [C]

Sol. $m_1 s_1 = m_2 s_2 \dots(1)$
 $m_1 s_1 + m_2 s_2 = (m_1 + m_2) s \dots(2)$
 $s = \frac{m_1 s_1 + m_2 s_2}{m_1 + m_2} = \frac{2 m_1 s_1}{m_1 + m_2} (\because m_1 s_1 = m_2 s_2)$
 $= \frac{2 m_1 s_1}{m_1 + \frac{m_1 s_1}{s_2}} = \frac{2 s_1 s_2}{s_1 + s_2}$

Q.42 The steam point and ice point of a mercury thermometer are marked as 80° and 10°. At what temperature on centigrade scale the reading of this thermometer will be 59° ?

- (A) 70° C (B) 60° C
(C) 80° C (D) None of these [A]

Sol. $\frac{T' - 10}{80 - 10} = \frac{T_C}{100}; T' = 59^\circ$

Q.43 300 g of water at 25°C is added to 100g of ice at 0°C. The amount of ice melts is -

- (A) 6.25 g
(B) 93.75 g
(C) 100 g
(D) none of the above [B]

Sol. heat release by water = $m s d \theta$
 $= 300 \times 1 \times 25$
 $= 7500 \text{ Cal.}$
amount of Ice melts from this heat
 $dQ = mL$

$$m = \frac{dQ}{L} = \frac{7500}{80} = 93.75 \text{ g}$$

Q.44 A temperature difference of 5°C on Celsius scale corresponding to the following temperature difference in the Fahrenheit scale -

- (A) 9° (B) 41°
(C) 2.8° (D) 15° [A]

Sol. $\frac{C}{5} = \frac{F - 32}{9}$
 $\frac{\Delta C}{5} = \frac{\Delta F}{9}$
 $\Delta F = 9 \times \frac{5}{9} = 9^\circ$

Q.45 250 gm of water and equal volume of alcohol of mass 200 gm are replaced successively in the same calorimeter and cool from 60°C to 55°C in 130 sec and 67 sec respectively. If the water equivalent of calorimeter is 10 gm. The specific heat of alcohol in cal/gm°C is -

- (A) 1.30 (B) 0.67
(C) 0.62 (D) None of these [C]

Sol. Hint : Rate of heat loss are equal in both case.

Q.46 2 kg ice at - 20°C is mixed with 5 kg water at 20°C in an insulating vessel having negligible heat capacity. Calculate the final mass of water remaining in container.

Given sp. heat water = 4.186 kJ K⁻¹ kg⁻¹

sp. heat Ice = 2.092 kJ K⁻¹ kg⁻¹

Latent heat of fusion of ice = 334.7 kJ Kg⁻¹

- (A) 7 kg (B) 6 kg
(C) 4 kg (D) 2 kg [B]

Sol. Heat that can be given to ice by water

$$= 5 \text{ kg} \times 1 \times (20 - 0)$$

$$Q = 100 \text{ k cal.}$$

Energy required to raise temp of ice from - 20°C to 0°C

$$E = 2 \text{ kg} \times 0.5 \times (0 - (-20)) = 20 \text{ k cal.}$$

Heat available for = Q - E

Melting ice = 100 - 20 = 80 k cal.

L = 80 cal/gram

$$\text{Ice that can be melted} = \frac{80 \text{ k cal}}{80 \text{ cal}} = 1000 \text{ gram} = 1$$

kg.

Hence water left in container = 6 kg

Q.47 Water of volume 2 litre in a container is heated with a coil of 1 kW at 27°C. The lid of the container is open and energy dissipates at rate of 160 J/s. In how much time temperature will rise from 27°C to 77°C ?

[Given specific heat of water is 4.2 kJ/kg]

- (A) 8 min 20 s (B) 6 min 2 s
(C) 7 min (D) 14 min [A]

Sol. Energy gained by water (in 1 s)

= energy supplied - energy lost

$$= (1000 \text{ J} - 160 \text{ J}) = 840 \text{ J}$$

Total heat required to raise the temperature of water from 27°C to 77°C is $ms\Delta\theta$.

Hence, the required time

$$t = \frac{ms\Delta\theta}{\text{rate by which energy is gained by water}}$$

$$= \frac{2 \times (4.2 \times 10^3) \times 50}{840}$$

$$= 500 \text{ s}$$

$$= 8 \text{ min } 20 \text{ s}$$

Q.48 One gram of ice is mixed with one gram of steam. After thermal equilibrium, the temperature of the mixture is -

- (A) 0°C (B) 100°C
(C) 55°C (D) 80°C [B]

Sol. Heat release when 1gm steam convert into water at 100°C is = mL_{cond}

$$= 1 \times 536 \text{ cal.}$$

heat required to convert ice into water

at 100°C = $mL_{\text{fus.}} + mS_w\theta$

$$= (1 \times 80) + 1 \times 1 \times 100$$

$$= 180 \text{ cal}$$

Since 536 cal is greater than 180 cal so partial condensation will take place and the final temperature will be 100°C

Q.49 If x grams of steam at 100°C becomes water at 100°C which converts y grams of ice at 0°C into water at 100°C, then the ratio x/y will be -

- (A) $\frac{1}{3}$ (B) $\frac{27}{4}$ (C) 3 (D) $\frac{4}{27}$

Sol. [A] $x \times 540 = y \times 80 + y \times 1 \times 100$

$$\Rightarrow 540x = 180y \quad \text{or} \quad \frac{x}{y} = \frac{1}{3}$$

Q.50 Ice point and steam point on a particular scale reads 10° and 80° respectively. The temperature on °F scale when temperature on new scale is 45° is -

- (A) 50°F (B) 112°F
(C) 122°F (D) 138°F [C]

Sol. Relation between the two scales

$$\frac{t-10}{80-10} = \frac{F-32}{180}$$

$$F = \frac{18}{7} (t-10) + 32$$