



# COORDINATION COMPOUNDS

## CHEMISTRY


### Single Correct Answer Type




- The IUPAC name of  $\text{Na}_3[\text{Co}(\text{ONO})_6]$  is:
  - Sodium cobaltinitrite
  - Sodium hexanitritocobaltate(III)
  - Sodium hexanitrocobalt(III)
  - Sodium hexanitritocobaltate(II)
- $\text{CuSO}_4$  decolourises on addition of KCN, the product is:
  - $\text{Cu}(\text{CN})_4^{2-}$
  - $[\text{Cu}(\text{CN})_4]^{3-}$
  - $\text{Cu}(\text{CN})_2$
  - $\text{CuCN}$
- Exchange of coordination group by a water molecule in complex molecule results in:
  - Ionization isomerism
  - Ligand isomerism
  - Hydration isomerism
  - Geometrical isomerism
- The type of isomerism found in urea molecule is
  - Chain
  - Position
  - Tautomerism
  - None of these
- The IUPAC name of the compound  is
  - Butane-2-aldehyde
  - 2-methyl butanal
  - 2-ethyl propanal
  - None of the above
- Anisol is a product obtained from phenol by the reaction known as:
  - Coupling
  - Etherification
  - Oxidation
  - Esterification
- Which of the following is diamagnetic in nature?
  - $[\text{Fe}(\text{CN})_6]^{3-}$
  - $[\text{NiCl}_4]^{2-}$
  - $[\text{Ni}(\text{CO})_4]$
  - $[\text{MnCl}_4]^{2-}$
- Which is the strongest field ligand?
  - $\text{CN}^-$
  - $\text{NO}_2^-$
  - $\text{NH}_3$
  - en
- Nitrobenzene on reduction with Zn and aq.  $\text{NH}_4\text{Cl}$  gives:
  - Aniline
  - Nitrosobenzene
  - N*-phenyl hydroxylamine
  - Hydrazobenzene
- The IUPAC name of  $[\text{Co}(\text{NH}_3)_5 \text{ONO}]^{2+}$  ion is
  - Pentaammine nitrito cobalt (IV) ion
  - Pentaammine nitro cobalt (III) ion
  - Pentaammine nitrito cobalt (III) ion
  - Pentaammine nitro cobalt (IV) ion
- The compound which does not show paramagnetism is
  - $\text{NO}_2$
  - NO
  - $[\text{Ag}(\text{NH}_3)_2]\text{Cl}$
  - $[\text{Cu}(\text{NH}_3)_4\text{Cl}_2]$
- Which of the following is expected to undergo nitration more easily and readily to furnish the corresponding nitro derivatives employing the usual nitrating mixture?
  - $\text{C}_6\text{H}_6$
  - $\text{C}_6\text{H}_5\text{NO}_2$
  - $\text{C}_6\text{H}_5\text{CH}_3$
  - $\text{C}_6\text{H}_5 \cdot \text{CCl}_3$
- The number of unpaired electrons calculated in  $[\text{Co}(\text{NH}_3)_6]^{3+}$  and  $[\text{Co}(\text{F}_6)]^{3-}$  are
  - 4 and 4
  - 0 and 2
  - 2 and 4
  - 0 and 4
- The IUPAC name of  is

- a) 4-hydroxy amino benzene carboxylic acid      b) 4-(N-hydroxy) imino benzene carboxylic acid  
 c) 4-hydroxy imino cyclohexanoic acid      d) 4-(N-hydroxy) imino cyclohexane-1-carboxylic acid
15. The IUPAC name of the coordination compound  $K_2[Zn(OH)_4]$  is  
 a) Potassium tetrahydrozine (II)      b) Dipotassium tetrahydroxo(II)  
 c) Potassium tetrahydrozincate (II)      d) Potassium tetrahydrozincate (III)
16. Arrange in order of decreasing trend towards  $S_E$  reactions, Chlorobenzene, Benzene, Anilium chloride, Toluene:  
 I.      (II)      (III)      (IV)  
 a)  $II > I > III > IV$       b)  $III > I > II > IV$       c)  $IV > II < I > III$       d)  $I > II > III > IV$
17. Toluene is nitrated and the resulting product is reduced with tin and hydrochloric acid. The product so obtained is diazotised and then heated with cuprous bromide. The reaction mixture so formed contains:  
 a) Mixture of *o*- and *m*-bromotoluenes  
 b) Mixture of *o*- and *p*-bromotoluenes  
 c) Mixture of *o*- and *p*-dibromobenzenes  
 d) Mixture of *o*- and *p*-bromoanilines
18. A positive carbylamine test is given by:  
 a) *N,N*-dimethylaniline  
 b) 2,4-dimethylaniline  
 c) *N*-methyl-*o*-methylaniline  
 d) *p*-methyl benzylamine
19.  $CN^-$  is strong field ligand. This is due to the fact that  
 a) It carries negative charge  
 b) It is a pseudohalide  
 c) It can accept electrons from metal species  
 d) It forms high spin complexes with metal species.
20. Which of the following is not true for ligand metal complex?  
 a) Highly charged ligand forms strong bond  
 b) Greater the ionization potential of central metal, the stronger is the bond  
 c) Larger the permanent dipole moment of ligand, the more stable is the bond  
 d) Larger the ligand, the more stable is the metal-ligand bond
21. The nitration of nitrobenzene with fuming  $HNO_3$  will give:  
 a) TNB      b) 1,3-dinitrobenzene      c) Picric acid      d) 1,4-dinitrobenzene
22. A ligand can also be regarded as  
 a) Lewis acid      b) Bronsted base      c) Lewis base      d) Bronsted acid
23. The correct statement with respect to the complexes  $Ni(CO)_4$  and  $[Ni(CN)_4]^{2-}$  is  
 a) Nickel is in the same oxidation state in both  
 b) Both have tetrahedral geometry  
 c) Both have square planar geometry  
 d) Have tetrahedral and square planar geometry respectively
24. Which one of the following has lowest value of paramagnetic behaviour?  
 a)  $[Cr(CN)_6]^{3-}$       b)  $[Mn(CN)_6]^{3-}$       c)  $[Fe(CN)_6]^{3-}$       d)  $[Co(CN)_6]^{3-}$
25. In the reaction;



the products are:







26. An octahedral complex is formed when central metal atom undergoes hybridization amongst the....orbitals.

a)  $sp^3$

b)  $dsp^2$

c)  $sp^3d$

d)  $sp^3d^2$



28. Biological oxidation of C<sub>6</sub>H<sub>6</sub> taking place in body of dog, gives:

a) Benzoic acid

b) Toluic acid

c) Maleic acid

d) Muconic acid

29. Ammonia forms the complex ion [Cu(NH<sub>3</sub>)<sub>4</sub>]<sup>2+</sup> with copper ions in the alkaline solutions but not in acidic solutions. What is the reason for it?

a) In acidic solutions hydration protects copper ions

b) In acidic solutions protons coordinate with ammonia molecules forming NH<sub>4</sub><sup>+</sup> ions and NH<sub>3</sub> molecules are not available

c) In alkaline solutions insoluble Cu(OH)<sub>2</sub> is precipitated which is soluble in excess of any alkali

d) Copper hydroxide is an amphoteric substance

30. Which of the following has the highest molar conductivity in solution?

a) [Pt(NH<sub>3</sub>)<sub>6</sub>]Cl<sub>4</sub>

b) [Pt(NH<sub>3</sub>)<sub>5</sub>Cl]Cl<sub>3</sub>

c) [Pt(NH<sub>3</sub>)<sub>4</sub>Cl<sub>2</sub>]Cl<sub>2</sub>

d) [Pt(NH<sub>3</sub>)<sub>3</sub>Cl<sub>3</sub>]Cl

31. Which of the following is not *meta* directing group?

a) —SO<sub>3</sub>H

b) —NO<sub>2</sub>

c) —CN

d) —NH<sub>2</sub>

32. Which of the following is an organometallic compound?

a) Lithium methoxide

b) Lithium acetate

c) Lithium dimethylamine

d) Methyl lithium

33. Which among the following is very strong *o*-, *p*-directing group?

a) —Cl

b) —OR

c) —NH<sub>2</sub>

d) —NHR

34. The type of hybridisation in tetrahedral complexes of metal atom is

a)  $dsp^2$

b)  $d^2sp$

c)  $sp^3$

d)  $sp^2$

35. Chlorobenzene on heating with NaOH at 300°C under pressure gives:

a) Phenol

b) Benzaldehyde

c) Chlorophenol

d) None of these

36. The coordination number of Fe in  $[\text{Fe}(\text{CN})_6]^{4-}$ ,  $[\text{Fe}(\text{CN})_6]^{3-}$  and  $[\text{FeCl}_4]^-$  are respectively.  
 a) 2, 3, 3                      b) 6, 6, 4                      c) 6, 3, 3                      d) 6, 4, 6

37. Consider the following statements

- I. Chain and position isomerism are not possible together between two isomers  
 II. Tautomerism is a chemical phenomenon which is catalysed by acid as well as base  
 III. Tautomers are always metamers  
 IV. Tautomers are always functional isomers

Select the correct answer by using the codes given below

- a) Only III is correct                      b) III and IV are correct  
 c) I, II and III are correct                      d) I, II and IV are correct

38. What is the EAN of nickel in  $[\text{Ni}(\text{CN})_4]^{2-}$ ?

- a) 32                      b) 35                      c) 34                      d) 36



39. Which of the following alcohols is dehydrated most readily with conc.  $\text{H}_2\text{SO}_4$ ?

- a)  $p\text{-O}_2\text{NC}_6\text{H}_4\text{CH}(\text{OH})\text{CH}_3$   
 b)  $p\text{-ClC}_6\text{H}_4\text{CH}(\text{OH})\text{CH}_3$   
 c)  $p\text{-CH}_3\text{OC}_6\text{H}_4\text{CH}(\text{OH})\text{CH}_3$   
 d)  $\text{C}_6\text{H}_5\text{CH}(\text{OH})\text{CH}_3$

40. The compound having tetrahedral geometry is

- a)  $[\text{Ni}(\text{CN})_4]^{2-}$                       b)  $[\text{Pd}(\text{CN})_4]^{2-}$                       c)  $[\text{PdCl}_4]^{2-}$                       d)  $[\text{NiCl}_4]^{2-}$

41. Identify 'Z' in the change;



42. Which of the following is most acidic?

- a)  $p$ -cresol                      b)  $p$ -chlorophenol                      c)  $p$ -nitrophenol                      d)  $p$ -aminophenol

43. Benzoylacetato beryllium exhibit isomerism of the type

- a) Structural                      b) Geometrical                      c) Optical                      d) Conformational

44. Which one of the following has a square planar geometry?

(At. No. Fe=26, Co=27, Ni=28, Pt=78)

- a)  $[\text{CoCl}_4]^{2-}$                       b)  $[\text{FeCl}_4]^{2-}$                       c)  $[\text{NiCl}_4]^{2-}$                       d)  $[\text{PtCl}_4]^{2-}$

45. The number of ions formed on dissolving one molecule of  $\text{FeSO}_4(\text{NH}_4)_2\text{SO}_4 \cdot 6\text{H}_2\text{O}$  in water is:

- a) 4                      b) 5                      c) 3                      d) 6


46. A solution of potassium ferrocyanide would contains-ions

- a) 2                      b) 3                      c) 4                      d) 5

47. Which of the following is not considered as an organometallic compound?

- a) Grignard reagent                      b)  $cis$ -platin                      c) Zeise's salt                      d) Ferrocene


48. When phenol is reacted with  $\text{CHCl}_3$  and  $\text{NaOH}$  followed by acidification, salicylaldehyde is obtained. Which of the following species are involved in the above mentioned reaction as intermediates?





49. Number of geometrical isomers for the molecule








73. The complex ion which has no 'd'-electron in the central metal atom is :  
 a)  $[\text{MnO}_4]^-$                       b)  $[\text{Co}(\text{NH}_3)_6]^{3+}$                       c)  $[\text{Fe}(\text{CN})_6]^{3-}$                       d)  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$
74. The shape of cobalt hexaammine cation, which has its central cobalt atom surrounded by six ammonia molecules is:  
 a) Tetrahedral                      b) Octahedral                      c) Square planar                      d) Trigonal
75. Which ligand is capable of forming low spin as well as high spin complexes?  
 a) CO                      b)  $\text{NO}_2^-$                       c)  $\text{CN}^-$                       d)  $\text{NH}_3$
76.   
 The IUPAC name of \_\_\_\_\_ is  
 a) 7-ethyl-2, 4, 5, 6-tetramethyl-deca-1, 9-diene                      b) 7-ethyl-2, 4, 5, 6-tetramethyl-deca-1, 8-diene  
 c) 4-ethyl-4, 5, 6, 7-tetramethyl-deca-1, 9-diene                      d) 7-(1-propenyl)-2, 3, 4, 5-tetramethyl-non-1-ene
77. IUPAC name of  $[\text{Pt}(\text{NH}_3)_3\text{Br}(\text{NO}_2)\text{Cl}]\text{Cl}$  is  
 a) Triamminechlorobromonitro platinum (IV) chloride  
 b) Triamminebromonitrochloro platinum (IV) chloride  
 c) Triamminebromochloronitro platinum (IV) chloride  
 d) Triamminenitrochlorobromo platinum (IV) chloride
78. An aromatic ether is not cleaved by HI even at 525 K. The compound is:  
 a)  $\text{C}_6\text{H}_5\text{OCH}_3$                       b)  $\text{C}_6\text{H}_5\text{OC}_6\text{H}_5$                       c)  $\text{C}_6\text{H}_5\text{OC}_3\text{H}_7$                       d) 
79. Phenol does not react with:  
 a)  $\text{Na}_2\text{CO}_3$                       b) NaOH                      c)  $\text{NaHCO}_3$                       d) KOH
80.  $[\text{EDTA}]^{4-}$  is a  
 a) Monodentate ligand                      b) Bidentate ligand  
 c) Quadridentate ligand                      d) Hexadentate ligand
81.  $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]\text{Br}_2$  and  $[\text{Pt}(\text{NH}_3)_4\text{Br}_2]\text{Cl}_2$  are related to  
 a) Optical isomer                      b) Linkage isomers                      c) Coordinate isomers                      d) Ionization isomers
82. Ferrocene is an example of  
 a) Sand-wiched complex  
 b) Pi-bonded complex  
 c) A complex in which all the five carbon atoms of cyclopentadiene anion are bonded to the metal  
 d) All of the above
83. Which compound is zero valent metal complex?  
 a)  $[\text{Cu}(\text{NH}_3)_4]\text{SO}_4$                       b)  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$                       c)  $[\text{Ni}(\text{CO})_4]$                       d)  $\text{K}_3[\text{Fe}(\text{CN})_6]$
84. Which of the following compounds is 2, 2, 3-trimethyl hexane?  
 a)  $(\text{CH}_3)_3\text{CCH}(\text{CH}_3)\text{CH}_2\text{CH}_3$                       b)  $(\text{CH}_3)_3\text{CCH}_2(\text{CH}_3)_2$   
 c)  $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{C}(\text{CH}_3)_3$                       d)  $(\text{CH}_3)_3\text{CCH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3$
85. The formula of phenoxy benzene is:  
 a)  $\text{C}_6\text{H}_5\text{C}_6\text{H}_5$                       b)  $\text{C}_6\text{H}_5-\text{O}-\text{C}_6\text{H}_5$                       c)  $\text{C}_6\text{H}_5-\text{O}-\text{C}_6\text{H}_6$                       d) None of these

86. Ziegler-Natta catalyst is an organometallic compound containing  
 a) Iron                              b) Titanium                              c) Rhodium                              d) Zirconium
87. Ziegler-Natta catalyst is  
 a)  $(\text{Ph}_3\text{P})_3\text{RhCl}$                               b)  $\text{K}[\text{PtCl}_3(\text{C}_2\text{H}_4)]$                               c)  $[\text{Al}_2(\text{C}_2\text{H}_6)_6 + \text{TiCl}_4]$                               d)  $[\text{Fe}(\text{C}_2\text{H}_5)_2]$
88. The tendency to show complex formation is maximum in ....elements.  
 a) *s*-block                              b) *p*-block                              c) *d*-block                              d) *f*-block
89. EDTA has coordination number  
 a) 3                              b) 4                              c) 5                              d) 6
90. Which of the following is used in Friedel-Craft's acylation reaction?



- b)  $\text{CH}_3\text{CH}_2\text{Cl}$                               c)  $\text{CH}_3\text{COOCH}_3$                               d)  $\text{CH}_3\text{Cl}$


91. The correct IUPAC name of  $\text{Mn}_3(\text{CO})_{12}$  is

- a) Dodacacarbonyl manganate (0)                              b) Dodacacarbonyl manganate (II)  
 c) Didacacarbonyl trimanganese (0)                              d) Manganic dodecacarbonyl (0)

92. The  $\pi$  -bonded organometallic compound which has ethene as one of its component is





- a) Zeise's salt                              b) Ferrocene                              c) Dibenzene chromium                              d) Tetraethyl tin

93. IUPAC name of the compound



- a) Ethyl-2-methyl-2-(*m*-nitro) phenyl propanoate                              b) Ethyl-2-methyl-2-(*o*-nitro) phenyl propanoate  
 c) Ethyl-2-methyl-2-(3-nitro) phenyl propanoate                              d) Ethyl-2-methyl-2-(3-nitro) phenyl propanoic acid

94. What is the product obtained in the following reaction:



95.  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$  is called:

- a) Hexaammine cobalt (III) chloride  
 b) Amino cobalt chloride (III)  
 c) Cobalt chloride hexaammine  
 d) Hexaammine trisobalt chloride


96. The complexes  $[\text{PtCl}_2(\text{NH}_3)_4]\text{Br}_2$  and  $[\text{PtBr}_2(\text{NH}_3)_4]\text{Cl}_2$  are example for isomerism

- a) Geometrical                              b) Optical                              c) Ionization                              d) Linkage

97. Geometrical shapes of the complexes formed by the reaction of  $\text{Ni}^{2+}$  with  $\text{Cl}^-$ ,  $\text{CN}^-$  and  $\text{H}_2\text{O}$ , respectively, are

- a) Octahedral, tetrahedral and square planar                              b) Tetrahedral, square planar and octahedral  
 c) Square planar, tetrahedral and octahedral                              d) Octahedral, square planar and octahedral

98. Identify the correct order of reactivity in electrophilic substitution reactions of the following compounds:







- a)  $1 > 2 > 3 > 4$                               b)  $4 > 3 > 2 > 1$                               c)  $2 > 1 > 3 > 4$                               d)  $2 > 3 > 1 > 4$

99. The centric formula for benzene was proposed by:





- a) Dewar  
b) Armstrong and Baeyer  
c) Ladenberg  
d) Kekule
100. Which is the correct statement?  
a) Benzyl alcohol is more acidic than phenol  
b) Ethanol is a powerful oxidizing agent  
c) Phenol is more acidic than propanol  
d) Ethane has high boiling point than ethanol
101. Phenol on sulphonation gives:  
a) *o*-phenol sulphonic acid  
b) *p*-phenol sulphonic acid  
c) *m*-phenol sulphonic acid  
d) Mixture of *o*- and *p*-phenol sulphonic acids
102. Which of the following organometallic compound is  $\sigma$  and  $\pi$  bonded?  
a)  $\text{Fe}(\text{CH}_3)_3$                       b)  $[\text{Co}(\text{CO})_5\text{NH}_3]^{2+}$                       c)  $[\text{Fe}(\eta^5 - \text{C}_5\text{H}_5)_2]$                       d)  $\text{K}[\text{PtCl}_3(\eta^2 - \text{C}_2\text{H}_4)]$
103. The number of double bonds in BHC (gammexane) is:  
a) 1                                      b) 2                                      c) 3                                      d) Zero
104. Given the molecular formula of the hexa coordinated complexes (A)  $\text{CoCl}_3 \cdot 6\text{NH}_3$  (B)  $\text{CoCl}_3 \cdot 5\text{NH}_3$  (C)  $\text{CoCl}_3 \cdot 4\text{NH}_3$ . If the number of coordinated  $\text{NH}_3$  molecules in A, B and C respectively are 6, 5 and 4, primary valency in (A), (B) and (C) are  
a) 0, 1, 2                              b) 3, 2, 1                              c) 6, 5, 4                              d) 3, 3, 3
105. Type of isomerism shown by  $[\text{Cr}(\text{NH}_3)_5\text{NO}_2]\text{Cl}_2$  is  
a) Optical                              b) Ionisation                              c) Geometrical                              d) Linkage
106.  $[\text{Sc}(\text{H}_2\text{O})_6]^{3+}$  ion is  
a) Colourless and diamagnetic                              b) Coloured and octahedral  
c) Colourless and paramagnetic                              d) Coloured and paramagnetic
107. Which one of the following octahedral complexes will not show geometrical isomerism? (A and B are monodentate ligands)  
a)  $[\text{MA}_4\text{B}_2]$                               b)  $[\text{MA}_5\text{B}]$                               c)  $[\text{MA}_2\text{B}_4]$                               d)  $[\text{MA}_3\text{B}_3]$
108. The IUPAC name of the following compound is  

$$\begin{array}{c} \text{O}=\text{C}-\text{CH}-\text{CH}_2 \\ | \quad | \quad | \\ \text{OH} \quad \text{NH}_2 \quad \text{OH} \end{array}$$
  
a) 3-amino-2-hydroxy propanoic acid                              b) 2-aminopropan-3-ol-1-oic acid  
c) 2-amion-3-hydroxy propanoic acid                              d) Aminohydroxy propanoic acid
109. Which of the following complex ion is not expected to absorb visible light?  
a)  $[\text{Ni}(\text{CN})_4]^{2-}$   
b)  $[\text{Cr}(\text{NH}_3)_6]^{3+}$   
c)  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$   
d)  $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$
110. The correct sequence of activating power of a group in benzene is:  
a)  $-\text{NH}_2 > -\text{NHCOCH}_3 > -\text{CH}_3$   
b)  $-\text{NH}_2 < -\text{NHCOCH}_3 < -\text{CH}_3$   
c)  $-\text{NH}_2 > -\text{NHCOCH}_3 < -\text{CH}_3$   
d)  $-\text{NH}_2 < -\text{NHCOCH}_3 > -\text{CH}_3$
111. The pair of compounds having metals in their highest oxidation state is  
a)  $\text{MnO}_2, \text{FeCl}_3$                               b)  $[\text{MnO}_4]^- , \text{CrO}_2\text{Cl}_2$   
c)  $[\text{Fe}(\text{CN})_6]^{3-} , [\text{Co}(\text{CN})_3]$                               d)  $[\text{NiCl}_4]^{2-} , [\text{CoCl}_4]^-$
112. Total number of geometrical isomers for the complex  $[\text{RhCl}(\text{CO})(\text{PPh}_3)(\text{NH}_2)]$  is  
a) 1                                      b) 2                                      c) 3                                      d) 4

113. The reaction of chloroform with alc. KOH and *p*-toluidine forms:



114. Which order is correct in spectrochemical series of ligands?



115. The IUPAC name of compound  $\text{K}_3[\text{Fe}(\text{CN})_5\text{NO}]$  is

a) Pentacyano nitrosyl potassium ferrate(II)

b) Potassium cyano pentanitrosyl ferrate(II)

c) Potassium pentacyanonitrosyl ferrate (III)

d) Potassium pentacyanonitrosyl ferrate (II)

116. The colour of  $[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$  is due to:

a) Transfer of an electron from one Ti to another

b) Presence of water molecule

c) Excitation of electrons from *d* – *d*

d) Intramolecular vibration

117. The oxidation number of Fe in  $\text{K}_4[\text{Fe}(\text{CN})_6]$  is





a) +3

b) +4



c) +2

d) –2

118. Correct structures of [E][S]-5-bromo-2,7-dimethyl, non-4-ene is



119. Name the metal *M* which is extracted on the basis of following reactions,



a) Nickel

b) Silver

c) Copper

d) Mercury

120. EAN of Cr in  $[\text{Cr}(\text{NH}_3)_6]\text{Cl}_3$  is:

a) 32

b) 33

c) 34

d) 35

121. The complex  $[\text{Pt}(\text{NH}_3)_6]\text{Cl}_4$  furnishes:

a) 5 ions

b) 6 ions

c) 4 ions

d) 2 ions

122. Ammoniacal solution of  $\text{Ni}(\text{CN})_2$  reacts with  $\text{C}_6\text{H}_6$  to produce a light violet coloured crystalline compound of the formula:

a)  $\text{Ni}(\text{CN})_2 \cdot \text{C}_6\text{H}_5$

b)  $\text{C}_6\text{H}_5\text{CH}_3$

c)  $\text{Ni}(\text{CN})_2\text{C}_6\text{H}_6$

d)  $\text{Ni}(\text{CN})_2\text{NH}_3 \cdot \text{C}_6\text{H}_6$

123. Ammonia forms the complex ion  $[\text{Cu}(\text{NH}_3)_4]^{2+}$  with copper ions in alkaline solution but not in acidic solution. What is the reason for it?

a) In acidic solutions, hydration protects copper ions

b) In alkaline solution, insoluble  $\text{Cu}(\text{OH})_2$  is precipitated which in excess of any alkali

- c) Copper hydroxide is an amphoteric substance  
 d) In acidic solutions, protons coordinate with ammonia molecules forming  $\text{NH}_4^+$  ions and  $\text{NH}_3$  molecules are not available

124. Which of the following shows geometrical isomerism?

- a) 1, 2-dichloroethane  
 b) 1, 2-dimethylcyclopropane  
 c)  $\text{CH}_3\text{CH} \begin{array}{l} \diagup \text{CO-NH} \\ \diagdown \text{NH-CO} \end{array} \text{CHCH}_3$   
 d) All of the above






125. The shape of the complex  $[\text{Ag}(\text{NH}_3)_2]^+$  is:

- a) Octahedral  
 b) Square planar  
 c) Tetrahedral  
 d) Linear

126. The  $\pi$ -bonded organometallic compound which has ethane as one of its component is

- a) Dibenzene chromium  
 b) Zeise salt  
 c) Ferrocene  
 d) Tetraethyl tin

127. The major product of the following reaction is:



128. Which is true in the case of  $\text{Ni}(\text{CO})_4$  complex?


- a) Hybridization of Ni is  $sp^3$   
 b) Tetrahedral shape of the molecule  
 c) Diamagnetic  
 d) All are correct

129. The reaction,  $\text{C}_6\text{H}_5\text{N}_2\text{Cl} \xrightarrow{\text{Cu}_2\text{Cl}_2/\text{HCl}} \text{C}_6\text{H}_5\text{Cl} + \text{N}_2$  is called:

- a) Etard's reaction  
 b) Sandmeyer's reaction  
 c) Wurtz-Fittig reaction  
 d) Perkin's reaction

130. Which of the following does not show optical isomerism?

- a)  $[\text{Co}(\text{en})_3]^{3+}$       b)  $[\text{Co}(\text{en})_2\text{Cl}_2]^+$       c)  $[\text{Co}(\text{NH}_3)_3\text{Cl}_3]^0$       d)  $[\text{Co}(\text{en})\text{Cl}_2(\text{NH}_3)_2]^+$




Having the IUPAC name as

- a) 1, 2-dimethyl cyclobutane      b) 2, 3-dimethyl cyclobutene  
c) 2, 3-dimethyl butane      d) 1, 2-dimethyl cyclobut-1-ene

132. Which of the following ions is produced when we prepare nitrating mixture by mixing together concentrated  $\text{HNO}_3$  and concentrated  $\text{H}_2\text{SO}_4$ ?

- a)  $\text{NO}_2^-$       b)  $\text{NO}_2^+$       c)  $\text{NO}_3^-$       d)  $\text{SO}_3^+\text{H}$

133. The correct IUPAC name of



- a) 1-bromo-2-chloro-6-fluoro-4-iodobenzene      b) 1-bromo-6-chloro-2-fluoro-4-iodobenzene  
c) 2-bromo-1-chloro-3-fluoro-5-iodobenzene      d) 2-bromo-3-chloro-1-fluoro-5-iodobenzene

134.  $[\text{Co}(\text{NH}_3)_4(\text{NO}_2)_2]\text{Cl}$  exhibits:

- a) Ionization isomerism, geometrical isomerism and optical isomerism  
b) Linkage isomerism, geometrical isomerism and optical isomerism  
c) Linkage isomerism, ionization isomerism and optical isomerism  
d) Linkage isomerism, ionization isomerism and geometrical isomerism

135. Which of the following complexes are not correctly matched with hybridisation of their central metal ion?

1.  $[\text{Ni}(\text{CO})_4]$        $sp^3$   
2.  $[\text{Ni}(\text{CO})_4]^{2-}$        $sp^3$   
3.  $[\text{CoF}_6]^{3-}$        $d^2sp^3$   
4.  $[\text{Fe}(\text{CN})_6]^{3-}$        $sp^3d^2$

Select the correct answer using the codes given below

- a) 1 and 2      b) 1 and 3      c) 2 and 4      d) 2, 3 and 4


136. Which of the following is an explosive?

- a)  $\text{PCl}_5$       b)  $\text{HNO}_3$       c)  $\text{C}_6\text{H}_5\text{OH}$       d) 2,4,6-trinitrophenol

137. The coordination number of Cr in  $[\text{Cr}(\text{NH}_3)_3(\text{H}_2\text{O})_3]\text{Cl}_3$  is:

- a) 3      b) 4      c) 6      d) 2

138. The major product obtained when 3-phenyl-1, 2-propane-diol is heated with  $\text{H}_2\text{SO}_4$  is:

- a)  $\text{C}_6\text{H}_5-\text{CH}_2-\text{CO}-\text{CH}_3$   
b)  $\text{C}_6\text{H}_5-\text{CH}_2-\text{CH}_2-\text{CHO}$   
c)  $\text{C}_6\text{H}_5-\text{CH}_2-\text{CH}=\text{CH}_2$   
d) 

139. Rate of substitution in phenol is:





- a) Slower than as in benzene  
b) Faster than as in benzene  
c) Equal to that as in benzene  
d) None of the above

140. Magnetic moment of  $[\text{Ag}(\text{CN})_2]^-$  is zero. How many unpaired electrons are there?

- a) Zero      b) 4      c) 3      d) 1

141. Chlorophyll is a coordination compound having central atom of:

- a) Ca      b) Mg      c) Na      d) K

142. Which of the following statements is incorrect?
- In  $K_3[Fe(CN)_6]$ , the ligand has satisfied only the secondary valency of ferric ion.
  - In  $K_3[Fe(CN)_6]$ , the ligand has satisfied both primary and secondary valencies of ferric ion.
  - In  $K_4[Fe(CN)_6]$ , the ligand has satisfied both primary and secondary valencies of ferrous ion.
  - In  $[Cu(NH_3)_4]SO_4$ , the ligand has satisfied only the secondary valency of copper.
143. Maximum number of open chain isomers that an alkene can have with the molecular formula  $C_4H_8$  is
- 5
  - 4
  - 3
  - 2
144. Which one is the wrong statement?
- Open chain compounds are called aliphatic
  - Unsaturated compounds contain multiple bonds in them
  - Saturated hydrocarbons are called alkene
  - Aromatic compounds possess a characteristic aroma
145. According to postulates of Werner's theory for coordination compounds, which of the following is true?
- Primary valencies are ionizable
  - Secondary valencies are ionizable
  - Only primary valencies are non-ionizable
  - Primary and secondary valencies are non-ionizable
146. Atomic numbers of Cr and Fe are respectively 24 and 26. Which of the following is paramagnetic with the spin of the electron?
- $[Cr(CO)_6]$
  - $[Fe(CO)_5]$
  - $[Fe(CN)_6]^{4-}$
  - $[Cr(NH_3)_6]^{3+}$
147. Which of the following structures correspond to the product expected, when excess of  $C_6H_6$  reacts with  $CH_2Cl_2$  in presence of anhy.  $AlCl_3$  ?
- 
  - 
  - 
  - 
148. Which of the following will give a pair of enantiomorphs?
- $[Co(en)_2Cl_2]Cl$
  - $[Cr(NH_3)_6][Co(CN)_6]$
  - $[Pt(NH_3)_4][PtCl_6]$
  - $[Co(NH_3)_4Cl_2]NO_2$
149. The crystal field splitting energy for octahedral ( $\Delta_0$ ) and tetrahedral ( $\Delta_t$ ) complexes is related to
- $\Delta_t = \frac{4}{9}\Delta_0$
  - $\Delta_t = \frac{1}{2}\Delta_0$
  - $\Delta_0 = 2\Delta_t$
  - $\Delta_0 = \frac{4}{9}\Delta_t$
150. The correct name of the compound  $[Cu(NH_3)_4](NO_3)_2$ , according to IUPAC system is:
- Cuprammonium nitrate
  - Tetraamminecopper(II) dinitrate
  - Tetraamminecopper(II) nitrate
  - Tetraamminecopper(I) dinitrate
151. Which among the following will not show chain isomerism?
- $C_3H_8$
  - $C_4H_{10}$
  - $C_5H_{12}O$
  - $C_5H_{10}O$
152. Phenol (1 mole) reacts with bromine to give s-tribromophenol. How much bromine is needed?
- 1.5 mole
  - 3.0 mole
  - 4.5 mole
  - 6.0 mole
153. Dimethyl glyoxime forms a coloured complex with
- Ag
  - Ni
  - Cr
  - Zn
154. Which has regular tetrahedral geometry?
- $[Ni(CN)_4]^{2+}$
  - $SF_4$
  - $[BF_4]^-$
  - $XeF_4$
155. In haemoglobin the iron shows oxidation state:
- +2
  - +3
  - +1
  - +4
156. For the given complex  $[CoCl_2(en)(NH_3)_2]^+$ , the number of geometrical isomers, the number of optical isomers and total number of isomers of all type possible respectively are
- 2, 2 and 4
  - 2, 2 and 3
  - 2, 0 and 2
  - 0, 2 and 2
157. Which can show aromatic character?

- a) Furan                      b) Pyrrol                      c) Benzene                      d) All of these
158. Of the following complexes, the one with the largest value of the crystal field splitting is:  
 a)  $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$                       b)  $[\text{Ru}(\text{CN})_6]^{3-}$                       c)  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$                       d)  $[\text{Fe}(\text{NH}_3)_6]^{3+}$
159. The specific rotation of a pure enantiomer is  $+16^\circ$ . Its observed rotation if it is isolated from a reaction with 25% racemisation and 75% retention is  
 a)  $-12^\circ$                       b)  $+12^\circ$                       c)  $+16^\circ$                       d)  $-16^\circ$
160. Lithium tetrahydridoaluminate is correctly represented as:  
 a)  $\text{Al}[\text{LiH}_4]$                       b)  $\text{Al}_2[\text{LiH}_4]_3$                       c)  $\text{Li}[\text{AlH}_4]$                       d)  $\text{Li}[\text{AlH}_4]_2$
161. Which of the following compounds is generally used for hydrogenation of alkenes?  
 a)  $\text{Ni}(\text{CO})_4$                       b)  $[(\text{C}_6\text{H}_5)_3\text{P}]_3\text{RhCl}$                       c)  $(\text{CH}_3)_3\text{Al}$                       d)  $(\text{C}_5\text{H}_5)_2\text{Fe}$
162. The end product of the reaction,  
 $\text{C}_6\text{H}_6 + \text{Cl}_2 \xrightarrow{\text{Sunlight}}$  is:  
 a)  $\text{C}_6\text{H}_5\text{Cl}$                       b) *o*- $\text{C}_6\text{H}_4\text{Cl}_2$                       c)  $\text{C}_6\text{H}_6\text{Cl}_6$                       d) *p*- $\text{C}_6\text{H}_4\text{Cl}_2$
163.  $[\text{Pt}(\text{NH}_3)_6]\text{Cl}_4$  complex gives  
 a) 4 ions                      b) 3 ions                      c) 2 ions                      d) 5 ions
164. Which does not obey EAN rule?  
 a)  $[\text{Cu}(\text{NH}_3)_4]^{2+}$                       b)  $[\text{Zn}(\text{OH})_4]^{2-}$                       c)  $[\text{HgI}_4]^{2-}$                       d)  $\text{Fe}(\text{CO})_5$
165. Oxidation number of Fe in  $\text{K}_3[\text{Fe}(\text{CN})_6]$  is:  
 a) +3                      b) +2                      c) +10                      d) 1
166. Which of the following is not an organometallic compound?  
 a)  $\text{NaOC}_2\text{H}_5$                       b)  $(\text{CH}_3)_3\text{Al}$                       c)  $(\text{C}_2\text{H}_5)_4\text{Pb}$                       d)  $\text{RMgX}$
167. Considering  $\text{H}_2\text{O}$  as weak field ligand, the number of unpaired electrons in  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$  will be (Atomic no. of Mn=25)  
 a) Three                      b) Five                      c) Two                      d) Four
168. The value of 'spin only' magnetic moment for one of the following configuration is 2.84 BM the correct one  
 a)  $d^4$  (in weak ligand field)                      b)  $d^4$  (in strong ligand field)  
 c)  $d^3$  (in weak as well as in strong field)                      d)  $d^5$  (in weak ligand field)
169. Fluorobenzene ( $\text{C}_6\text{H}_5\text{F}$ ) can be synthesized in the laboratory:  
 a) By heating phenol with HF and KF  
 b) From aniline by diazotisation followed by heating the diazonium salt with  $\text{HBF}_4$   
 c) By direct fluorination of benzene with  $\text{F}_2$  gas  
 d) By reacting bromobenzene with NaF solution
170. Which compound burns with a sooty flame?  
 a)  $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$   
 b)  $\text{C}_6\text{H}_5\text{COOH}$   
 c)  $\text{CH}_3\text{OH}$   
 d)  $\text{CH}_3\text{COC}_2\text{H}_5$
171. How many EDTA (ethylenediaminetetraacetic acid) molecules are required to make an octahedral complex with a  $\text{Ca}^{2+}$  ion?  
 a) Six                      b) Three                      c) One                      d) Two
172. Intramolecular rearrangement of phenyl esters to give *o*- and *p*-derivatives in presence of  $\text{AlCl}_3$  is known as:  
 a) Friedel-Craft's reaction  
 b) Fries rearrangement  
 c) Esterification  
 d) Coupling
173. Which reaction can produce  $\text{R}-\text{CO}-\text{Ar}$  species?  
 a)  $\text{ArCOCl} + \text{H}-\text{Ar} \xrightarrow{\text{AlCl}_3}$                       b)  $\text{COCl} + \text{RMgX} \rightarrow$                       c)  $\text{RCOCl} + \text{H}-\text{Ar} \xrightarrow{\text{AlCl}_3}$                       d)  $\text{R} + \text{CrO}_3 \rightarrow$
174. Acidic character of phenol is due to:  
 a) Resonance of phenoxide ion




188. When phenol is treated with excess bromine water, it gives:

- a) *m*-bromophenol  
 b) *o*- and *p*-bromophenol  
 c) 2,4-dibromophenol  
 d) 2,4,6-tribromophenol

189. Which have octahedral shape ( $d^2sp^3$ ) hybridization of central atom?


- a)  $[\text{Cr}(\text{NH}_3)_6]^{2+}$       b)  $[\text{Fe}(\text{CN})_6]^{3-}$       c)  $[\text{Cu}(\text{NH}_3)_6]^+$       d) All are correct

190. Which of the following molecules/species are aromatic in character?



- a) 2      b) 3      c) 4      d) 1

191. Among the following compounds ;



the order of basicity is :

- a) IV > III > II > I      b) II > I > III > IV      c) III > IV > II > I      d) I > III > IV > II

192.

The correct name of  $\text{CO}_3\text{Fe} \begin{array}{c} \diagup \text{CO} \diagdown \\ \text{--- CO ---} \\ \diagdown \text{CO} \diagup \end{array} \text{Fe CO}_3$  is :

- a) Tri- $\mu$ -carbonyl *bis*-(tricarbonyl)iron (0)  
 b) Hexacarbonyl iron (III)  $\mu$ -tricarbonyl ferrate(0)  
 c) Tricarbonyl iron(0)  $\mu$ -tricarbonyl iron(0) tricarbonyl  
 d) Nonacarbonyl iron

193. Which is high spin complex?

- a)  $[\text{CoCl}_6]^{3-}$       b)  $[\text{FeF}_6]^{3-}$       c)  $[\text{Co}(\text{NH}_3)_6]^{2+}$       d) All are correct

194. The correct IUPAC name of tartaric acid is

- a) 1, 4-dicarboxy-2, 3-dihydroxy ethane      b)  $\alpha, \alpha'$ -dihydroxy butane-1,4-dioic acid  
 c) 1, 4-dihydroxybutane-2, 3-dioic acid      d) 2, 3-dihydroxybutane-1, 4-dioic acid


195. What is the overall formation equilibrium constant for the ion  $[\text{ML}_4]^{2-}$  ion, given that  $\beta_4$  for this complex is  $2.5 \times 10^{13}$ ?

- a)  $2.5 \times 10^{13}$       b)  $5 \times 10^{-13}$       c)  $2.5 \times 10^{-14}$       d)  $4.0 \times 10^{-13}$

196. The oxidation state of Cr in  $[\text{Cr}(\text{NH}_3)_4 \text{Cl}_2]^+$  is

- a) 0      b) +1      c) +2      d) +3

197. Which of the following compounds has the most acidic nature?



198. The oxidation state of Mo in its oxo-complex species  $[\text{Mo}_2\text{O}_4(\text{C}_2\text{H}_4)_2(\text{H}_2\text{O})_2]^{2-}$  is:

- a) +2      b) +3      c) +4      d) +5

199.  $\text{CH}_3\text{MgI}$  is an organometallic compound due to

- a) Mg—I bond      b) C—I bond      c) C—Mg bond      d) C—H bond

200. The effective atomic number of Cr (At. No.=24) in  $[\text{Cr}(\text{NH}_3)_6]\text{Cl}_3$  is

- a) 35      b) 27      c) 33      d) 36

201. When aniline is heated with benzaldehyde, the product is:

- a) Benzoin      b) Schiff's base      c) Unsaturated acid      d) Azoxy benzene

202. Slow heating of salicylic acid gives:  
a) Benzoic acid                      b) Phenol                                      c) Benzaldehyde                              d) None of these


203. According to Hückel, monocyclic compounds will show aromaticity when:

- a) It has  $4\pi$ -electrons
- b) It has no  $\pi$ -electron
- c) It has  $4\pi+2$  electrons
- d) It has  $(4n + 2)\pi$ -electrons

204. When phenol is distilled with zinc dust, it gives:





- a) Benzene                                      b) Toluene                                      c)  $C_6H_5CHO$                                       d) None of these

205. The IUPAC name of the given structure



- a) N-chloro-N-bromoethanamide                                      b) N-bromo-N-chloroethanamide
- c) N-bromo-N-chloroacetamide                                      d) N-chloro-N-bromoacetamide

206. Acetophenone when reacted with a base  $C_2H_5ONa$ , yields a stable compound which has the structure:




207. Which of the following has maximum resonance energy?

- a) Diphenyl                                      b) Benzene                                      c) Naphthalene                                      d) Phenanthrene

208. Benzene sulphonic acid on treating with  $P_2O_5$  gives:

- a) Salicylic acid                                      b) Benzoic acid                                      c) Acid anhydride                                      d) Sodium benzoate

209. Compounds with following formula will show



- a) Position and functional isomerism
- b) Chain and positional isomerism
- c) Chain and functional isomerism
- d) None of the above combinations





210. Which of the following statements is correct?


- a) In  $K_3[Fe(CN)_6]$ , the ligand has satisfied both primary and secondary valencies of ferric ion
- b) In  $(Cu(NH_3)_4)SO_4$ , the ligand has satisfied only the secondary valency of copper
- c) In  $K_3[Fe(CN)_6]$ , the ligand has satisfied only the secondary valency of ferric ion
- d) Both (b) and (c)

211. Which statement is not correct?

- a)  $Fe(CO)_5$  reacts with  $Br_2Cl_4$

- b) Carbonyl complexes are usually formed with transition metals  
 c) All transition metals form monometallic carbonyls  
 d) The decomposition of  $\text{Ni}(\text{CO})_4$  to give Ni is used in the extraction of Ni by Mond's process
212. The complex showing a spin-only magnetic moment of 2.82 BM is  
 a)  $\text{Ni}(\text{CO})_4$                       b)  $[\text{NiCl}_4]^{2-}$                       c)  $\text{Ni}(\text{PPh}_3)_4$                       d)  $[\text{Ni}(\text{CN})_4]^{2-}$
213. The IUPAC name of  $[\text{CoCl}(\text{NO}_2)(\text{en})_2]\text{Cl}$  is:  
 a) Chloronitro-*bis*(ethylenediamine) cobaltic(III) chloride  
 b) Chloronitro-*bis*(ethylenediamine)cobalt(II) chloride  
 c) Chloro-*bis*(ethylenediamine)nitrocobalt(III) chloride  
 d) *Bis*-(ethylenediamine)chloronitrocobalt(III) chloride
214. The product of acid catalysed hydration of 2-phenyl propene is:  
 a) 3-phenyl-2-propanol  
 b) 1-phenyl-2-propanol  
 c) 2-phenyl-2-propanol  
 d) 2-phenyl-1-propanol
215. Carboic acid is the name used for:  
 a) Opium                      b) Phenol                      c) Chloroform                      d)  $\text{H}_2\text{CO}_3$
216. The major product of the following reaction





217. The oxidation number of cobalt in  $K[Co(CO)_4]$  is

- a) -1                                      b) +3                                      c) +1                                      d) -3

218. Formaldehyde-phenol resin is:

- a) Orlon                                      b) Nylon                                      c) Teflon                                      d) Bakelite

219. Among the ligands  $NH_3$ , en,  $CN^-$  and CO, the correct order of their increasing field strength, is

- a)  $CO < NH_3 < en < CN^-$                                       b)  $NH_3 < en < CN^- < CO$   
 c)  $CN^- < NH_3 < CO < en$                                       d)  $en < CN^- < NH_3 < CO$

220. Cyclopentadienyl anion is aromatic due to the presence of:

- a) 6 $\pi$ -electrons                                      b) 10  $\pi$ -electrons                                      c) 4  $\pi$ -electrons                                      d) 12  $\pi$ -electrons


221. The IUPAC name of  $K_4[Fe(CN)_6]$  is

- a) Potassium ferrocyanide                                      b) Potassium hexa cyanoferrate (I)  
 c) Tetra potassium hexa cyanoferrate (II)                                      d) Potassium hexa cyanoferrate (II)

222. Which xylene is most easily sulphonated?

- a) *Ortho*                                      b) *Para*                                      c) *Meta*                                      d) All at the same rate

223. The IUPAC name of following polyfunctional compound is



- a) 2,4-dioxo cyclohexanoic acid                                      b) 2,4-dioxo cycloheptanoic acid  
 c) 4-formyl-2-oxo cyclohexane-1-carboxylic acid                                      d) 2,4-dioxo cyclohexane-1-carboxylic acid

224. Alkyl groups are *o*- and *p*-directing because of:


- a) Resonance effect  
 b) Inductive effect  
 c) Resonance effect through hyperconjugation  
 d) All of the above


225. Racemic modification can be resolved by

- a) The use of enzymes                                      b) Fractional crystallisation  
 c) Fractional distillation                                      d) None of the above

226. Which of the following structure contain 1 primary and 7 secondary hydrogen atoms?

- a) 
$$\begin{array}{c} CH_3 - CH - CH_2 - CH_2 \\ | \qquad \qquad | \\ CH_3 \qquad \qquad CH - CH_3 \\ | \\ CH_3 \\ | \\ CH_3 \end{array}$$
- b) 
$$\begin{array}{c} CH_3 - CH_2 - CH_2 - CH \\ | \\ CH_2 - CH_3 \end{array}$$
- c) 
$$\begin{array}{c} CH_3 \\ | \\ CH_3 - CH_2 - CH - CH_2 - CH_3 \end{array}$$
- d)  $CH_3 - CH_2 - CH_2 - CH_2$



227. Which of the following compounds does not dissolve in conc.  $\text{H}_2\text{SO}_4$  even on warming?  
 a) Ethylene                      b) Benzene                      c) Hexane                      d) Aniline
228. In the complex  $\text{Fe}(\text{CO})_x$ , the value of  $x$  is and it is:  
 a) 3, octahedral                      b) 4, tetrahedral                      c) 5, trigonal pyramidal                      d) 6, square pyramidal
229. The empirical formula of naphthalene is:  
 a)  $\text{CH}_2$                       b)  $\text{C}_5\text{H}_4$                       c)  $\text{C}_2\text{H}$                       d)  $\text{C}_n\text{H}_{2n}$
230. The chemical formula of diammine silver (I) chloride is  
 a)  $[\text{Ag}(\text{NH}_3)\text{Cl}]$                       b)  $[\text{Ag}(\text{NH}_3)_3]\text{Cl}$                       c)  $[\text{Ag}(\text{NH}_4)_2]\text{Cl}$                       d)  $[\text{Ag}(\text{NH}_3)_2]\text{Cl}$
231. For the square planar complex  $[\text{M}(a)(b)(c)(d)]$  (where,  $\text{M}$ =central metal and  $a, b, c,$  and  $d$  are monodentate ligands), the number of possible geometrical isomers are  
 a) 1                      b) 2                      c) 3                      d) 4
232. Which group is *meta* directing?  
 a)  $-\text{CCl}_3$                       b)  $-\text{OH}$                       c)  $-\text{NH}_2$                       d)  $-\text{CH}_3$
233. The IUPAC name of the compound  $[\text{Cu}(\text{NH}_3)_4(\text{NO}_3)_2]$  is:  
 a) Cuprammonium nitrate  
 b) Dinitratotetraamminecopper(II)  
 c) Tetraamminecopper(II) dinitrite  
 d) Tetraamminecopper(III) dinitrite
234. Coordination number of Fe in the complexes  $[\text{Fe}(\text{CN})_6]^{4-}$ ,  $[\text{Fe}(\text{CN})_6]^{3-}$  and  $[\text{FeCl}_4]^-$  would be respectively  
 a) 6, 4, 6                      b) 6, 6, 4                      c) 6, 3, 3                      d) 2, 3, 3
235. Which statement is true for cyclohexane?  
 a) It has two possible isomers  
 b) It has three conformations  
 c) Boat conformation is most stable  
 d) Chair and boat conformations differ in energy by 44 kJ/mol
236. Ligands in a complex salt are:  
 a) Anions linked by coordinate bonds to a central metal atom or ion  
 b) Cations linked by coordinate bonds to a central metal atom or ion  
 c) Molecules linked by coordinate bonds to a central metal atom or ion  
 d) Ions or molecules linked by coordinate bonds to a central metal atom or ion
237. The IUPAC name of  $\text{CH}_3\text{COCH}_2\text{COOC}_2\text{H}_5$  is  
 a) Ethyl butanoate                      b) Ethyl-(3-oxo)butanoate  
 c) Ethyl butan-1-oate-2-one                      d) Ethyl butan-4-oate-2-one
238. When benzene is treated with CO and HCl in presence of anhydrous aluminium chloride, benzaldehyde is formed. This reaction is known as:  
 a) Friedel-Craft's reaction  
 b) Rosenmund's reaction  
 c) Stephen's reaction  
 d) Gattermann-Koch's reaction
239.  $[\text{Cr}(\text{NH}_3)_6]^{3+}$  ion is:  
 a) Paramagnetic                      b) Diamagnetic                      c) Square planar                      d) None of these
240. The following compound can exhibit  
  
 a) Tautomerism                      b) Optical isomerism  
 c) Geometrical isomerism                      d) Geometrical and optical isomerism
241. Which complex is diamagnetic?

- a)  $[\text{Fe}(\text{CN})_6]^{4-}$                       b)  $[\text{Cu}(\text{NH}_3)_4]^{3+}$                       c)  $[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$                       d) None of these
242. *Meso*-tartaric acid is optically inactive due to the presence of  
 a) Molecular symmetry                      b) Molecular asymmetry  
 c) External compensation                      d) Two asymmetric C-atoms

243. Complex forming tendency increases with:  
 a) Increase in size of cation  
 b) Decrease in size of cation  
 c) Increase in size of anion  
 d) None of the above

244. Ziegler-Natta catalyst is  
 a)  $(\text{Ph}_3\text{P})_3\text{RhCl}$                       b)  $\text{Al}_2(\text{C}_2\text{H}_6)_6 + \text{TiCl}_4$   
 c)  $\text{Fe}(\text{C}_2\text{H}_5)_2$                       d)  $\text{K}[\text{PtCl}_3(\text{C}_2\text{H}_4)]$

245. Among the following compounds the one that is most reactive towards electrophilic nitration is:  
 a) Toluene                      b) Benzene                      c) Benzoic acid                      d) Nitrobenzene

246. Phenol on oxidation gives chloranil. The oxidant used is:  
 a)  $\text{K}_2\text{S}_2\text{O}_8$                       b)  $\text{KMnO}_4$                       c)  $\text{KClO}_3 + \text{HCl}$                       d) None of these


247. The IUPAC name of the compound
- $$\begin{array}{ccccccc} & & \text{CH}_3 & - & \text{CH} & - & \text{CH}_2 & - & \text{CH}_3 \\ & & | & & | & & & & \\ \text{CH}_3 & - & \text{CH}_2 & - & \text{CH} & - & \text{CH} & - & \text{CH} & - & \text{CH}_2 & - & \text{CH}_2 & - & \text{CH}_3 \\ & & | & & | & & & & \\ & & \text{CH}_3 & & \text{CH}_2 & - & \text{CH}_3 & & \end{array}$$
- is

- a) 3-sec-butyl-5-ethyl-3-methyloctane                      b) 4-sec-butyl-5-ethyl-3-methyloctane  
 c) 5-sec-butyl-4-ethyl-3-methyloctane                      d) 4-sec-butyl-3-ethyl-5-methyloctane
248. All the common *m*-directing groups.....the benzene ring towards electrophilic substitution reactions.  
 a) Deactivate                      b) Activate                      c) Both (a) and (b)                      d) None of these
249. Among the following, the coloured compound is :  
 a)  $\text{CuCl}$                       b)  $\text{K}_3\text{C}_4(\text{CN})_4$                       c)  $\text{CuF}_2$                       d)  $[\text{Cu}(\text{CH}_3\text{CN})_4]\text{BF}_3$
250. The existence of two different coloured complexes with the composition of  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+$  is due to:  
 a) Linkage isomerism  
 b) Geometrical isomerism  
 c) Coordination isomerism  
 d) Ionisation isomersim

251.  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]$  possesses:  
 a) Square planar geometry  
 b) Tetrahedral geometry  
 c) Tetrahedral nature  
 d) Octahedral geometry


252. Which one does not belong to ligand?  
 a)  $\text{PH}_3$                       b)  $\text{NO}^+$                       c)  $\text{BF}_3$                       d)  $\text{Cl}^-$

253. Product formed in the reaction;  
 Phenol  $\xrightarrow[\text{Pyridine} - \text{SO}_3/(\text{CH}_3\text{CO})_2\text{O}]{(\text{CH}_3)\text{SO}}$  Product; is:



254. Which one of the following has square planar structure?  
 a)  $[\text{Ni}(\text{CN})_4]^{2-}$                       b)  $[\text{Ni}(\text{CO})_4]$                       c)  $[\text{NiCl}_4]^{2-}$                       d) All of these

255. 4-methyl benzene sulphonic acid reacts with sodium acetate to give:




256. Phthalein test is characteristics of ...and is given by it.

- a) Alcohols                                      b) Phenols                                      c) Aldehydes                                      d) Ketones


257. Which of the following compounds would exhibit coordination isomerism?

- a) [Cr(H<sub>2</sub>O)<sub>6</sub>]Cl<sub>2</sub>                                      b) [Cr(NH<sub>3</sub>)<sub>6</sub>][Co(CN)<sub>6</sub>]                                      c) [Cr(en)<sub>2</sub>]NO<sub>2</sub>                                      d) [Ni(NH<sub>3</sub>)<sub>6</sub>][BF<sub>4</sub>]<sub>2</sub>

258. In a reaction of aniline a coloured product C was obtained.



The structure of C would be:



259. The carboxyl functional group (-COOH) is present in:

- a) Picric acid  
b) Barbituric acid  
c) Ascorbic acid  
d) Aspirin

260. Which of the following is an example of electrophilic substitution reaction?

- a) Acylation                                      b) Alkylation                                      c) Benzoylation                                      d) All of these

261. The number of ions given by [Co(NH<sub>3</sub>)<sub>4</sub>]Cl<sub>3</sub> in aqueous solution is:

- a) 2                                      b) 3                                      c) 1                                      d) 4

262. Which of the following is an organometallic compound?

- a) Ti(OC<sub>6</sub>H<sub>5</sub>)<sub>4</sub>                                      b) Ti(OCOCH<sub>3</sub>)<sub>4</sub>                                      c) Ti(OC<sub>2</sub>H<sub>5</sub>)<sub>4</sub>                                      d) Ti(C<sub>2</sub>H<sub>5</sub>)<sub>4</sub>

263. A solution of CuCl in NH<sub>4</sub>OH is used to measure the amount of which gas in a sample by simply measuring change in volume?

- a) CO<sub>2</sub>                                      b) H<sub>2</sub>                                      c) CO                                      d) All of these

264. On passing benzene vapour through a tube at 700-800°C or through molten lead we get:

- a) Diphenyl                                      b) Phenol                                      c) Toluene                                      d) Benzaldehyde

265. Picric acid is a yellow coloured compound. Its chemical name is:


- a) *m*-nitrobenzoic acid                                      b) 2,4,6-trinitrophenol                                      c) Trinitrotoluene                                      d) Trinitroaniline

266. The ideal starting material for the synthesis of *m*-chloronitro benzene is:

- a) Benzene                      b) Chlorobenzene                      c) Toluene                      d) Nitrobenzene
267. In a reaction involving ring substitution of  $C_6H_5Y$ , the major product is *meta*-isomer. The group  $Y$  can be:
- a)  $-NH_2$                       b)  $-COOH$                       c)  $-CH_3$                       d)  $-Cl$

268. When ammonia is added to green aqueous solution of nickel(II) sulphate, the colour of the solution changes to blue violet. This is caused by:
- a) Nickel undergoing a change in oxidation state  
 b) Ammonia molecules replacing water molecules surrounding nickel  
 c) Change in coordination number of nickel  
 d) Change in pH value of the solution

269. The compound, whose stereo chemical formula is written below, exhibits  $x$ -geometrical isomers and  $y$ -optical isomers. The value of  $x$  and  $y$  are



- a) 4 and 4                      b) 2 and 2                      c) 2 and 4                      d) 4 and 2
270. Among the following-phenol, benzoic acid, nitrobenzene and toluene, the compound that undergoes nitration readily is:
- a) Benzoic acid                      b) Toluene                      c) Phenol                      d) Nitrobenzene

271. Which one is organometallic compound?

- a) Lithium acetate                      b) Lithium methoxide  
 c) Lithium dimethyl amide                      d) Methyl lithium

272. What are the products formed when an equimolar mixture of benzaldehyde and formaldehyde is heated with concentrated NaOH?

- a)  $C_6H_5-CH_2-OH$  and  $H-COONa$   
 b)  $C_6H_5-COONa$  and  $CH_3-OH$   
 c)  $C_6H_5-CH_2-COONa$   
 d)  $C_6H_5-COOH$  and  $CH_3-ONa$





273. Gammexane (a  $\gamma$ -isomer of) is:

- a) BHC  
 b) Benzene hexachloride  
 c) Lindane  
 d) All of these

274. Number of electrons gained by Pd in  $[PdCl_4]^{2-}$ :


- a) 4                      b) 8                      c) 10                      d) 0





275. Which of the following is considered to be an anticancer species?



276. For benzaldehyde which of the following is incorrect?

- a) It is an aromatic aldehyde

- b) It is used in perfumery  
 c) On oxidation it yields benzoic acid  
 d) On reduction it yields phenol
277. The main source of aromatic compounds is:  
 a) Wood                                      b) Petroleum                                      c) Coal                                      d) Both (b) and (c)
278. Phenol on hydrogenation in presence of a nickel catalyst at 160°C gives:  
 a) Benzene                                      b) Cyclohexane                                      c) Cyclohexanol                                      d) *n*-hexanol
279. The IUPAC name of compound
- 
- is
- a) Hexane-1, 2, 5-tricarbonitrile                                      b) Hexane-1, 3, 6-tricarbonitrile  
 c) Butane-1, 2, 4-tricarbonitrile                                      d) Butane-1, 3, 4-tircarbonitrile
280. *Fac-mer* isomerism is associated with which one of the following complexes? (*M*=central metal)  
 a)  $[M(AA)_2]$                                       b)  $[MA_3B_3]$                                       c)  $[M(AA)_3]$                                       d)  $[MABCD]$
281. Which of the following is the correct order of stability of the following four distinct conformation of *n* butane?  
 a) Staggered > Gauche > Partially eclipsed > Fully eclipsed  
 b) Gauche > Staggered > partially eclipsed > Fully eclipsed  
 c) Staggered > Partially eclipsed > Gauche > Fully eclipsed  
 d) Fully eclipsed > Staggered > Partially eclipsed > Gauche
282. *o*-nitrophenol can form hydrogen bonds within the molecule. It thus, has:  
 a) Very high m.p.                                      b) Very high viscosity                                      c) Low m.p.                                      d) none of these
283. The element which does not form mononuclear carbonyl is:  
 a) Fe                                      b) Mn                                      c) Ni                                      d) W
284. Which of the following is hexadentate ligand?  
 a) Ethylene diamine                                      b) Ethylene diamine tetra acetic acid  
 c) 1,10-phenanthroline                                      d) Acetyl acetato
285. The molecular formula of a saturated compound is  $C_2H_4Cl_2$ . The formula permits the existence of two  
 a) Functional isomers                                      b) Position isomers                                      c) Optical isomers                                      d) *cis* – *trans* isomers
286. An octahedral complex is formed when hybrid orbitals of the following type are involved  
 a)  $sp^3$                                       b)  $dsp^2$                                       c)  $d^2sp^3$                                       d)  $sp^2d^2$
287. The IUPAC name of the given compound  $CH_3 - CH = CH - COOC_2H_5$  is  
 a) Ethyl propenoate                                      b) Ethyl-2-butenolate  
 c) Ethyl-1-butenolate                                      d) Propene ethyl methanoate
288. Which product is not obtained by heating wood or coal in the absence of air?  
 a) Coal-tar                                      b) Naphthalene                                      c) Benzene                                      d) Wax
289. Dry distillation of calcium benzoate with calcium formate gives:  
 a) Acetaldehyde                                      b) Benzoic acid                                      c) Benzaldehyde                                      d) Benzoic anhydride
290. Which will give  $Fe^{3+}$  ions in solution?  
 a)  $[Fe(CN)_6]^{3-}$   
 b)  $Fe_2(SO_4)_3$   
 c)  $[Fe(CN)_6]^{4-}$   
 d)  $NH_4(SO_4)_2 \cdot FeSO_4 \cdot 6H_2O$
291. Each metal possesses:  
 a) Primary valencies satisfied by anions only  
 b) Secondary valencies satisfied by donor molecules  
 c) Coordination number  
 d) All of the above
292. Aspirin is:  
 a) Antibiotic                                      b) Antipyretic                                      c) Sedative                                      d) Psychedelic


293. Hybridisation, shape and magnetic moment of  $K_3[Co(CO_3)_3]$  is
- a)  $d^2 sp^3$ , octahedral, 4.9 BM                      b)  $sp^3 d^2$ , octahedral, 4.9 BM  
c)  $dsp^2$ , square planer, 4.9 BM                      d)  $sp^3$ , tetrahedral, 4.9 BM
294. Among the following complexes ( $K-P$ ),  
 $K_3[Fe(CN)_6]$  ( $K$ ),  $[Co(NH_3)_6]Cl_3$  ( $L$ ),  
 $Na_3[Co(ox)_3]$  ( $M$ )  
 $[Ni(H_2O)_6]Cl_2$  ( $N$ ), and  
 $[Zn(H_2O)_6](NO_3)_2$  ( $P$ ) the diamagnetic complexes
- a)  $K, L, M, N$                       b)  $K, M, O, P$                       c)  $L, M, O, P$                       d)  $L, M, N, O$
295. Aniline when diazotised in cold and then treated with dimethyl aniline gives a coloured product. Its structure would be:
- a) 
- b) 
- c) 
- d) 
296. Pyridine possesses:
- a) Aromatic nature  
b) Unsaturated aliphatic nature  
c) Alicyclic nature  
d) Aliphatic nature
297. A reagent used for identifying nickel ion is:
- a) Potassium ferrocyanide  
b) Phenolphthalein  
c) Dimethyl glyoxime  
d) EDTA
298. Aniline was diazotised and subsequently reduced with stannous chloride and hydrochloric acid to yield:
- a) Phenyl aniline                      b) Phenyl hydrazine                      c) *p*-amino azobenzene                      d) Diazoamino benzene
299. The reaction of toluene with  $Cl_2$  in presence of  $FeCl_3$  gives predominantly:
- a) *m*-chlorobenzene  
b) Benzoylchloride  
c) Benzyl chloride  
d) *o*- and *p*-chlorobenzene
300. Which statement is not correct in the case of  $[Co(NH_3)_6]^{3+}$  complex?
- a) It is octahedral in shape  
b) It involves  $d^2 sp^2$ -hybridization  
c) It has diamagnetic nature  
d) None of the above
301. Pick out the complex compound in which the central metal atom obeys EAN rule strictly
- a)  $K_4[Fe(CN)_6]$                       b)  $K_3[Fe(CN)_6]$                       c)  $[Cr(H_2O)_6]Cl_3$                       d)  $[Cu(NH_3)_4]SO_4$
302. Amongst the following, the compound that can be most readily sulphonated is:
- a) Benzene                      b) Methoxy benzene                      c) Toluene                      d) Chlorobenzene
303. *p*-chloroaniline and anilium hydrochloride can be distinguished by:
- a)  $P_2O_5$                       b)  $AgNO_3$                       c) Carbylamine test                      d) Sandmeyer's reaction
304. Pyrogallol is..... trihydroxy benzene.

- a) 1, 2, 4                      b) 1, 2, 3                      c) 1, 3, 5                      d) None of these

305. Phenol is weakly acidic but does not react with  $\text{NaHCO}_3$  like carboxylic acids hence:

- a) Phenol is weaker than carbonic acid  
b) Phenol is stronger than acid  
c) Phenol is stronger than carboxylic acid  
d) None of the above

306. *p*-cresol reacts with chloroform in alkaline medium to give compound (A) which adds hydrogen cyanide to form compound (B). The latter on acidic hydrolysis gives chiral carboxylic acid. The acid is:



307. The number of isomeric xylenes is:


- a) 2                      b) 3                      c) 4                      d) 1

308. The IUPAC name of  $[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}_2]\text{Cl}$  is:

- a) Tetrahydrodichlorochromium(III) chloride  
b) Tetraaquodichlorochromium(III) chloride  
c) Tetraaquodichlorochromium(I) chloride  
d) None of the above

309. Among the following metal carbonyls, C—O bond order is lowest in

- a)  $[\text{Mn}(\text{CO})_6]^+$                       b)  $[\text{Fe}(\text{CO})_5]$                       c)  $[\text{Cr}(\text{CO})_6]$                       d)  $[\text{V}(\text{CO})_6]^-$



- a) 3-propyl-1,3-pentadiene                      b) 3,3-dipropyl-1,3-pentadiene  
c) 3,3-diethenyl penta-1,4-diene                      d) 4,4-diethenyl penta,1,2-diene


311. Which of the following shell, form an outer octahedral complex?

- a)  $d^4$                       b)  $d^8$                       c)  $d^6$                       d) None of these


312. Friedel-Craft's reaction of bromobenzene with methyl iodide gives:

- a) *o*-bromotoluene  
b) *p*-bromotoluene  
c) *o*- and *p*-bromotoluene  
d) *m*-bromotoluene

313. An organic compound  $\text{C}_7\text{H}_8\text{O}$  is neither soluble in  $\text{NaOH}$  nor gives blue colour with  $\text{FeCl}_3$ , is:

- a)  $\text{C}_6\text{H}_5 \cdot \text{CH}_2\text{OH}$                       b)                       c)  $\text{C}_6\text{H}_5 \cdot \text{OCH}_3$                       d) None of these

314. Which exist as a pair of mirror image isomers?




315. Benzene double bonds are not so reactive as those of hexatriene because:


- a) The three double bonds are caged in a ring  
b) Benzene is aromatic and has six  $\pi$ -resonating electrons  
c) Benzene has no double bond

d) Benzene is non-polar

316. The most stable ion is



317. Which of the following is/are three isomers?



a) Only (i)

b) Only (ii)

c) Only (iii)

d) All (i), (ii) and (iii)

318. In the coal-tar distillation of middle oil, the aromatic compounds present are:


a) Benzene, naphthalene, anthracene

b) Naphthalene, pyridine, phenol

c) Naphthalene, pyridine

d) None of the above

319. The correct order of increasing reactivity of C—X bond towards nucleophilic in the following compound is:



a)  $\text{I} < \text{II} < \text{IV} < \text{III}$

b)  $\text{II} < \text{III} < \text{I} < \text{IV}$

c)  $\text{IV} < \text{III} < \text{I} < \text{II}$

d)  $\text{III} < \text{II} < \text{I} < \text{IV}$

320. Which of the following system is most stable for a chelate?

a) Two fused cyclic system

b) Three fused cyclic system

c) Four fused cyclic system

d) Five fused cyclic system

321. Which of the following reaction take place when a mixture of concentrated  $\text{HNO}_3$  and  $\text{H}_2\text{SO}_4$  reacts on benzene at 300 K?

a) Sulphonation

b) Nitration

c) Hydrogenation

d) Dehydration

322. Consider the following reaction:






a) Benzene

b) Toluene

c) Benzaldehyde

d) Benzoic acid

323. The shortest C—O bond order exists in:



324. Between *p*-nitrophenol and salicylaldehyde, solubility in base is:

a) Almost nil in both cases

b) Higher in *p*-nitrophenol

c) Higher for salicylaldehyde

d) Equal in nature

325. (+) and (-) forms of optically active compounds are different in

a) Boiling points

b) Melting points

c) Specific gravity

d) Specific rotation

326. Benzene on treatment with dry HCN and HCl in presence of anhy.  $\text{AlCl}_3$  followed by hydrolysis forms:

a) Chlorobenzene

b) Benzoic acid

c) Benzaldehyde

d) Cyanobenzene

327. In which of the following compounds does the central atom obey EAN rule?



d) All of these

328. Pick the correct name of  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$

a) Chloropentammine cobalt (III) chloride

b) Chloropentammine cobalt (III)

c) Chloropentammine cobalt (II) chloride



d) Pentammine chloro cobalt(III) chloride

329. The geometry of  $\text{Ni}(\text{CO})_4$  and  $\text{Ni}(\text{PPh}_3)_2\text{Cl}_2$  are

- a) Square planar and tetrahedral respectively  
c) Tetrahedral and square planar respectively

- b) Both tetrahedral  
d) Both square planar

330. Select pair of chain isomers from the following



- a) I and II                      b) II and III                      c) I and IV                      d) II and III

331. Which ligand produces a high crystal field splitting (a strong ligand field)?

- a) CO                              b)  $\text{NO}_2^-$                               c)  $\text{CN}^-$                               d) All are correct

332. Benzene reacts with *n*-propyl chloride in the presence of anhydrous  $\text{AlCl}_3$  to give predominantly:

- a) Isopropyl benzene  
b) No reaction  
c) *n*-propylbenzene  
d) 3-propyl-1-chlorobenzene

333. Which of the following coordination compounds would exhibit optical isomerism?

- a) Pentaamminenitrocobalt (III) iodide                      b) Diamminedinitroplatinum (II)  
c) *trans*-dicyanobis (ethylenediamine)                      d) Tris-(ethylenediamine) cobalt(III) bromide

334. What is the magnetic moment of  $\text{K}_3[\text{FeF}_6]$ ?

- a) 3.87 BM                      b) 4.89 BM                      c) 5.91 BM                      d) 6.92 BM

335. The EAN of Cr in  $[\text{Cr}(\text{SCN})_6]^{3-}$  is:

- a) 35                              b) 33                              c) 34                              d) 37

336. Which has maximum paramagnetic character?

- a)  $[\text{Fe}(\text{CN})_6]^{4-}$                       b)  $[\text{Cu}(\text{H}_2\text{O})_4]^{2+}$                       c)  $[\text{Cu}(\text{NH}_3)_4]^{2+}$                       d)  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$

337. Phenol, when it first reacts with concentrated sulphuric acid and then with concentrated nitric acid, gives:

- a) Nitrobenzene  
b) 2, 4, 6-trinitrobenzene  
c) *o*-nitrophenol  
d) *p*-nitrophenol

338. Activation of benzene ring by  $-\text{NH}_2$  in aniline can be reduced by treating with:

- a) Dil. HCl                      b) Ethyl alcohol                      c) Acetic acid                      d) Acetyl chloride

339. Sulphonation of benzoic acid produces mainly:

- a) *o*-sulphobenzoic acid  
b) *m*-sulphobenzoic acid  
c) *p*-sulphobenzoic acid  
d) *o-p*-disulphobenzoic acid

340. The IUPAC name for the complex  $[\text{Co}(\text{NO}_2)(\text{NH}_3)_5]\text{Cl}_2$  is

- a) Nitrito -N- pentamminecobalt (III) chloride                      b) Nitrito -N- pentamminecobalt (II) chloride  
c) Pentammine nitrito-N- cobalt (II) chloride                      d) Pentaammine nitrito-N- cobalt (III) chloride

341. The ionisation isomer of  $[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}(\text{NO}_2)]\text{Cl}$  is

- a)  $[\text{Cr}(\text{H}_2\text{O})_4(\text{O}_2\text{N})]\text{Cl}_2$                       b)  $[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}_2](\text{NO}_2)$   
c)  $[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}(\text{ONO})]\text{Cl}$                       d)  $[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}_2(\text{NO}_2)] \cdot \text{H}_2\text{O}$

342. Salicylic acid, aspirin, nylon, plastics and picric acid have a common raw material, namely:

- a) Methane                      b) Formic acid                      c) Phenol                      d) Alcohol

343. Ulmann's reaction is used for the preparation of:

- a) Diphenyl                      b) Iodobenzene                      c) Toluene                      d) Naphthalene

344. Which of the following statements is/are incorrect for *D* - (+) -glyceradehyde?


- a) The symbol *D* not indicates the dextrorotatory nature of the compound

- b) The sign (+) indicates the dextrorotatory nature of the compound  
 c) The symbol *D* indicates that hydrogen atom lies left to the chiral centre in the Fischer projection diagram  
 d) The symbol *D* indicates that hydrogen atom lies right to the chiral centre in the Fischer projection diagram

345. Complexes with  $\text{CN}^-$  ligands are usually:

- a) High spin complexes    b) Low spin complexes    c) Both (a) and (b)    d) None of these

346. The IUPAC of




- a) 2-cyclopentyl propane    b) 1, 1-dimethyl-1-cyclopentyl methane  
 c) 1-(1-methyl) ethyl cyclopentane    d) None of the above

347. Which ion is paramagnetic?

- a)  $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$     b)  $[\text{Fe}(\text{CN})_6]^{4-}$     c)  $[\text{Ni}(\text{CO})_4]$     d)  $[\text{Ni}(\text{CN})_4]^{2-}$

348.



- Configuration of \_\_\_\_\_ are  
 a) *R, R*    b) *R, S*    c) *S, S*    d) *S, R*

349. Dow process is used for the conversion of chlorobenzene to:

- a) Benzene    b) Nitrobenzene    c) Phenol    d) Gammexane

350. Phenolphthalein is produced on heating phthalic anhydride and conc. sulphuric acid with:

- a) Salicylic acid    b) Phenol    c) Phenacetin    d) Phenanthrene


351. Benzene is converted to toluene by:

- a) Friedel-Crafts reaction  
 b) Grignard reaction  
 c) Wurtz reaction  
 d) Perkin's reaction

352. The number of ions formed when hexamine copper (II) sulphate is dissolved in water is?

- a) 1    b) 2    c) 4    d) 6

353. In a set of reactions *m*-bromobenzoic acid gave a product *D*, Identify the product *D*:



- a)    b)    c)    d)


354. In  $[\text{Cr}(\text{C}_2\text{O}_4)_3]^{3-}$ , the isomerism shown is:

- a) Ligand    b) Optical    c) Geometrical    d) Ionization

355. The hybridization of Fe in  $\text{K}_4[\text{Fe}(\text{CN})_6]$  complex is:

- a)  $d^2 sp^2$     b)  $d^2 sp^3$     c)  $dsp^2$     d)  $sp^3$

356.



- a) Hex-3-yn-5-ene    b) Hex-5-en-3-yne    c) Hex-3-yn-1-ene    d) Hex-1-en-3-yne

357. Nickel metal is in highest oxidation state in:

- a)  $\text{Ni}(\text{CO})_4$                       b)  $\text{K}_2\text{NiF}_6$                       c)  $[\text{Ni}(\text{NH}_3)_6](\text{BF}_4)_2$                       d)  $\text{K}_4[\text{Ni}(\text{CN})_6]$

358. Which of the following complexes show six coordination number?

- a)  $[\text{Zn}(\text{CN})_4]^{2-}$                       b)  $[\text{Ni}(\text{NH}_3)_4]^{2+}$                       c)  $[\text{Cu}(\text{CN})_4]^{2-}$                       d)  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$

359. Which of the following statements is wrong?

- a) The IUPAC name of alkenes ends with suffix-ene  
 b) The IUPAC name of alkynes ends with suffix-yne  
 c) The IUPAC name of acid amide is alkanamide  
 d) The substituents get lower number in comparison to principal functional group

360. The possible number of isomers for the complex  $[\text{MCl}_2\text{Br}_2]\text{SO}_4$  is:

- a) 1                                      b) 2                                      c) 4                                      d) 5

361.  $\text{K}_3[(\text{Al})(\text{C}_2\text{O}_4)_3]$  is called

- a) Potassium aluminium (III) oxalate                      b) Potassium alumino oxalate  
 c) Potassium trioxalato aluminate (VI)                      d) Potassium trioxalato aluminate (III)




362. In  $\text{Fe}(\text{CO})_5$ , the Fe — C bond possesses

- a)  $\pi$  — Character only                      b) Both  $\sigma$  and  $\pi$  — characters  
 c) Ionic characters                      d)  $\sigma$  — Character only

363. The reaction,  $[\text{Fe}(\text{CNS})_6]^{3-} \rightarrow [\text{FeF}_6]^{3-}$  taken place with

- a) Decrease in magnetic moment                      b) Increase in magnetic moment  
 c) Decrease in coordination number                      d) Increase in coordination number

364. Which chloro derivative of benzene among the following would undergo hydrolysis most readily with aqueous NaOH to furnish the corresponding hydroxyl derivative?



d)  $\text{C}_6\text{H}_5\text{Cl}$

365. Some salts although containing two different metallic elements give test for only one of them in solution. Such salts are:

- a) Complex salts                      b) Double salts                      c) Normal salts                      d) None of these

366. Mixture X = 0.02 mole of  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Br}$  and 0.02 mole of  $[\text{Co}(\text{NH}_3)_5\text{Br}]\text{SO}_4$  was prepared in 2 litre of solution.

1 litre of mixture X + excess  $\text{AgNO}_3 \rightarrow Y$ .

1 litre of mixture X + excess  $\text{BaCl}_2 \rightarrow Z$ .

No. of moles of Y and Z are.

- a) 0.01, 0.01                      b) 0.02, 0.01                      c) 0.01, 0.02                      d) 0.02, 0.02

367. The hybridization of central metal ion and shape of Wilkinson's catalyst is

- a)  $sp^3d$ , trigonal bipyramidal                      b)  $sp^3$ , tetrahedral  
 c)  $dsp^2$ , square planar                      d)  $d^2sp^2$ , octahedral

368. The d-electron configurations of  $\text{Cr}^{2+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Fe}^{2+}$  and  $\text{Co}^{2+}$  are  $d^4$ ,  $d^5$ ,  $d^6$  and  $d^7$  respectively. Which one of the following will exhibit minimum paramagnetic behaviour?

- a)  $[\text{Cr}(\text{H}_2\text{O})_6]^{2+}$   
 b)  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$   
 c)  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$   
 d)  $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$   
 (At. Nos. Cr = 24, Mn = 25, Fe = 26, Co = 27)

369. An enantiomerically pure acid is treated with racemic mixture of an alcohol having one chiral carbon. The ester formed will be
- a) Optically active mixture                      b) Pure enantiomer  
c) *meso* compound                                      d) Racemic mixture

370. Which of the following ring is most strained?

- a) Cyclohexane                      b) Cyclopentane                      c) Cyclobutane                      d) Cyclopropane

371. Formylchloride has not been prepared so far. Which can function as formylchloride in formylation?

- a) HCHO + HCl                      b) HCOOCH<sub>3</sub> + HCl                      c) CO + HCl                      d) HCONH<sub>2</sub> + HCl


372. In hexacyanomanganate (II) ion the Mn-atom assumes  $d^2sp^3$ -hybrid state. The number of unpaired electrons in the complex is:

- a) 1                      b) 2                      c) 3                      d) 0


373. Which one of the following does not give a white precipitate with silver nitrate solution?

- a) [Co(NH<sub>3</sub>)<sub>6</sub>]Cl<sub>3</sub>                      b) [Co(NH<sub>3</sub>)<sub>5</sub>Cl]Cl<sub>2</sub>                      c) [Co(NH<sub>3</sub>)<sub>4</sub>Cl<sub>2</sub>]Cl                      d) [Co(NH<sub>3</sub>)<sub>3</sub>Cl<sub>3</sub>]

374. In a set of reactions, ethyl benzene yielded a product *D*.



*D* would be:



375. The oxidation number of Pt in [Pt(C<sub>2</sub>H<sub>4</sub>)Cl<sub>3</sub>] is

- a) +1                      b) +2                      c) +3                      d) +4


376. Among [Fe(H<sub>2</sub>O)<sub>6</sub>]<sup>3+</sup>, [Fe(CN)<sub>6</sub>]<sup>3-</sup>, [FeCl<sub>6</sub>]<sup>3-</sup> species, the hybridization state of the Fe atom are, respectively

- a)  $d^2sp^3, d^2sp^3, sp^3d^2$                       b)  $sp^3d^2, d^2sp^3, d^2sp^3$                       c)  $sp^3d^2, d^2sp^3, sp^3d^2$                       d) None of these

377. Of the following complex ions, which is diamagnetic in nature?

- a) [CoF<sub>6</sub>]<sup>3-</sup>                      b) [NiCl<sub>4</sub>]<sup>2-</sup>                      c) [Ni(CN)<sub>4</sub>]<sup>2-</sup>                      d) [CuCl<sub>4</sub>]<sup>2-</sup>

378. The IUPAC name of compound



- a) 2-methoxycarbonylbenzoic acid                      b) Methyl-2-carboxy benzoate  
c) 2-carboxy phenyl ethanoate                      d) *o*-carboxyphenyl acetate

379. Which of the following are produced from coal-tar?

- a) Synthetic dyes                      b) Drugs                      c) Perfumes                      d) All of these


380. Chlorine is least reactive in:

- a) CH<sub>3</sub>Cl                      b) CH<sub>2</sub>=CHCl                      c) C<sub>6</sub>H<sub>5</sub>Cl                      d) C<sub>2</sub>H<sub>5</sub>Cl



381. Correct IUPAC name of compound


(CH<sub>3</sub>)<sub>2</sub>C(CH<sub>2</sub>CH<sub>3</sub>)CH<sub>2</sub>CH(Cl)CH<sub>3</sub> is

- a) 5-chloro-3,3-dimethylhexane                      b) 3-chloro-2-ethyl-2-methylpentane  
c) 2-chloro-4-ethyl-4-methylpentane                      d) None of the above




In the above reaction 'X' stands for:

- a)  $\text{NH}_2$                       b) Cl                      c)  $\text{SnCl}_2$                       d)  $\text{NH}_3\text{Cl}^+$
383. Which follows EAN rule?  
 a)  $\text{Fe}(\text{CO})_5$                       b)  $\text{Ni}(\text{CO})_4$                       c)  $\text{K}_4[\text{Fe}(\text{CN})_6]$                       d) All are correct
384. Which one is bidentate ligand?  
 a)  $\text{C}_2\text{O}_4^{2-}$                       b)  $\text{NH}_2 \cdot \text{CH}_2 \cdot \text{CH}_2 \cdot \text{NH}_2$                       c) Both (a) and (b)                      d) None of these
385. The reagent used for conversion of benzene diazonium chloride to benzene is:  
 a)  $\text{H}_3\text{PO}_2 + \text{H}_2\text{O}$                       b)  $\text{Na}_2\text{SnO}_2 + \text{NaOH}$                       c)  $\text{C}_2\text{H}_5\text{OH}$                       d) All of these
386. Which will not give the usual test for iron?  
 a)  $\text{K}_2\text{Fe}_2(\text{SO}_4)_4 \cdot 24\text{H}_2\text{O}$   
 b)  $(\text{NH}_4)_2\text{Fe}(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$   
 c)  $\text{K}_3[\text{Fe}(\text{CN})_6]$   
 d)  $\text{Fe}_2(\text{SO}_4)_3$
387.  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Br}$  and  $[\text{Co}(\text{NH}_3)_5\text{Br}]\text{SO}_4$  are a pair of ..... isomers.  
 a) Ionisation                      b) Ligand                      c) Coordination                      d) Hydrate
388. The first organic compound prepared in the laboratory was  
 a) Acetic acid                      b) Acetylene                      c) Urea                      d) Methane
389. Aniline on heating with conc.  $\text{H}_2\text{SO}_4$  at 460 K gives:  
 a) Aniline sulphate  
 b) Benzene sulphonic acid  
 c) Sulphanilic acid  
 d) None of the above
390. Which of the following statements regarding phenols is not correct?  
 a) Phenols are stronger acid than water and alcohols  
 b) Phenols are weaker acids than carboxylic acids  
 c) Phenols are soluble in both aqueous  $\text{NaOH}$  and aqueous  $\text{NaHCO}_3$   
 d) Phenoxide ions are more stable than the corresponding phenols
391. Which would decolourise cold, aq. potassium permanganate solution?  
 a) Benzoic acid                      b) Cinnamic acid                      c) *p*-toluic acid                      d) *m*-toluic acid
392. The magnetic moment of  $\text{K}_3[\text{Fe}(\text{CN})_6]$  is found to be 1.7 BM. How many unpaired electron (s) is/are present per molecule?  
 a) 1                      b) 2                      c) 3                      d) 4
393. The IUPAC name of the compound  
 is  
 a) Bicyclo [2,5,0] nonane                      b) Bicyclo [5,0,2] nonane  
 c) Bicyclo [5,2,0] nonane                      d) Bicyclo [0,2,5] nonane
394. The IUPAC name of the compound  
 is  
 a) 2-oxocyclohexane-1-carboxylic acid                      b) Cyclohexane-2-oxo-1-carboxylic acid  
 c) 6-oxocyclohexane-1-carboxylic acid                      d) None of the above
395. The IUPAC name of



is

- a) Spiro [3.2.1] octane    b) Bicyclo [3.2.2] octane    c) Bicyclo [3.2.1] octane    d) None of these
396. Which of the following deactivates benzene substitution?  
 a)  $-NHR$     b)  $-OH$     c)  $-OR$     d)  $-COOR$
397. Aniline, chloroform and alc. KOH on heating give:  
 a) Phenyl isocyanide    b) Phenyl cyanide    c) Chlorobenzene    d) Phenol
398. In the chemical reactions,




the compounds "A" and "B" respectively are :

- a) Nitrobenzene and chlorobenzene  
 b) Nitrobenzene and fluorobenzene  
 c) Phenol and benzene  
 d) Benzenediazonium chloride and fluorobenzene
399. The incorrect statement for IUPAC system of nomenclature is  
 a) In an organic compound, the longest carbon chain is always selected for assigning the root word  
 b) There is no compound with the name 3-ethyl pentane  
 c) Out of  $-NH_2$  and  $-OH$  groups present in an organic compound,  $-NH_2$  is treated as substituent  
 d) Different alkyl groups are written alphabetically while, writing the IUPAC name
400. When sodium benzene sulphonate is fused with sodium hydroxide (solid), followed by hydrolysis the product formed is:  
 a) Benzene    b) Sod. phenoxide    c) Benzene thiophenol    d) Phenol
401. The correct order of stability of conformations of cyclohexane is  
 a) Chair > twist boat > boat    b) Twist boat > chair > boat  
 c) Boat > chair > twist boat    d) Boat > twist boat > chair
402. Phenol with dilute  $HNO_3$  gives:  
 a) *meta* and *para* nitrophenol  
 b) *ortho* and *para* nitrophenol  
 c) Trinitrophenol  
 d) *ortho* and *meta* nitrophenol
403. The increasing order of boiling points of compounds given below is:  
 (I) 1,2-dihydroxy benzene  
 (II) 1,3-dihydroxy benzene  
 (III) 1,4-dihydroxy benzene  
 (IV) Hydroxyl benzene  
 a) I < II < III < IV    b) I < II < IV < III    c) IV < I < II < III    d) IV < II < I < III
404. The pair of the compounds in which both the metals are in the higher possible oxidation state is  
 a)  $CrO_2Cl_2, MnO_4^-$     b)  $[Co(CN)_6]^{3-}, MnO_3$   
 c)  $TiO_3, MnO_2$     d)  $[Fe(CN)_6]^{3-}, [Co(CN)_6]^{3-}$
405. The number of ions given by  $K_2[PtCl_6]$  in aqueous solution is:  
 a) 2    b) 3    c) 4    d) Zero
406. Which of the following are functional isomers?  
 a)  $CH_3CH_2Cl$  and  $CH_3CH_2Br$     b)  $CH_3CHBr_2$  and  $CH_2Br_2 \cdot CH_2Br$   
 c)  $C_2H_5OC_2H_5$  and  $CH_3OC_3H_7$     d)  $CH_3CH_2CHO$  and  $CH_3-CH-CH_2$




407. Phenol is:  
 a) Strongly acidic                      b) Weakly acidic                      c) Strongly basic                      d) Weakly basic
408. The correct IUPAC name of  $KAl(SO_4)_2 \cdot 12H_2O$  is:  
 a) Aluminium potassium sulphate-12-water  
 b) Potassium aluminium(III) sulphate-12-water  
 c) Potassium aluminate(III) sulphatehydrate  
 d) Aluminium(III) potassium sulphate hydrate-12

409. A complex shown below can exhibit:



- a) Optical isomerism only  
 b) Geometrical isomerism only  
 c) Both optical and geometrical isomerism  
 d) None of the above
410. The IUPAC name of the complex  $[Co(NH_3)_4Cl_2]Cl$  is  
 a) Dichloro tetraammine cobalt (III) chloride                      b) Tetraammine dichloro cobalt(III) chloride  
 c) Tetraammine dichloro cobalt (II) chloride                      d) Tetraammine dichloro cobalt (IV) chloride
411. The correct decreasing order of their reactivity towards hydrolysis is:

(i)  $C_6H_5COCl$



- a) (i) > (ii) > (iii) > (iv)                      b) (iv) > (ii) > (i) > (iii)                      c) (ii) > (iv) > (i) > (iii)                      d) (ii) > (iv) > (iii) > (i)
412. Nitrobenzene is generally used for:  
 a) Preparing shoe polish                      b) Preparing floor polish                      c) Preparing aniline                      d) All of these
413. In the coordination compound,  $K_4[Ni(CN)_4]$ , the oxidation state of nickel is  
 a) -1                      b) 0                      c) +1                      d) +2
414. Salicylic acid as compared to benzoic acid:  
 a) Is more acidic                      b) Has same acidity                      c) Has less acidity                      d) None of these
415. Which ligand is expected to be bidentate?  
 a)  $C_2O_4^{2-}$                       b)  $CH_3C \equiv N$                       c)  $Br^-$                       d)  $CH_3NH_2$
416. Which one of the following is most reactive towards aqueous NaOH?  
 a)  $C_6H_5Cl$                       b)  $C_6H_5CH_2Cl$                       c)  $C_6H_5Br$                       d)  $BrC_6H_4Br$
417. Which is not an aromatic compound?  
 a) Pyridine                      b) Naphthalene                      c) Xylene                      d) Cyclohexane
418. Which one of the following is wrongly matched?  
 a)  $[Cu(NH_3)_4]^{2+}$  – Square planar                      b)  $[Ni(CO)_4]$  – Neutral ligand  
 c)  $[Fe(CN_6)]^{3-}$  –  $sp^3 d^2$                       d)  $[Co(en)_3]^{3+}$  – Follows EAN rule

419. Stereoisomers have different

- a) Molecular formula                      b) Structural formula  
 c) Configuration                      d) Molecular mass

420. Which of the following will show optical isomerism?

- a)  $[\text{Cu}(\text{NH}_3)_4]^{2+}$
- b)  $[\text{ZnCl}_4]^{2-}$
- c)  $[\text{Cr}(\text{C}_2\text{O}_4)_3]^{3-}$
- d)  $[\text{Co}(\text{CN})_6]^{3-}$

421. A complex of cobalt has five ammonia molecules, one nitro group and two chlorine atoms for each cobalt atom. One mole of this compound produces three mole ions in aqueous solution which on treating with excess of  $\text{AgNO}_3$  give two mole of  $\text{AgCl}$ . The formula of the compound is:

- a)  $[\text{Co}(\text{NH}_3)_4\text{NO}_2\text{Cl}][(\text{NH}_3)_2\text{Co}(\text{NH}_3)_5\text{Cl}][\text{ClNO}_2]$
- b)  $[\text{Co}(\text{NH}_3)_5\text{NO}_2]\text{Cl}_2$
- c)  $[\text{Co}(\text{NH}_3)_5][(\text{NO}_2)_2\text{Cl}_2]$
- d)  $[\text{Co}(\text{NH}_3)_5][(\text{NO}_2)_2\text{Cl}_2]$

422. Which one group is trivalent in nature?

- a) Benzo
- b) Benzal
- c) Benzyl
- d) All of these

423. Benzene contains double bonds but does not give addition reactions because:

- a) Double bonds in benzene are strong
- b) Double bonds change their position rapidly
- c) Resonance lowers the energy of benzene molecule and leads to greater stabilization
- d) None of the above


424. Low spin complex of  $d^6$ -cation in an octahedral field will have the following energy:

- a)  $\frac{-12}{5}\Delta_0 + P$
- b)  $\frac{-12}{5}\Delta_0 + 3P$
- c)  $\frac{-2}{5}\Delta_0 + 2P$
- d)  $\frac{-2}{5}\Delta_0 + P$

( $\Delta_0$  = Crystal field splitting energy in an octahedral field,  $P$  = Electron pairing energy)

425.  $\text{C}_7\text{H}_8\text{O}$  show how many isomers?

- a) 2
- b) 3
- c) 4
- d) 5



The above structural formula refers to:

- a) BHC
- b) DNA
- c) DDT
- d) RNA

427. The compound



Have its IUPAC name as

- a) Octa dec-9-enoic acid
- b) Oleic acid
- c) Ethyl hexadec-9-enoic acid
- d) All of these

428. The type of isomerism present in nitropentaammine-chromium (III) chloride is :

- a) Optical
- b) Linkage
- c) Ionization
- d) polymerization




429. Which complex compound possesses  $sp^3d^2$  hybridisation?



- a)  $[\text{Fe}(\text{NH}_3)_6]^{3+}$
- b)  $[\text{Fe}(\text{CN})_6]^{4-}$
- c)  $[\text{Fe}(\text{CN})_6]^{3-}$
- d)  $[\text{Fe}(\text{Cl})_6]^{3-}$

430. Amongst the following carboxylic acids the strongest acid is:

- a) Benzoic acid
- b) *o*-methoxybenzoic acid
- c) *m*-nitrobenzoic acid
- d) *p*-nitrobenzoic acid

431. When EDTA solution is added to  $Mg^{2+}$  ion solution, then which of the following statements is not true?
- Four coordinate sites of  $Mg^{2+}$  are occupied by EDTA and remaining two sites are occupied by water molecules.
  - All six coordinate sites of  $Mg^{2+}$  are occupied.
  - $P^H$  of the solution is decreased.
  - Colourless  $[Mg - EDTA]^{2-}$  chelate is formed.
432. The energy difference between chair and the boat conformation of cyclohexane is
- 29.7 kJ
  - 44 kJ
  - 151 kJ
  - 36 kJ
433. Compounds having the same molecular formula but different properties are called
- Isotopes
  - Isobars
  - Isomers
  - Isomorphs
434.  $CH_3OCH_2CH_2CH_2OCH_2CH_3$  is
- Ethylmethylpropyl diether
  - Ethylmethoxypropyl ether
  - 3-ethoxy-1-methoxy propane
  - 1-ethoxy-3-methoxy propane
435. The benzene molecule contains:
- Six  $sp^2$ -hybridized carbons
  - Three  $sp^2$ -hybridized carbons
  - Six  $sp^3$ -hybridized carbons
  - Three  $sp^3$ -hybridized carbons
436. The correct order of stability of conformations of  $NH_2 - CH_2 - CH_2 - OH$  is
- Gauche > eclipsed > anti
  - Gauche > anti > eclipsed
  - Eclipsed > gauche > anti
  - Anti > eclipsed > gauche
437. The solubility of  $AgCN$  increases by the addition of  $KCN$  because of:
- Complex formation
  - Redox change
  - Salt formation
  - None of these
438. Alicyclic compounds are
- Aromatic cyclic compounds
  - Aliphatic cyclic compounds
  - Both (a) and (b)
  - None of the above
439. Which of the following compounds reacts slower than benzene in electrophilic bromination?
- $C_6H_5 - NO_2$
  - $C_6H_5 - NH_2$
  - $C_6H_5 - OH$
  - $C_6H_6 - CH_3$
440. The fraction of chlorine precipitated by  $AgNO_3$  solution from  $[Cu(NH_3)_5Cl]Cl_2$  is:
- 1/2
  - 2/3
  - 1/3
  - 1/4
441. Number of possible optical isomers in  $[Co(en)_2Cl_2]^+$  is
- 2
  - 3
  - 4
  - 6
442. Dimethyl glyoxime gives a red precipitate with  $Ni^{2+}$  which is used for its detection. To get this precipitate readily, the best pH range is
- < 1
  - 3 - 4
  - 9 - 11
  - 2 - 3
443. Predict the product:





444. Replacement of Cl of chlorobenzene to give phenol requires drastic conditions but chlorine of 2, 4-dinitrochlorobenzene is readily replaced because:

- a)  $\text{NO}_2$  makes the electron rich ring at *ortho* and *para* positions  
 b)  $\text{NO}_2$  withdraws electrons at *meta* position  
 c)  $\text{NO}_2$  donate electrons at *meta*-position  
 d)  $\text{NO}_2$  withdraws electrons at *ortho* and *para* positions

445. Salicylic acid on heating with soda lime forms:

- a) Phenol                                      b) Benzyl alcohol                                      c) Benzene                                      d) Benzoic acid





446. Which of the following is an organometallic compound?

- a)  $\text{Ti}(\text{C}_2\text{H}_5)_4$                                       b)  $\text{Ti}(\text{OC}_2\text{H}_5)_4$                                       c)  $\text{Ti}(\text{OCOCH}_3)_4$                                       d)  $\text{Ti}(\text{OC}_6\text{H}_5)_4$





447. Which kind of isomerism is exhibited by octahedral  $\text{Co}(\text{NH}_3)_4\text{Br}_2\text{Cl}$  ?

- a) Geometrical and ionisation                                      b) Geometrical and optical  
 c) Optical and ionisation                                      d) Geometrical only

448. Which of the following is the strongest base?



449. Which of the following will be aromatic?



450. The correct symbol relating the two Kekule structure of benzene is:

- a)  $\rightarrow$                                       b)  $\rightleftharpoons$                                       c)  $\leftrightarrow$                                       d)  $\rightleftharpoons$

451. Benzaldehyde can be obtained by the hydrolysis of:

- a) Benzyl chloride                                      b) Benzal chloride                                      c) Benzonitrile                                      d) Benzoic acid

452. Which of the following has an optical isomer?

- a)  $[\text{Co}(\text{en})(\text{NH}_3)_2]^{2+}$                                       b)  $[\text{Co}(\text{H}_2\text{O})_4(\text{en})]^{3+}$                                       c)  $[\text{Co}(\text{en})_2(\text{NH}_3)_2]^{3+}$                                       d)  $[\text{Co}(\text{NH}_3)_3\text{Cl}]^+$

453. Chromium carbonyl is:

- a)  $\text{Cr}(\text{CO})_4$                                       b)  $\text{Cr}(\text{CO})_5$                                       c)  $\text{Cr}(\text{CO})_6$                                       d) None of these

454. Which of the following reagents may be used to distinguish between phenol and benzoic acid?





- a) Aqueous NaOH                                      b) Tollen's reagent                                      c) Molisch reagent                                      d) Neutral  $\text{FeCl}_3$



455. Which of the following complex species do not involve  $d^2sp^3$ -hybridization?

- a)  $[\text{CoF}_6]^{3-}$                                       b)  $[\text{Co}(\text{NH}_3)_6]^{3+}$                                       c)  $[\text{Fe}(\text{CN})_6]^{3-}$                                       d)  $[\text{Cr}(\text{NH}_3)_6]^{3+}$

456. Which one of the following shows maximum value of paramagnetic behaviour?



473. Among the following, identify the species with an atom of +6 oxidation state:  
 a)  $[\text{MnO}_4]^-$                       b)  $[\text{Cr}(\text{CN})_6]^{3-}$                       c)  $[\text{NiF}_6]^{2-}$                       d)  $\text{CrO}_2\text{Cl}_2$
474. Which of the following alkanes contain primary, secondary, tertiary and quaternary carbon atom together?  
 a)  $(\text{CH}_3)_3\text{CH}$                       b)  $(\text{C}_2\text{H}_5)_3\text{CH}$                       c)  $(\text{CH}_3)_3\text{CCH}_2\text{CH}(\text{CH}_3)_2$                       d)  $(\text{CH}_3)_4\text{C}$
475. The hardness of water is estimated by:  
 a) Conductivity method                      b) EDTA method                      c) Titrimetric method                      d) Distillation method
476.  $\text{I}_2$  is stirred in between two liquids,  $\text{C}_6\text{H}_6$  and water. It:  
 a) Dissolves more in  $\text{C}_6\text{H}_6$   
 b) Dissolves more in  $\text{H}_2\text{O}$   
 c) Dissolve equally  
 d) Dissolves in neither  $\text{C}_6\text{H}_6$  nor water
477. The number of tertiary C-atoms in 2,2,4,4-tetra methyl pentane is  
 a) 1                      b) 2                      c) 3                      d) 4
478. Hydrogenation of benzoyl chloride in the presence of Pd on  $\text{BaSO}_4$  gives:  
 a) Benzyl alcohol                      b) Benzaldehyde                      c) Benzoic acid                      d) Phenol
479. The Clemmensen reduction of benzaldehyde gives:  
 a)  $\text{C}_6\text{H}_5\text{NH}_2$                       b)  $\text{C}_6\text{H}_5\text{OH}$                       c)  $\text{C}_6\text{H}_5\text{CH}_3$                       d)  $\text{C}_6\text{H}_5\text{COOH}$
480. Which of the following ligand has lowest  $\Delta_o$  value?  
 a)  $\text{CN}^-$                       b)  $\text{CO}$                       c)  $\text{F}^-$                       d)  $\text{NH}_3$
481. Which one of the following has an optical isomer?  
 (en=ethylenediamine)  
 a)  $[\text{Zn}(\text{en})(\text{NH}_3)_2]^{2+}$                       b)  $[\text{Co}(\text{en})_3]^{3+}$                       c)  $[\text{Co}(\text{H}_2\text{O})_4(\text{en})]^{3+}$                       d)  $[\text{Zn}(\text{en})_2]^{2+}$
482. Trichloroacetaldehyde,  $\text{CCl}_3\text{CHO}$  reacts with chlorobenzene in presence of sulphuric acid and produces:
- a) 
- b) 
- c) 
- d) 
483. Which fraction of coal-tar is rich in arene?  
 a) Light oil                      b) Heavy oil                      c) Green oil                      d) Middle oil
484. The coordination number and oxidation number of X in the following compound  $[\text{X}(\text{SO}_4)(\text{NH}_3)_5]\text{Cl}$  will be  
 a) 10 and 3                      b) 2 and 6                      c) 6 and 3                      d) 6 and 4
485. Benzyl chloride is formed by treating toluene with  $\text{Cl}_2$  in:  
 a) Presence of light  
 b) Absence of light  
 c) Treating benzene with anhy.  $\text{AlCl}_3$


- d) Treating benzene with  $\text{As}_2\text{S}_3$
486. Which complex cannot ionize in solution?  
 a)  $[\text{CoCl}_3(\text{NH}_3)_3]$       b)  $\text{K}_4(\text{Fe}(\text{CN})_6)$       c)  $\text{K}_2[\text{Pt}(\text{F}_6)]$       d)  $[\text{Pt}(\text{NH}_3)_6]\text{Cl}_4$
487.  $[\text{Ni}(\text{CN})_4]^{2-}$ ,  $[\text{MnBr}_4]^{2-}$  and  $[\text{CoF}_6]^{3-}$ , geometry, hybridisation and magnetic moment of the ions respectively, are  
 a) Tetrahedral, square planar, octahedral :  
 $sp^3, dsp^2, sp^3d^2$ : 5.9, 0, 4.9  
 b) Tetrahedral, square planar, octahedral :  
 $dsp^2, sp^3, sp^3d^2$ : 0, 5.9, 4.9  
 c) Square planar, tetrahedral, octahedral :  
 $dsp^2, sp^3, d^2sp^3$ : 5.9, 4.9, 0  
 d) Square planar, tetrahedral, octahedral :  
 $dsp^2, sp^3, sp^3d^2$ : 0, 5.9, 4.9
488. Ozonolysis of benzene gives:  
 a) 1 molecule of glyoxal  
 b) 2 molecules of glyoxal  
 c) 3 molecules of glyoxal  
 d) None of these
489. In benzene, C—C bond length is 1.39 Å; the C—H bond length is:  
 a) 1.39      b) 1.08      c) 1.54      d) 1.46
490. The IUPAC name of following compound is  
  
 a) N,N-dimethyl, 3-methyl pentan-3-amine      b) 3-N,N-dimethyl, 3-methyl pentanamine  
 c) 3-methyl-3-N, N-dimethyl pentane      d) 3-methyl-3-N, N-dimethyl butane
491. Which of the following may be used as food preservative?  
 a) Benzene  
 b) Ethylene  
 c) Sodium benzoate  
 d) Sodium metaaluminate
492. Which compound is formed when sodium phenoxide is heated with ethyl iodide?  
 a) Phenetole      b) Ethyl phenyl alcohol      c) Phenol      d) None of these
493. In metal carbonyl (organometallic) complexes, the M—C bond is  
 a) Ionic      b) Covalent with ionic character  
 c) Covalent      d) Coordinate covalent
494. Octahedral complex  
 is  
 a) *cis*      b) *trans*      c) *mer*      d) *fac*
495. The correct order of magnetic moments (spin only values in BM) among the following is  
 (Atomic no. Mn=25, Fe=26, Co=27)  
 a)  $[\text{MnCl}_4]^{2-} > [\text{CoCl}_4]^{2-} > [\text{Fe}(\text{CN})_6]^{4-}$       b)  $[\text{MnCl}_4]^{2-} > [\text{Fe}(\text{CN})_6]^{4-} > [\text{CoCl}_4]^{2-}$   
 c)  $[\text{Fe}(\text{CN})_6]^{4-} > [\text{MnCl}_4]^{2-} > [\text{CoCl}_4]^{2-}$       d)  $[\text{Fe}(\text{CN})_6]^{4-} > [\text{CoCl}_4]^{2-} > [\text{MnCl}_4]^{2-}$
496. Aniline and methyl amine can be differentiated by:  
 a) Diazotisation followed by coupling with phenol  
 b) Reaction with chloroform and aqueous solution of KOH  
 c) Reaction with  $\text{HNO}_2$

d) None of the above





497. The functional group present in cresols is:

- a) Alcoholic ( $-\text{OH}$ )      b) Aldehydic ( $-\text{CHO}$ )      c) Phenolic ( $-\text{OH}$ )      d) Carboxylic ( $-\text{COOH}$ )





498. In the reaction;



the structure of the product T is:



499. Which one of the following compounds is most acidic?




500. The most unstable configuration of cyclohexane is

- a) Boat      b) Chair      c) Twist boat      d) Half chair

501. In which compound synergic effect is present?

- a)  $[\text{Ni}(\text{CO})_4]$       b)  $[\text{NiCl}_4]^{2-}$       c)  $[\text{CuCl}_4]^{2-}$       d)  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$

502. The IUPAC name of the compound



- a) 4-amino-2-ethyl pent-1-ene      b) 2-ethyl pentan-4-amine  
c) Amino-4-pentene      d) 4-ethyl pent-4-en-2-amine


503. Aqua regia reacts with Pt to yield:

- a)  $\text{Pt}(\text{NO}_3)_4$       b)  $\text{H}_2[\text{PtCl}_6]$       c)  $\text{PtCl}_4$       d)  $\text{PtCl}_2$

504.  $\text{K}_3[\text{Al}(\text{C}_2\text{O}_4)_3]$  is called:

- a) Potassium aluminooxalate  
b) Potassium alumino(III) oxalate  
c) Potassium trioxalatoaluminate  
d) Potassium trioxalatoaluminate(III)

505. The IUPAC name of



- a) 6-oxo-1,2,2-trimethyl bicyclo [2.2.1] heptane      b) 1,7,7-trimethyl bicyclo [2.2.1] heptan-2-one  
c) 1,5,5-trimethyl bicyclo [2.1.1] hexane-2-one      d) 1,7,7-trimethyl bicyclo [2.1.2] heptan-2-one

506. Nitration of toluene takes place at:

- a) *ortho* position  
b) *meta* position  
c) *para* position  
d) Both *ortho* and *para* position

507. Estimation of calcium and magnesium is done by

- a) EDTA      b) Oxalate      c) Phosphate      d) None of these


508. How many enantiomer pairs are obtained by monochlorination of 2, 3-dimethyl butane?

- a) Four      b) Two      c) Three      d) One

509. Common reactions of benzene and its derivatives are:


- a) Electrophilic addition reactions  
b) Electrophilic substitution reactions  
c) Nucleophilic substitution reactions  
d) Nucleophilic addition reactions

510. The IUPAC name of the compound



- a) 1, 3, 5-triheptene      b) 2, 4, 6-triheptene  
c) 2, 4, 6-heptatriene      d) Hepta-1, 3, 5-triene

511. Name of compound



- a) 1, 2, 3-triformylpentane      b) Propane-1, 2, 3-tricarbaldehyde  
c) 3-formylpentane-1, 5-dial      d) Propane-1, 2, 3-trial

512. The attacking species in aromatic sulphonation is:

- a)  $\text{SO}_3$       b)  $\text{H}_3\text{SO}_4^+$       c)  $\text{HSO}_4$       d)  $\text{SO}_2^+$

513. Which one of the following compound does not react with bromine?

- a) Ethyl amine      b) Propene      c) Phenol      d) Chloroform

514. The magnetic moment (spin only) of  $[\text{Ni Cl}_4]^{2-}$  is

- a) 1.82 BM      b) 5.46 BM      c) 2.82 BM      d) 1.41 BM

515.



undergoes electrophilic substitution reaction preferentially :

- a) At position-2      b) At position-3      c) At position-4      d) At positions-2 and 4

516. Ionization of  $\text{K}[\text{Ag}(\text{CN})_2]$  will give:


- a)  $\text{K}^+$  and  $[\text{Ag}(\text{CN})_2]^-$  ion  
b) KCN and AgCN  
c)  $\text{K}^+$ ,  $\text{Ag}^+$ ,  $\text{CN}^-$   
d) None of the above


517. The coordination number and oxidation state of Cr in  $\text{K}_3[\text{Cr}(\text{C}_2\text{O}_4)_3]$  are respectively




- a) +6 and +3      b) 3 and 0      c) 4 and +2      d) 3 and +3

518. A complex of platinum, ammonia and chlorine produces four ions per molecule in the solution. The structure consistent with the observation is:  
 a)  $[\text{Pt}(\text{NH}_3)_4]\text{Cl}_4$       b)  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_4]$       c)  $[\text{Pt}(\text{NH}_3)_5\text{Cl}]\text{Cl}_3$       d)  $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}_2$
519. The type of magnetism exhibited by  $[\text{Mn}(\text{H}_2\text{O})^{2+}]$  ion is  
 a) Paramagnetism      b) Diamagnetism      c) Both (a) and (b)      d) None of these
520. According to effective atomic number rule the central metal acquires:  
 a) Inert gas configuration  
 b) Duplet  
 c) Octet  
 d) Quartet
521.  $\text{K}_3\text{CoF}_6$  is high spin complex. What is the hybrid state of Co-atom in this complex?  
 a)  $sp^3d$       b)  $sp^3d^2$       c)  $d^2sp^3$       d)  $dsp^2$

522. The correct structure of ethylenediaminetetraacetic acid (EDTA) is



523.  $[\text{Co}(\text{NH}_3)_5\text{Br}]\text{SO}_4$  and  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Br}$  are examples of which type of isomerism?  
 a) Linkage      b) Optical      c) Geometrical      d) Ionisation
524. The coordination number of a central metal atom in a complex is determined by  
 a) The number of ligands around a metal ion bonded by  $\sigma$  – bonds  
 b) The number of ligands around a metal ion bonded by  $\pi$  –bonds  
 c) The number of ligands around a metal ion bonded by  $\sigma$  –and  $\pi$  – bonds both  
 d) The number of only anionic ligands bonded to the metal ion
525. Action of benzoic acid with hydrazoic acid in presence of  $\text{N}_3\text{H}$  gives:  
 a) Aniline      b) Benzamide      c) Phenyl cyanide      d) All of these
526. Which ion shows usually the coordination number 6?  
 a)  $\text{Cr}^{3+}$       b)  $\text{Fe}^{3+}$       c)  $\text{Fe}^{2+}$       d) All of these
527. Which of the following represents hexadentate ligand?  
 a) 2, 2-bipyridyl      b) DMG      c) Ethylenediamine      d) None of these
528. Nitrobenzene can be prepared from benzene by using a mixture of conc.  $\text{HNO}_3$  and conc.  $\text{H}_2\text{SO}_4$ . In the mixture, nitric acid acts as a/an:  
 a) Catalyst      b) Reducing agent      c) Acid      d) Base
529. The value of the 'spin only' magnetic moment for one of the following configurations is 2.84 BM. The correct one is  
 a)  $d^5$  (in strong ligand field)      b)  $d^3$  (in weak as well as strong ligand fields)  
 c)  $d^4$  (in weak ligand field)      d)  $d^4$  (in strong ligand field)
530. The IUPAC name of the compound  
 is  
 a) 6-(3-oxobutyl) cyclohexan-1-one      b) 6-(2-oxobutyl) cyclohexan-1-one  
 c) 2-(3-oxobutyl) cyclohexan-1-one      d) 2-(2-oxobutyl) cyclohexan-1-one
531. Hybridisation, shape and magnetic moment of  $[\text{Ni}(\text{CN})_4]^{2-}$  ion

- a)  $dsp^2$ , square planar, zero  
 c)  $sp^2d^2$ , octahedral, zero
- b)  $dsp^2$ , square planar, 1.73  
 d)  $d^2sp^3$ , octahedral, 1.73
532. Choose the IUPAC name of 
- a) Dicyclobutane  
 c) Bicyclo [2.2.1] hexane
- b) Bicyclo [2.2.0] hexane  
 d) None of these
533. Which of the following is a heterocyclic compound?
- a) Phenanthrene  
 b) Thiophene  
 c) Phenol  
 d) Aniline
534.  $[Sc(H_2O)_6]^{3+}$  ion is
- a) Colourless and diamagnetic  
 c) Colourless and paramagnetic
- b) Coloured and octahedral  
 d) Coloured and paramagnetic
535. Benzene reacts with  $CH_3Cl$  in the presence of anhydrous  $AlCl_3$  to form:
- a) Xylene  
 b) Toluene  
 c) Chlorobenzene  
 d) Benzylchloride
536. The magnetic moment of  $[Co(NH_3)_6]Cl_3$  is
- a) 1.73  
 b) 2.83  
 c) 6.6  
 d) Zero
537. The correct order of reactivity towards electrophilic substitution is:
- a) Phenol > Benzene > Chlorobenzene > Benzoic acid  
 b) Benzoic acid > Chlorobenzene > Benzene > Phenol  
 c) Phenol > Chlorobenzene > Benzene > Benzoic acid  
 d) Benzoic acid > Phenol > Benzene > Chlorobenzene
538. The product formed by the reaction of  $C_6H_5CN$  and  $CH_2N_2$  is:
- a)  b)  c)  d) None of these
539. Increasing order of expected enol content
- a)  $CH_3COCH_2CHO > CH_3COCH_3 > CH_3CHO > CH_3COCH_2COCH_3$   
 b)  $CH_3COCH_2COCH_3 > CH_3COCH_2CHO > CH_3COCH_3 > CH_3CHO$   
 c)  $CH_3CHO > CH_3COCH_3 > CH_3COCH_2CHO > CH_3COCH_2COCH_3$   
 d)  $CH_3COCH_3 > CH_3COCH_2COCH_3 > CH_3CHO > CH_3COCH_2CHO$
540. Out of the following the metal which forms polynuclear carbonyl is:
- a) Na  
 b) Mg  
 c) Mn  
 d) All of these
541. Picric acid and benzoic acid can be distinguished by:
- a) Aqueous  $NaHCO_3$   
 b) Aqueous NaOH  
 c) Aqueous  $FeCl_3$   
 d) Aqueous  $Na_2CO_3$
542. The compound having the lowest oxidation state of iron is
- a)  $K_4Fe(CN)_6$   
 b)  $K_2FeO_4$   
 c)  $Fe_2O_3$   
 d)  $Fe(CO)_5$
543. The name of  $[Pt(NH_3)_4Cl_2]^{2+}$ ,  $[PtCl_4]^{2-}$  is
- a) Tetramminedichloroplatinum(IV) tetrachloro platinate(II)  
 b) Dichloroplatinum (IV) tetrachloroplatinate  
 c) Tetrachloroplatinum (II) tetrammineplatinate  
 d) Tetrachloroplatinum (II) dichlorotetraammine platinate
544. *m*-dihydroxybenzene is also called:
- a) Catechol  
 b) Resorcinol  
 c) Quinol  
 d) Pyrogallol
545. The ion which exhibits green colour
- a)  $Cu^{2+}$   
 b)  $Mn^{2+}$   
 c)  $Co^{2+}$   
 d)  $Ni^{2+}$
546.  $X \xrightarrow{Cl_2}$  Benzotrichloride  $\xrightarrow{\text{Hydrolysis}}$  Y  
 X and Y respectively are:
- a) Benzene, benzaldehyde  
 b) Toluene, benzaldehyde  
 c) Toluene, benzoic acid  
 d) Benzene, benzoic acid


547. Geometrical isomerism is found in coordination compounds having coordination number:

- a) 2                                      b) 3                                      c) 4 (tetrahedral)                      d) 6

548. Which one of the following complexes is not expected to exhibit isomerism?


- a)  $[\text{Ni}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}$     b)  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$                       c)  $[\text{Ni}(\text{NH}_3)_2\text{Cl}_2]$                       d)  $[\text{Ni}(\text{en})_3]^{2+}$

549. The correct acidity order of the following is:



- a) (III) > (IV) > (II) > (I)  
 b) (IV) > (III) > (I) > (II)  
 c) (III) > (II) > (I) > (IV)  
 d) (II) > (III) > (IV) > (I)

550. Identify 'Z' in the reaction;



- a)
- b)
- c)
- d)


551. Pure aniline is a:

- a) Brown coloured liquid  
 b) Colourless liquid  
 c) Brown coloured solid  
 d) Colourless solid

552. Aromatic compounds undergo most easily:


- a) Nucleophilic substitution  
 b) Electrophilic substitution  
 c) Nucleophilic addition  
 d) Electrophilic addition





553. The colour of  $\text{CoCl}_3 \cdot 5\text{NH}_3 \cdot \text{H}_2\text{O}$  is:  
 a) Orange yellow      b) Orange      c) Green      d) Pink
554. The value of  $x$  on the  $[\text{Ni}(\text{CN})_4]^x$  is:  
 a) +2      b) -2      c) Zero      d) +4
555. Complexes with halide ligands are generally:  
 a) High spin complexes      b) Low spin complexes      c) Both (a) and (b)      d) None of these
556. The hybridization involved in  $[\text{CoF}_6]^{3-}$  is:  
 a)  $d^2sp^3$       b)  $d^3sp^2$       c)  $dsp^3$       d)  $sp^3d^2$



Will have the name

- a) N-ethyl-N-methylethanamine      b) N,N-diethylmethanamine  
 c) N,N-diethylethanamide      d) None of the above
558. The oxidation state of Fe in the brown ring complex  $[\text{Fe}(\text{H}_2\text{O})_5\text{NO}]\text{SO}_4$  is  
 a) +3      b) 0      c) +2      d) +1
559. The metal ion in complex A has EAN identical to the atomic number of krypton. A is  
 (At. no. of Cr=24, Fe=26, Pd=46)  
 a)  $[\text{Pd}(\text{NH}_3)_6]\text{Cl}_4$       b)  $[\text{Cr}(\text{NH}_3)_5\text{Cl}]\text{SO}_4$       c)  $\text{Na}_4[\text{Fe}(\text{CN})_6]$       d)  $\text{K}_3[\text{Fe}(\text{CN})_6]$
560. Which one of the following is expected to exhibit optical isomerism [en = ethylenediamine]?  
 a) *trans* -  $[\text{Co}(\text{en})_2\text{Cl}_2]$       b) *cis* -  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$   
 c) *cis* -  $[\text{Co}(\text{en})_2\text{Cl}_2]$       d) *Trans* -  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$
561. What is the magnetic moment of  $\text{K}_3[\text{FeF}_6]$ ?  
 a) 5.91 BM      b) 4.89 BM      c) 3.87 BM      d) 6.92 BM
562. Identify 'Y' in the change;



- a)       b)   
 c)       d) 

563. Among the following statements on the nitration of aromatic compounds, the false one is:  
 a) The rate of nitration of benzene is almost the same as that of hexadeuterobenzene  
 b) The rate of nitration of toluene is greater than that of benzene  
 c) The rate of nitration of benzene is greater than that of hexadeuterobenzene  
 d) Nitration is an electrophilic substitution reaction
564. The bond length of C—O bond in carbon monoxide is 1.128 Å. The C—O bond in  $\text{Fe}(\text{CO})_5$  is:  
 a) 1.15 Å      b) 1.128 Å      c) 1.72 Å      d) 1.118 Å
565. Which one is not correct for homologous series?  
 a) All members are represented by same general formula  
 b) All members have same chemical properties  
 c) All members have same physical properties  
 d) All members have same functional group



is named in IUPAC as

- a) 2, 3-dimethyl bicyclo [2.2.1] hept-5-ene      b) 1, 2-dimethyl bicyclo [2.2.1] hept-4-ene  
 c) 5, 6-dimethyl bicyclo [2.2.1] hept-2-ene      d) 4, 5-dimethyl bicyclo [2.2.1] hept-1-ene
567. Ferric ion forms a prussian blue coloured solution due to the formation of:

- a)  $K_4[Fe(CN)_6]$       b)  $Fe(CNS)_3$       c)  $Fe_4[Fe(CN)_6]_3$       d)  $K_3[Fe(CN)_6]$
568. What is the magnetic moment of  $[FeF_6]^{3-}$ ?  
 a) 5.92      b) 5.49      c) 2.34      d) 4
569. Which of the following can exhibit geometrical isomerism?  
 a)  $[MnBr_4]^{2-}$       b)  $[Pt(NH_3)_3Cl]^+$       c)  $[PtCl_2 \cdot P(C_2H_5)_3]_2$       d)  $[Fe(H_2O)_5NO]^{2+}$
570. A compound contains 2 dissimilar asymmetric C-atoms. The number of optical isomers are  
 a) 2      b) 3      c) 4      d) 5
571. Coordination number of Ni in  $[Ni(C_2O_4)_3]^{4-}$  is:  
 a) 3      b) 6      c) 4      d) 5
572. Which compound exhibits optical isomerism?  
 a) Pentaamminenitrocobalt (III) iodide  
 b) Diamminedichloroplatinum (II)  
 c) *Trans*-dicyano-bis-(ethylenediamine) chromium (III) chloride  
 d) Tris-(ethylenediamine)cobalt (III) bromide

573. Ruthenium carbonyl is:  
 a)  $Ru(CO)_4$       b)  $Ru(CO)_5$       c)  $Ru(CO)_8$       d)  $Ru(CO)_6$
574. Oxidation state of nitrogen is incorrectly given for

Compound	Oxidation state
----------	-----------------

- |                         |    |              |    |
|-------------------------|----|--------------|----|
| a) $[Co(NH_3)_5Cl]Cl_2$ | 0  | b) $NH_2OH$  | -1 |
| c) $(N_2H_5)_2SO_4$     | +2 | d) $Mg_3N_2$ | -3 |

575. Which of the following can participate in linkage isomerism?  
 a)  $NH_3$       b)  $H_2O$       c)  $H_2NCH_2CH_2NH_2$       d)  $NO_2^-$
576. *Ortho*-nitrophenol is less soluble in water than *p*- and *m*-nitrophenols because:  
 a) *o*-nitrophenol shows intramolecular H-bonding  
 b) *o*-nitrophenol shows intermolecular H-bonding  
 c) Melting point of *o*-nitrophenol is lower than those of *m*- and *p*-isomers  
 d) *o*-nitrophenol is more volatile in steam than those of *m*- and *p*-isomers
577. Among the following most basic compound is:  
 a) Benzyl amine      b) Aniline      c) Acetanilide      d) *p*-nitro aniline
578. The EAN of platinum in potassium hexachloroplatinate (IV) is:  
 a) 46      b) 86      c) 36      d) 84
579. The number of ions formed when copper ammonium sulphate is dissolved in water is:  
 a) 1      b) 2      c) 4      d) Zero
580. Which of the following cannot show linkage isomerism?  
 a)  $NO_2^-$       b)  $NH_3$       c)  $CN^-$       d)  $SCN^-$
581. Xylenes on oxidation with acidic  $KMnO_4$  gives:  
 a) Phthalic acid      b) Isophthalic acid      c) Terephthalic acid      d) All of these
582. The ratio of  $\sigma$ - and  $\pi$ -bonds in benzene is:  
 a) 2      b) 4      c) 6      d) 8
583. The order of decreasing reactivity towards  $S_E$  reaction for the given compound is:

(i)  $C_6H_6$





(ii)  $C_6H_5CH_3$

(iii)  $C_6H_5Cl$

(iv)  $C_6H_5OH$

- a) (ii) > (iv) > (i) > (iii)      b) (iv) > (iii) > (ii) > (i)      c) (iv) > (ii) > (i) > (iii)      d) (i) > (ii) > (iii) > (iv)

584. Which of the following compounds is not optically active?




585. The number of geometrical isomers of  $[\text{Co}(\text{NH}_3)_3(\text{NO}_2)_3]$  are:  
 a) Zero                                      b) 2                                      c) 3                                      d) 4
586. Phenol is less acidic than:  
 a) Water                                      b) *p*-methoxyphenol                      c) *p*-nitrophenol                      d) Ethanol
587. In the reaction,  

$$\text{C}_6\text{H}_5\text{CH}_3 \xrightarrow{\text{Oxidation}} \text{A} \xrightarrow{\text{NaOH}} \text{B} \xrightarrow[\Delta]{\text{Soda lime}} \text{C}_\text{S}$$
  
 a)  $\text{C}_6\text{H}_5\text{OH}$                                       b)  $\text{C}_6\text{H}_6$                                       c)  $\text{C}_6\text{H}_5\text{COONa}$                       d)  $\text{C}_6\text{H}_5\text{ONa}$
588. Incorrect statement is  
 a) Ethane can have an infinite number of conformations  
 b) Cyclopropane molecule has considerable angle strain  
 c) Eclipsed form of ethane is less stable than staggered conformation  
 d) Staggered conformation possess maximum energy
589. The complex  $[\text{Co}(\text{NH}_3)_5\text{Br}]\text{SO}_4$  will give white ppt. with:  
 a)  $\text{PbCl}_2$                                       b)  $\text{AgNO}_3$                                       c)  $\text{KI}$                                       d) None of these
590. Which of the following complexes exhibits the highest paramagnetic behaviour?  
 a)  $[\text{Fe}(\text{en})(\text{bpy})(\text{NH}_3)_2]^{2+}$   
 b)  $[\text{Co}(\text{OX})_2(\text{OH})_2]^-$   
 c)  $[\text{Ti}(\text{NH}_3)_6]^{3+}$   
 $[\text{V}(\text{gly})_2(\text{OH})_2(\text{NH}_3)_2]^+$   
 d) Where, gly = glycine, en = ethylenediamine and bpy = bipyridylmoities  
 (At. No. Ti=22, V=23, Fe=26, Co=27)
591. The coordination number in a/an ..... complex may increase to 8.  
 a) Cobalt                                      b) Osmium                                      c) Nickel                                      d) Iron
592. Compound used for covering wounds caused by bite of mad dog is:  
 a) Benzoic acid                                      b) Aniline                                      c) Phenol                                      d) Salicylic acid
593. Cinnamic acid on decarboxylation gives:  
 a) Benzene                                      b) Toluene                                      c) Styrene                                      d) Benzaldehyde
594. In which of the following pairs both the complex show optical isomerism?  
 a) *Cis*- $[\text{Cr}(\text{C}_2\text{O}_4)_2\text{Cl}_2]^{3-}$ ; *cis*- $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]$                       b)  $[\text{PtCl}(\text{dien})]\text{Cl}$ ,  $[\text{NiCl}_2\text{Br}_2]^{2-}$   
 c)  $[\text{Co}(\text{NO}_3)_3(\text{NH}_3)_3]$ , *cis*- $[\text{Pt}(\text{en})_2\text{Cl}_2]$                       d)  $[\text{Co}(\text{en})_3]\text{Cl}_3$ , *cis*- $[\text{Co}(\text{en})_2\text{Cl}_2]\text{Cl}$
595. The name of the ring structure complex compound formed between metal ion and polydentate ligand is  
 a) Simple complex                      b) Chelate complex                      c) Polynuclear complex                      d) None of the above
596. IUPAC name of  

$$\begin{array}{c} \text{Cl}_2\text{CH} - \text{CH} - \text{CH} - \text{CCl}_3 \\ | \quad | \\ \text{C}_2\text{H}_5 \quad \text{C}_2\text{H}_5 \end{array}$$
 is  
 a) 1,1,1,4,4-pentachloro-2,3-diethyl-butane  
 b) 3-(dichloromethyl)-4-(trichloromethyl)-hexane  
 c) 3-(trichloromethyl)-4-(dichloromethyl)-hexane  
 d) 1,1,4,4,4-pentachloro-2,3-diethyl butane
597. Which statement is wrong with regard to acetaldehyde and benzaldehyde?  
 a) Both react with hydroxylamine to form oximes  
 b) Both react with HCN to form cyanohydrin  
 c) Both react with NaOH to form polymers  
 d) Both react with hydrazine to form hydrazones
598. The coordination number of Cu in complex  $[\text{Cu}(\text{H}_2\text{O})_4]^{2+}$  is  
 a) 4                                      b) 3                                      c) 2                                      d) 1
599. Which reaction sequence would be best to prepare 3-chloroaniline from benzene?  
 a) Chlorination, nitration, reduction  
 b) Nitration, chlorination, reduction

- c) Nitration, reduction, chlorination  
 d) Nitration, reduction, acetylation, chlorination, hydrolysis
600. The complexes  $(\text{Co}(\text{NH}_3)_6)[\text{Cr}(\text{C}_2\text{O}_4)_3]$  and  $[\text{Cr}(\text{NH}_3)_6][\text{Co}(\text{C}_2\text{O}_4)_3]$
- a) Geometrical isomerism  
 b) Ionization energy  
 c) Coordination isomerism  
 d) Linkage isomerism

601. The reaction,



is an example of:

- a) Substitution reaction  
 b) Addition reaction  
 c) Condensation reaction  
 d) Elimination reaction
602. Given the molecular formula of the hexa coordinated complexes is

- (A)  $\text{CoCl}_3 \cdot 6\text{NH}_3$   
 (B)  $\text{CoCl}_3 \cdot 5\text{NH}_3$   
 (C)  $\text{CoCl}_3 \cdot 4\text{NH}_3$

If the number of coordinated  $\text{NH}_3$  molecules in *A*, *B* and *C* respectively are 6, 5 and 4 the primary valency in (*A*), (*B*) and (*C*) are

- a) 6, 5, 4  
 b) 3, 2, 1  
 c) 0, 1, 2  
 d) 3, 3, 3
603.  $\text{C}_6\text{H}_{14}$  has two tertiary carbons. The IUPAC name is
- a) *n*-hexane  
 b) 2-methylpentane  
 c) 3-methylpentane  
 d) 2,3-dimethylbutane

604. The compound  $[\text{Co}(\text{NO}_2)(\text{NH}_3)_5]\text{Cl}_2$  and  $[\text{Co}(\text{ONO})(\text{NH}_3)_5]\text{Cl}_2$  are examples of:

- a) Geometrical isomers  
 b) Linkage isomers  
 c) Ligand isomers  
 d) Ionization isomers


605. Which is not a  $\pi$ -bonded complex?

- a) Zeise salt  
 b) Ferrocene  
 c) Dibenzene chromium  
 d) Tetraethyl lead


606. When phenol is treated with  $\text{PCl}_5$ , the yield of chlorobenzene is generally poor because of the formation of:

- a) Benzoyl chloride  
 b) *p*-chlorophenol  
 c) *o*-chlorophenol  
 d) Tertiary phosphate





607. Which will show tautomerism?



608. The IUPAC name of compound



- a) *N*-phenylaminoethanone  
 b) *N*-phenylethanamide  
 c) *N*-phenylmethanamide  
 d) *N*-phenylaminomethane
609. Which one of the following is most reactive towards electrophilic reagent?



610. Which of the following shows  $dsp^2$  hybridisation?



611. Which one of the following is not an explosive?

a) Trinitrolycerine

b) o-aminotoluene

c) Dynamite

d) TNT

612. Spin only magnetic moment of the compound  $Hg[Co(SCN)_4]$  is

a)  $\sqrt{3}$

b)  $\sqrt{15}$

c)  $\sqrt{24}$

d)  $\sqrt{8}$

613. When phenol is treated with  $NH_3$  and  $ZnCl_2$ , it changes to:


a) Aniline

b) Salicylic acid


c) Cyclohexanol

d) None of these

614. In which complex is the transition metal in zero oxidation state?



615. The species having tetrahedral shape is



616. An imperfect complex of a complex compound is 100% ionized; the compound is called:

a) Double salt

b) Complex salt



c) Acid salt

d) Normal salt

617. For which transition metal ions are low spin complexes impossible?



d) All are correct



Select the complexes which are diamagnetic.

a) (A), (B) and (C)

b) (B), (C) and (D)

c) (A), (C) and (D)

d) (A), (B) and (D)

619. Wilkinson's catalyst,  $(Ph_3P)_3RhCl$  is used for

a) Hydrogenation of oils

b) Hydrogenation of alkynes

c) Hydrogenation of alkenes

d) Polymerization of alkenes

620. Among the following compounds, the most acidic is:

a) p-nitrophenol

b) p-hydroxybenzoic acid

c) o-hydroxybenzoic acid

d) p-toluic acid

621. An aromatic primary amine with cold nitrous acid leads to the formation of:

a) Alcohol

b) Nitrite

c) Diazonium salt

d) Benzene

622. Chlorobenzene gives DDT when it reacts with:


a) Phenol

b) Naphthalene





c) Chloral

d) Acetaldehyde

623. Under suitable conditions  $C_6H_5CH_2OH$  (A),  $C_6H_5OH$  (B) and  $C_6H_5COOH$  (C) can act as acids. The increasing order of their acidic strengths is:



624. Which is considered to be an anticancer species?



625. The compound required for the formation of thermosetting polymer with methanal is:

a) Phenol





b) Benzene

c) Benzaldehyde

d) All of these

626. Which one of the following has highest number of isomers?

- a)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]^{2+}$       b)  $[\text{Co}(\text{en})_2\text{Cl}_2]^+$       c)  $[\text{Ru}(\text{NH}_3)_4\text{Cl}^-]$       d)  $[\text{In}(\text{PP}_3)_2\text{H}(\text{CO})]^{2+}$
627. Which group is *o*- and *p*-directing?  
 a)  $-\text{NO}_2$       b)  $-\text{SO}_3\text{H}$       c)  $-\text{COOH}$       d)  $-\text{NHCOCH}_3$
628. When benzyl chloride is boiled with aqueous solution of lead nitrate in current of carbon dioxide, the main product is:  
 a) Benzoic acid      b) Benzyl alcohol      c) Benzaldehyde      d) Nitrobenzene
629. Ligands in complex compounds  
 a) Donates electron pair      b) Accept electron pair  
 c) Neither accept electron pair nor donate      d) All of the above
630. Aniline is separated by:  
 a) Fractional crystallisation  
 b) Fractional distillation  
 c) Steam distillation  
 d) Vacuum distillation
631. In which of the following octahedral complexes of Co (at. No. 27), will be magnitude of  $\Delta_0$  be the highest?  
 a)  $[\text{Co}(\text{CN})_6]^{3-}$       b)  $[\text{Co}(\text{C}_2\text{O}_4)_3]^{3-}$       c)  $[\text{Co}(\text{H}_2\text{O})_6]^{3+}$       d)  $[\text{Co}(\text{NH}_3)_6]^{3+}$
632. The IUPAC name of  $\text{K}_2[\text{PtCl}_6]$  is  
 a) Hexachloroplatinate potassium      b) Potassium hexachloroplatinate (IV)  
 c) Potassium hexachloroplatinate      d) Potassium hexachloroplatinum(IV)
633. Aqueous solution of nickel sulphate on treating with pyridine and then adding a solution of sodium nitrite gives dark blue crystals of:  
 a)  $[\text{Ni}(\text{py})_4]\text{SO}_4$       b)  $[\text{Ni}(\text{py})_2(\text{NO}_2)_2]$       c)  $[\text{Ni}(\text{py})_4(\text{NO}_2)_2]$       d)  $[\text{Ni}(\text{py})_3(\text{NO}_2)]_2\text{SO}_4$
634. Benzyl alcohol is obtained from benzaldehyde by:  
 a) Fittig's reaction      b) Cannizzaro's reaction      c) Kolbe's reaction      d) Wurtz's reaction
635. The structure of the compound that gives a tribromo derivative on treatment with bromine water is:



636. The coordination number and the oxidation state of the element 'E' in the complex  $[\text{E}(\text{en})_2(\text{C}_2\text{O}_4)]\text{NO}_2$  (where (en) is ethylene diamine) are, respectively:  
 a) 6 and 3      b) 6 and 2      c) 4 and 2      d) 4 and 3
637. Benzaldehyde reacts with  $\text{PCl}_5$  to give:  
 a) Benzyl chloride      b) Benzo trichloride      c) Benzal chloride      d) Chlorobenzene
638. Which one of the following complex ions has geometrical isomers?  
 a)  $[\text{Co}(\text{en})_3]^{3+}$       b)  $[\text{Ni}(\text{NH}_3)_5\text{Br}]^+$       c)  $[\text{Co}(\text{NH}_3)_2(\text{en})_2]^{3+}$       d)  $[\text{Cr}(\text{NH}_3)_4(\text{en})]^{3+}$
639. The strongest acid among the following aromatic compounds is:  
 a) *Ortho*-nitrophenol      b) *para*-chlorophenol      c) *para*-nitrophenol      d) *meta*-nitrophenol
640. The isomers observed in alkanes is  
 a) Metamerism      b) Chain isomerism  
 c) Position isomerism      d) Geometrical isomerism
641. The two compounds pentaamminesulphatocobalt (III) bromide and pentaamminesulphatocobalt(III) chloride represent:  
 a) Linkage isomerism  
 b) Ionization isomerism  
 c) Coordination isomerism  
 d) No isomerism
642. Both  $[\text{Ni}(\text{CO})_4]$  and  $[\text{Ni}(\text{CN})_4]^{2-}$  are diamagnetic. The hybridisation of nickel in the compounds

respectively are :





a)  $sp^3, sp^3$

b)  $sp^3, dsp^2$

c)  $dsp^2, sp^3$

d)  $dsp^3, dsp^2$

643. The following compounds on hydrolysis in aqueous acetone will give:



a) Mixture of (K) and (L)    b) Mixture of (K) and (M)    c) Only (M)

d) Only (K)

644. The number of  $\pi$ -electrons in cyclo hepta trienyl anion is:

a) 2

b) 3

c) 8

d) 5

645. In the Grignard reaction, which metal forms an organometallic bond?

a) Sodium

b) Titanium

c) Magnesium

d) Palladium

646. Aromatic hydrocarbons are the derivatives of:

a) Benzene

b) Methane

c) Normal series of paraffins

d) None of the above

647. Benzene easily shows:


a) Ring fission reactions since it is unstable

b) Addition reactions since it is unsaturated

c) Electrophilic substitution reactions due to stable ring and high  $\pi$ -electron density

d) Nucleophilic substitution reactions due to stable ring and minimum electron density

648. The IUPAC name of the compound



is


a) Tetra phenyl methane

c) 1,1,1,1-tetracyclohexyl methane

b) 1,1,1,1-tetraphenyl methane

d) Methyno-1,1,1,1-tetracyclohexane

649.



having the IUPAC name as

a) 2,4,4-trimethyl pentanal

c) 1,3,3-trimethyl butanal

b) 4,4,2-trimethyl pentanal

d) 3,3,1-trimethyl butanal

650. When benzoic acid is heated with soda lime, we get:

a) Phenol

b) Benzyl alcohol

c) Benzene

d) Benzaldehyde

651. If a compound absorbs violet colour from the sunlight, then the observed colour is:

a) Yellow

b) Orange

c) Blue

d) Green

652. Sulphonic acid is used in the manufacture of:

a) Antipyretics

b) Antitoxine

c) Antibiotics

d) Dyes

653. In the silver plating of Cu,  $K[Ag(CN)_2]$  is used instead of  $AgNO_3$ . The reason is:

a) A thin layer of Ag is formed on Cu

b) More heat is required



sodium nitrite in the presence of dilute HCl. The compound so formed is treated with fluoroboric acid which is subsequently heated dry. The final product is:

- a) *p*-bromofluorobenzene
- b) *p*-bromoaniline
- c) 2,4,6-tribromofluorobenzene
- d) 1,3,5-tribromobenzene


669. Which of the following is a common donor atom in ligands?

- a) Nitrogen
- b) Oxygen
- c) Arsenic
- d) Both (b) and (c)

670. The reaction of aniline with acetyl chloride in presence of NaOH gives:

- a) Acetanilide
- b) Aniline hydrochloride
- c) *p*-chloroaniline
- d) A red dye

671. In the reaction, the compound "X" is:



- a)  $\text{CH}_3\text{COOH}$
- b)  $\text{Br} \cdot \text{CH}_2\text{COOH}$
- c)  $(\text{CH}_3\text{CO})_2\text{O}$
- d)  $\text{CHO} \cdot \text{COOH}$

672. Which of the following will exhibit maximum ionic conductivity?

- a)  $\text{K}_4[\text{Fe}(\text{CN})_6]$
- b)  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$
- c)  $[\text{Cu}(\text{NH}_3)_4]\text{Cl}_2$
- d)  $[\text{Ni}(\text{CO})_4]$

673. Dipole moment of *p*-nitroaniline, when compared to nitrobenzene (X) and aniline (Y) will be:

- a) Greater than (X) and (Y)
- b) Smaller than (X) and (Y)
- c) Greater than (X) but smaller than (Y)
- d) Equal to zero


674. The structure of iron pentacarbonyl is:

- a) Square planar
- b) Trigonal bipyramidal
- c) Triangular
- d) None of these

675. Turnbull's blue is:

- a) Ferricyanide
- b) Ferrous ferricyanide
- c) Ferrous cyanide
- d) Ferri ferrocyanide

676. The correct IUPAC name of



- a) 2-hydroxypropane-1, 2, 3-tricarboxylic acid
- b) 3-carboxy-3-hydroxy-pentane-1, 5-dioic acid
- c) 2 carboxy-4 hydroxy-pentane-1, 5-dioic acid
- d) 3-carboxy-3-hydroxy-hexane-1, 6-dioic acid

677. The trivial name among the following is

- a) Acetone
- b) Acetylene
- c) Uric acid
- d) None of these

678. The IUPAC name of  $[\text{Pt}(\text{NH}_3)_4(\text{NO}_2)\text{Cl}]\text{SO}_4$  is

- a) Chloronitro tetrammine platinum (IV) sulphate
- b) Tetrammine chloronitro platinum (II) sulphate
- c) Tetrammine chloronitro platinum (IV) sulphate
- d) Chlorotetrammine nitroplatinum (IV) sulphate

679. The overlapping in benzene is in carbon-carbon orbitals of the type:

- a)  $p - p$
- b)  $sp - sp$
- c)  $sp^2 - sp^2$
- d)  $sp^3 - sp^3$



680. Change in composition of coordination sphere yields which type of isomer?

- a) Geometrical
- b) Ionization
- c) Optical
- d) None of these

681. The IUPAC name of  $\text{K}_2[\text{Ni}(\text{CN})_4]$  is

- a) Potassium tetracyanonickelate (II)
- b) Potassium tetracyanonickelate (III)
- c) Potassium tetracyanonickel (II)
- d) Potassium tetracyanonickel (III)

682. Aniline in a set of the following reactions yielded a coloured compound Y:



683. The effective atomic number rule is less likely to apply if the metal-ligand bond:

- Is extremely weak
- Has a covalent character
- Has a large amount of ionic character
- None is correct

684. Potassium ferrocyanide is an example of

- Tetrahedral
- Octahedral
- Square planar
- Linear

685. 1-phenyl, 2-chloropropane on treatment with aqueous KOH gives mainly:

- 1-phenylpropane
- 3-phenylpropane
- 1-phenylpropan-2-ol
- 1-phenylpropan-3-ol


686. Which class of compounds can exhibit geometrical isomerism?

- $\text{C}_6\text{H}_5\text{CH} = \text{NOH}$
- $\text{CH}_3\text{CH} = \text{CHCH}_3$
- $\text{HOOCCH} - \text{CH}_2 - \text{CHCOOH}$
- All of the above

687. The product of oxidation of aniline with  $\text{K}_2\text{Cr}_2\text{O}_7$  and conc.  $\text{H}_2\text{SO}_4$  will be:

- p-amino phenol
- p-benzoquinone
- Aniline black dye
- Phenyl hydroxylamine

688. Among the following the Newmann projections of *meso*-2, 3-butanediol are







- P, Q
- P, R
- R, S
- Q, S

689. A new carbon-carbon bond is formed in:

- Cannizzaro's reaction
- Friedel-Crafts reaction

- c) Clemmensen reduction  
d) None of the above

690. Which of the following compounds can exhibit tautomerism?



691. The most basic compound among the following is:

- a) Benzylamine      b) Aniline      c) Acetanilide      d) *p*-nitroaniline

692. Which of the following has least oxidation state of Fe?

- a)  $\text{K}_3[\text{Fe}(\text{OH})_6]$       b)  $\text{K}_2[\text{FeO}_4]$   
c)  $\text{FeSO}_4(\text{NH}_4)_2\text{SO}_4 \cdot 6\text{H}_2\text{O}$       d)  $[\text{Fe}(\text{CN})_6]^{3-}$

693. The spin only magnetic moment value (in Bohr magneton units) of  $\text{Cr}(\text{CO})_6$  is

- a) 0      b) 2.84      c) 4.90      d) 5.92

694. Which is an excellent antiseptic?

- a) Phenol      b) Benzyl alcohol      c) Benzaldehyde      d) Acetic acid

695. Scientist who explained the structures and isomerism in the complex compound was:

- a) Sidgwick      b) Pauling      c) Powell      d) Werner

696. The cation that does not form an ammine complex with excess of ammonia is:

- a)  $\text{Al}^{3+}$       b)  $\text{Ag}^+$       c)  $\text{Cu}^{2+}$       d)  $\text{Cd}^{2+}$

697. The complex ion which has the highest magnetic moment among the following is

- a)  $[\text{CoF}_6]^{3-}$       b)  $[\text{Co}(\text{NH}_3)_6]^{3+}$       c)  $[\text{Ni}(\text{NH}_3)_4]^{2+}$       d)  $[\text{Ni}(\text{CN})_4]^{2-}$

698. For square planar complex of platinum (II),  $[\text{Pt}(\text{NH}_3)(\text{Br})(\text{Cl})\text{Py}]^{2+}$ , how many isomeric forms are possible?

- a) Two      b) Three      c) Four      d) Six

699. Which of the following has highest boiling point?

- a) Benzene      b) Phenol      c) Toluene      d) Ethyl benzene

700. A nitrogen containing organic compound on heating with chloroform and alcoholic KOH evolved very unpleasant smelling vapours. The compound could be:

- a) Nitrobenzene      b) Benzamide      c) *N,N*-dimethyl amine      d) Aniline


701. Which of the following 0.1 M complex compound solutions will have the minimum electrical conductivity?

- a) Hexammine platinum (IV) chloride      b) Chloropenta ammine platinum (IV) chloride  
c) Dichloro tetrammine platinum (IV) chloride      d) Trichloro triammine platinum (IV) chloride

702. False statement is

- a) Aprotic solvents increase the enol content in tautomerism  
b) Any deviation from the normal bond angles introduces angle strain in molecule  
c) Diastereomers have identical physical properties  
d) Chain isomers can also be position isomers

703. The correct IUPAC name of the compound is






- a) 4-bromo-5-chloro-3-iodo hexane      b) 3-bromo-2-chloro-4-iodo hexane  
c) 3-bromo-4-iodo -2-chloro hexane      d) 2-bromo-3-bromo-4-iodo hexane

704. Benzyl chloride ( $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$ ) can be prepared from toluene by chlorination with:


- a)  $\text{SO}_2\text{Cl}_2$       b)  $\text{SOCl}_2$       c)  $\text{S}_2\text{Cl}_2$       d)  $\text{NaOCl}$

705. The compound 2,2'-bipyridine has the structure





706. The IUPAC name of




- a) 4-formyl-6-oxocyclohexane-1-carboxylic acid  
c) 6-oxo-4-formyl cyclohexane-1-carboxylic acid

- b) 2-oxo-4-formyl cyclohexane-1-carboxylic acid  
d) 4-formyl-2-oxo cyclohexane-1-carboxylic acid

707.

The diazonium salt  $\text{H}_3\text{C}-\text{C}_6\text{H}_3(\text{Br})-\text{N}_2\text{Cl}$ , gives,



- a)  $\text{HCl}/\text{CuCl}$                       b)  $\text{HNO}_2/\text{Cu}$                       c)  $\text{C}_2\text{H}_5\text{OH}/\text{Cu}$                       d)  $\text{SnCl}_2/\text{HCl}$

708. Diethylenetriamine is:

- a) Chelating agent                      b) Polydentate ligand                      c) Tridentate ligand                      d) All of these

709. The no. of ions given by  $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}_2$  in aqueous solution is:

- a) 2                                              b) 3                                              c) 4                                              d) 5

710. Aniline reacts with excess of bromine to give:

- a) Benzyl bromide and hydrobromic acid  
b) 2,4,6-tribromoaniline  
c) 2-bromotoluene and hydrobromic acid  
d) 2-bromophenol and hydrobromic acid

711. The coordination compounds,

$[\text{Co}(\text{NH}_3)_6]^{3+}[\text{Cr}(\text{CN})_6]^{3-}$   
and  $[\text{Cr}(\text{NH}_3)_6]^{3+}[\text{Co}(\text{CN})_6]^{3-}$  are example of

- a) Linkage isomerism                      b) Coordination isomerism  
c) Ionisation isomerism                      d) Geometrical isomerism



712. Both  $\text{Co}^{3+}$  and  $\text{Pt}^{4+}$  have a coordination number of six. Which of the following pairs of complexes will show approximately the same electrical conductance for their 0.001 M. aqueous solutions?




- a)  $\text{CoCl}_3 \cdot 4\text{NH}_3$  and  $\text{PtCl}_4 \cdot 4\text{NH}_3$                       b)  $\text{CoCl}_3 \cdot 3\text{NH}_3$  and  $\text{PtCl}_4 \cdot 5\text{NH}_3$   
c)  $\text{CoCl}_3 \cdot 6\text{NH}_3$  and  $\text{PtCl}_4 \cdot 5\text{NH}_3$                       d)  $\text{CoCl}_3 \cdot 6\text{NH}_3$  and  $\text{PtCl}_4 \cdot 3\text{NH}_3$

713. In SCN ligand if N is attached to central atom, the name of ligand is:

- a) Thiocyanato-N                      b) Cyanato-N                      c) Thiocyanato-S                      d) Cyanato-S

714. The product formed on heating









715. Oxidation of ethyl benzene by  $\text{KMnO}_4$  gives:

- a) Benzyl alcohol      b) Benzophenone      c) Acetophenone      d) Benzoic acid

716. One of the following statements regarding Reimer-Tiemann reaction is false:

- a) Reaction of phenol with  $\text{CHCl}_3$  and  $\text{KOH}$   
 b)  $\text{CCl}_2$  acts as a nucleophile  
 c) Reaction of phenol with  $\text{CCl}_4$  and  $\text{NaOH}$   
 d) Reaction of phenol with formaldehyde to form bakelite

717. The structure representing a heterocyclic compound is



718. Phenol reacts with  $\text{Br}_2$  in  $\text{CCl}_4$  at low temperature to give:

- a) *m*-bromophenol  
 b) *o*- and *p*-bromophenol  
 c) *p*-bromophenol  
 d) 2,4,6-tribromophenol


719. The correct name of the compound  $[\text{Cu}(\text{NH}_3)_4](\text{NO}_3)_2$ , according to IUPAC system is

- a) Cuprammonium nitrate      b) Tetrammine copper (II) dinitrate  
 c) Tetrammine copper (II) nitrate      d) Tetrammine copper (II) dinitrite

720. Nitroethane can exhibit one of the following kind of isomerism

- a) Metamerism      b) Optical activity      c) Tautomerism      d) Position isomerism

721. What would be the correct IUPAC name of




- a) 3,3-dimethyl-3-cyclopentyl propanal  
 b) 3-methyl-3-cyclopentyl butan-1-al  
 c) 1-(1-methyl-1-formyl) methylethyl cyclopropane  
 d) None of above

722. The number of unpaired electrons in the square planar  $[\text{Pt}(\text{CN})_4]^{2-}$  ion is

- a) 2      b) 1      c) 0      d) 3

723. The oxidation number of cobalt in  $\text{K}[\text{Co}(\text{CO})_4]$  is

- a) +1                                      b) +3                                      c) -1                                      d) -3
724. IUPAC name of  $\text{Na}_3[\text{Co}(\text{NO}_2)_6]$  is  
 a) Sodium hexanitrito cobaltate (II)                                      b) Sodium hexanitro cobaltate (III)  
 c) Sodium hexanitrito cobaltate (III)                                      d) Sodium cobaltinitrite(II)
725. The total number of possible isomers for the complex compound  $[\text{Cu}(\text{NH}_3)_4][\text{PtCl}_4]$   
 a) 6                                      b) 5                                      c) 4                                      d) 3
726. Benzaldehyde reacts with excess of anhydrous ethyl alcohol in the presence of HCl, gives:  
 a)  $\text{C}_6\text{H}_5\text{COCl}$                                       b)  $\text{C}_6\text{H}_5\text{COOC}_2\text{H}_5$                                       c)  $\text{C}_6\text{H}_5\text{CH}(\text{OC}_2\text{H}_5)_2$                                       d)  $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$
727. Which pair of isomerism is not possible together?  
 a) Chain and position                                      b) Functional and position  
 c) Tautomerism and functional                                      d) All of the above
728. Which type of conformation is shown by I and II?



- a) I is eclipsed, II is staggered                                      b) II is eclipsed, I is staggered  
 c) Both are eclipsed                                      d) Both are staggered
729. Which will give chiral molecule?  
 a)  $\text{CH}_3\text{COCl} \xrightarrow{\text{LiAlH}_4}$                                       b)  $\text{C}_2\text{H}_5\text{CHO} \xrightarrow[\text{H}^+/\text{H}_2\text{O}]{\text{CH}_3\text{MgBr}}$   
 c)  $(\text{CH}_3)_2\text{CHC}_2\text{H}_5 \xrightarrow{\text{Cu}}$                                       d)  $\text{H}_3\text{C}-\text{C}(\text{H})=\text{C}(\text{CH}_3)_2 \xrightarrow{\text{Cl}_2}$
730. The neutral ligand is:  
 a) Chloro                                      b) Hydroxo                                      c) Ammine                                      d) Oxalato
731. The effective atomic number of cobalt in the complex  $[\text{Co}(\text{NH}_3)_6]^{3+}$  is  
 a) 36                                      b) 24                                      c) 33                                      d) 30
732.  $\text{K}_4[\text{Fe}(\text{CN})_6]$  is used to detect the presence of:  
 a) Metallic ion                                      b) Ferrous ion                                      c) Ferric ion                                      d) None of these
733. *p*-nitro benzaldehyde reacts with concentrated NaOH solution at room temperature to give:  
 a) *p*-nitrobenzamide  
 b) *p*-nitro benzyl alcohol and sod. *p*-nitrobenzoate  
 c) Benzaldehyde  
 d) *p*-nitrotoluene
734. But-1-ene and cyclobutane exhibit  
 a) Ring chain isomerism  
 b) Position isomerism  
 c) Tautomerism  
 d) Functional isomerism
735. The groups satisfying the secondary valencies of a cation in a complex are called:  
 a) Ligands                                      b) Radicals                                      c) Primary valencies                                      d) None of these
736. Benzene was discovered by:  
 a) Cavendish                                      b) Faraday                                      c) Berzelius                                      d) Wöhler
737. The number of structural and configurational isomers of a bromo compound  $\text{C}_5\text{H}_9\text{Br}$  obtained by the addition of HBr on 2-pentyne respectively are  
 a) 1, 2                                      b) 2, 4                                      c) 4, 2                                      d) 2, 1

738. The primary valency of Fe in  $K_3[Fe(CN)_6]$  is:

- a) 3                                      b) 2                                      c) 1                                      d) Zero


739. Which complex compound obeys 18-electron rule?

- a)  $[V(CO)_5]$                               b)  $[Fe(NH_3)_6]^{2+}$                               c)  $[Ni(CO)_6]$                               d)  $[Mn(H_2O)_6]^{2+}$

740. Two isomers  $X$  and  $Y$  with the formula  $Cr(H_2O)_5ClBr_2$  were taken for experiment on depression in freezing point. It was found that one mole of  $X$  gave depression corresponding to 2 moles of particles and one mole of  $Y$  gave depression due to 3 moles of particles. The structural formula of  $X$  and  $Y$  respectively, are


- a)  $[Cr(H_2O)_5Cl]Br_2$ ;  $[Cr(H_2O)_4Br_2]Cl \cdot H_2O$                               b)  $[Cr(H_2O)_5Cl]Br_2$ ;  $[Cr(H_2O)_3ClBr_2] \cdot 2H_2O$   
 c)  $[Cr(H_2O)_5Br]BrCl$ ;  $[Cr(H_2O)_4ClBr]Br \cdot H_2O$                               d)  $[Cr(H_2O)_4Br_2]Cl \cdot H_2O$ ;  $[Cr(H_2O)_5Cl]Br_2$

741. The IUPAC name of



- a) 1-formyl-3-oxo-pentanoic acid                              b) 5-formyl-3-oxo pentanoic acid  
 c) 3-oxo-5-formyl pentanoic acid                              d) 3-oxo-1-formyl pentanoic acid

742. The two complexes given below are:



- a) Geometrical isomers                              b) Position isomers                              c) Optical isomers                              d) Identical


743. Which of the following statements is not correct?

- a) In oxyhaemoglobin  $Fe^{2+}$  is paramagnetic  
 b) During respiration the size of  $Fe^{2+}$  increases when it changes from diamagnetic to paramagnetic state  
 c) Four haeme groups are present in haemoglobin  
 d) Haeme is the prosthetic group and it is non-protein part

744. Chlorination of toluene in the presence of light and heat followed by treatment with aqueous NaOH gives:

- a) *o*-cresol                              b) *p*-cresol                              c) 2,4-dihydroxytoluene                              d) Benzoic acid


745. Which of the following has maximum probability of showing tautomerism?






746. The halide which undergoes nucleophilic substitution most readily is:

- a) *p*- $H_3CC_6H_4Cl$                               b) *o*- $H_3COC_6H_4Cl$                               c) *p*- $ClC_6H_4Cl$                               d) *p*- $O_2NC_6H_4Cl$

747. The major product (70% to 80%) of the reaction between *m*-dinitrobenzene with  $(NH_4)_2S_x$  is:



748. The '*E*'-isomer is



d) None of the above

749. The Baeyer angle strain is minimum in

- a) Cyclopropane                      b) Cyclobutane                      c) Cyclopentane                      d) Cyclohexane


750. Among the following ions, which one has the highest unpaired electrons?

- a)  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$                       b)  $[\text{Zn}(\text{H}_2\text{O})_6]^{2+}$                       c)  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$                       d)  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$

751. Which will give a white precipitate with  $\text{AgNO}_3$  in aqueous solution?

- a)  $[\text{Co}(\text{NH}_3)_5\text{Cl}](\text{NO}_2)_2$                       b)  $[\text{Pt}(\text{NH}_3)_6]\text{Cl}_4$                       c)  $[\text{Pt}(\text{en})\text{Cl}_2]$                       d)  $[\text{Cu}(\text{NH}_3)_4]\text{SO}_4$

752. The organic product formed in the reaction;



- a)  $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$  and  $\text{CH}_3\text{OH}$   
 b)  $\text{C}_6\text{H}_5\text{COOH}$  and  $\text{CH}_4$   
 c)  $\text{C}_6\text{H}_5\text{CH}_3$  and  $\text{CH}_3\text{OH}$   
 d)  $\text{C}_6\text{H}_5\text{CH}_3$  and  $\text{CH}_4$


753. Complexes with bidentate ligands are called:

- a) Ligands                      b) Chelates                      c) Complexes                      d) None of these

754. Excited state configuration of  $\text{Mn}^{2+}$  is

- a)  $t_{2g}^4$                       b)  $t_{2g}^3 e_g^2$                       c)  $t_{2g}^4 e_g^2$                       d)  $t_{2g}^5 e_g^0$

755. The IUPAC name of



- a) Ethyl acetylate                      b) Ethyl methyl butenoate  
 c) Ethyl acetoethanoate                      d) Ethyl (3-methyl) but-2-enoate

756. The compound which result from the coordination of carbon monoxide are known as

- a) Carbon permono                      b) Electronic                      c) Carbonyls                      d) None of these

757. The correct IUPAC name of  $\text{AlCl}_3(\text{EtOH})_4$  is:

- a) Aluminium(II) chloride-4-ethanol  
 b) Aluminium(III)chloride-4-ethanol  
 c) Aluminium(IV)chloride-4-hydroxy ethane  
 d) Aluminium chloride-4-ethanol

758. The IUPAC name of  $[\text{Co}(\text{NH}_3)_6][\text{Cr}(\text{C}_2\text{O}_4)_3]$  is

- a) Hexaamine cobalt (III) tris (oxalato ) chromium  
 b) Hexaamine cobalt (III) tris (oxalato ) chromate(III)  
 c) Hexaamine cobalt tris (oxalato ) chromium(III)  
 d) Hexaamine cobalt (III) chromium (III) oxalate

759. The insecticide, germicide gammexane is a formulation for:

- a) DDT  
 b) Benzene hexachloride  
 c) Hexachlorobenzene  
 d) Chloral

760. Among  $[\text{Ni}(\text{CO})_4]^{2-}$ ,  $[\text{Ni}(\text{CN})_4]^{2-}$ ,  $[\text{NiCl}_4]^{2-}$  species, the hybridisation states of the Ni atom are, respectively (Atomic no. of Ni=28)



- a)  $sp^3, dsp^2, dsp^2$                       b)  $sp^3, dsp^2, sp^3$                       c)  $sp^3, sp^3, dsp^2$                       d)  $dsp^2, sp^3, sp$

761. Which of the following complex ions is expected to absorb visible light?



d) Electron deficient molecule

776. The two isomers given below are



a) Enantiomers                      b) Diastereomers                      c) Measomers                      d) Position isomers

777. Which of the following has lowest boiling point?

a) Phenol                      b) *o*-nitrophenol                      c) *m*-nitrophenol                      d) *p*-nitrophenol

778. The IUPAC name of  $[\text{Ni}(\text{NH}_3)_4][\text{NiCl}_4]$  is

a) Tetrachloro nickel (II) - tetraammine nickel (II)  
b) Tetraammine nickel (II) - tetrachloro nickel (II)  
c) Tetraammine nickel (II) - tetrachloro nickelate (II)  
d) Tetrachloro nickel (II) - tetraammine nickelate (0)

779. All ligands are:

a) Lewis acid                      b) Lewis base                      c) Neutral                      d) None of these


780. Aspirin is known as:

a) Phenyl salicylate                      b) Acetyl salicylate                      c) Methyl salicylic acid                      d) Acetyl salicylic acid

781. Which of the following has on optical isomer?

a)  $[\text{Co}(\text{NH}_3)_3\text{Cl}]^+$                       b)  $[\text{Co}(\text{en})(\text{NH}_3)_2]^{2+}$                       c)  $[\text{Co}(\text{H}_2\text{O})_4(\text{en})]^{3+}$                       d)  $[\text{Co}(\text{en})_2(\text{NH}_3)_2]^{3+}$

782. The IUPAC name of the compound




a) 1,1,1-trichloro-2,2-diphenyl ethane                      b) 2,4,5-trichloro hexanol  
c) 2,2,2-trichloro bicyclo [4.4.0] nenone                      d) 2,2,2-trichloro-1,1-diphenyl ethane

783. The property by virtue of which a compound can rotate the plane of polarised light is known as

a) Polarisability                      b) Phosphorescence                      c) Optical activity                      d) Polarization

784. The molecules represented by the following two structures are







a) Epimers                      b) Diastereomers                      c) Enantiomers                      d) Identical

785. The IUPAC name of the coordination compound  $\text{K}_3[\text{Fe}(\text{CN})_6]$  is

a) Tripotassium hexacyanoiron (II)                      b) Potassium hexacyanoiron (II)  
c) Potassium hexacyanoferrate (III)                      d) Potassium hexacyanoferrate (II)

786. Which one of the following is an inner orbital complex as well as diamagnetic in nature?

- a)  $[\text{Cr}(\text{NH}_3)_6]^{3+}$       b)  $[\text{Co}(\text{NH}_3)_6]^{3+}$       c)  $[\text{Ni}(\text{NH}_3)_6]^{2+}$       d)  $[\text{Zn}(\text{NH}_3)_6]^{2+}$
787. How many unpaired electrons are present in the central metal ion of  $[\text{CoCl}_4]^{2-}$ ?
- a) 3      b) 4      c) 5      d) 2
788. Show the coordination number of the metal ion, its oxidation number, the number of electrons in  $d$ -orbitals and the number of unpaired electrons  $d$ -orbitals respectively in complex  $[\text{Co}(\text{H}_2\text{O})_4\text{SO}_3]\text{Cl}$ .
- a) 6, 3, 6, 4      b) 6, 3, 6, 0      c) 5, 3, 6, 4      d) 5, 3, 6, 0
789. Benzene reacts with.....to give acetophenone.
- a) Acetyl chloride  
b) Acetyl chloride in presence of anhy.  $\text{AlCl}_3$   
c) Anhy.  $\text{AlCl}_3$   
d) None of the above
790. Which group would you introduce into a drug or a dye to make it water soluble?
- a)  $-\text{NO}_2$       b)  $-\text{Cl}$       c)  $-\text{SO}_3\text{H}$       d)  $-\text{OH}$
791. In the coordination compound,  $\text{K}_4[\text{Ni}(\text{CN})_4]$ , oxidation state of nickel is
- a)  $-1$       b)  $+1$       c)  $0$       d)  $+2$
792. The IUPAC name of  $[\text{Cr}(\text{NH}_3)_4\text{Cl}_2]\text{NO}_3$  is:
- a) Tetraaminodichlorochromium (I) nitrate  
b) Tetraaminodichlorochromium (III) nitrate  
c) Dichlorotetraamminechromium (III) nitrate  
d) Tetraaminodichlorochromium (II) nitrate
793. Vanillin, used as a flavouring agent is:
- a) An aliphatic alcohol      b) An aromatic aldehyde      c) A hydrocarbon      d) A carbohydrate
794. Which of the following will exhibit optical isomerism?
- a)  $[\text{Cr}(\text{en})(\text{H}_2\text{O})_4]^{3+}$       b)  $[\text{Cr}(\text{en})_3]^{3+}$       c)  $\text{trans}-[\text{Cr}(\text{en})_2\text{Cl}_2]^+$       d)  $[\text{Cr}(\text{NH}_3)_6]^{3+}$
795. Which one is a mixed ketone?
- a) Benzophenone      b) Benzenone      c) Acetophenone      d) Dibenzyl ketone
796. Transition metals can form complexes in:
- a) Zero oxidation state      b) Cation form      c) Anion form      d) All of these
797. Toluene on oxidation with air in presence of  $\text{V}_2\text{O}_5$  yields:
- a) Phenol      b) Benzoic acid      c) Benzaldehyde      d) Benzyl alcohol
798.  $[\text{Pt}(\text{NH}_3)_4]\text{Cl}_2$  is
- a) Pyramidal      b) Pentagonal      c) Tetrahedral      d) Square planar
799. In  $\text{Fe}(\text{CO})_5$ , the  $\text{Fe}-\text{C}$  bond possess:
- a)  $\pi$ -character only  
b) Both  $\sigma$  and  $\pi$ -characters  
c) Ionic character  
d)  $\sigma$ -character only
800. Which molecule has tetrahedral geometry?
- a)  $[\text{Co}(\text{NH}_3)_6]^{3+}$       b)  $[\text{Ni}(\text{CN})_4]^{2-}$       c)  $\text{Fe}(\text{CO})_5$       d)  $[\text{NiCl}_4]^{2-}$
801.  $[\text{Co}(\text{NH}_3)_5\text{Br}]\text{SO}_4$  and  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Br}$  are the examples of:
- a) Linkage isomerism  
b) Geometrical isomerism  
c) Ionization isomerism  
d) Optical isomerism
802. The compounds  $R-\text{NO}_2$  and  $R-\text{ONO}$  are
- a) Geometrical isomers      b) Functional isomers  
c) Metamers      d) Optical isomers
803. Which of the following Fischer projection formula is same as D-glyceraldehyde?




804.  $[\text{Fe}(\text{NO}_2)_3\text{Cl}_3]$  and  $[\text{Fe}(\text{O}-\text{NO})_3\text{Cl}_3]$  shows

- a) Linkage isomerism  
c) Optical isomerism

- b) Geometrical isomerism  
d) None of the above

805.



The IUPAC name of the compound is

- a) 2-ethenyl-3-methyl cyclohexa-1, 3-diene  
c) 2, 6-dimethyl hepta-2, 5dienoic acid

- b) 2, 5-dimethyl hepta-2, 6-dienoic acid  
d) 2, 3-dimethyl epoxyethane

806. When benzene sulphonic acid and *p*-nitrophenol are treated with  $\text{NaHCO}_3$ , the gases released respectively are:

- a)  $\text{SO}_2, \text{NO}_2$                       b)  $\text{SO}_2, \text{NO}$                       c)  $\text{SO}_2, \text{CO}_2$                       d)  $\text{NO}_2, \text{CO}_2$

807. Which of the following is non-ionizable?

- a)  $[\text{Co}(\text{NH}_3)_3\text{Cl}_3]$                       b)  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$                       c)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$                       d)  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_2$





808. Increasing order of expected keto content

- a)  $\text{CH}_3\text{COC}_2\text{H}_5 > \text{CH}_3\text{CHO} > \text{CH}_3\text{COCH}_3 > \text{CH}_3\text{COCH}_2\text{COCH}_3$   
b)  $\text{CH}_3\text{COCH}_3 > \text{CH}_3\text{CHO} > \text{CH}_3\text{COC}_2\text{H}_5 > \text{CH}_3\text{COCH}_2\text{COCH}_3$   
c)  $\text{CH}_3\text{CHO} > \text{CH}_3\text{COC}_2\text{H}_5 > \text{CH}_3\text{COCH}_3 > \text{CH}_3\text{COCH}_2\text{COCH}_3$   
d)  $\text{CH}_3\text{COCH}_2\text{COCH}_3 > \text{CH}_3\text{CHO} > \text{CH}_3\text{COCH}_3 > \text{CH}_3\text{COC}_2\text{H}_5$


809. Which is colourless complex?

- a)  $\text{Cu}_2(\text{CH}_3\text{COO})_4 \cdot \text{H}_2\text{O}$   
b)  $\text{Cu}_2\text{Cl}_2$   
c)  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$   
d)  $[\text{Cu}(\text{NH}_3)_4]\text{SO}_4 \cdot \text{SO}_4 \cdot 4\text{H}_2\text{O}$

810. Which is not a reasonable structure for dimethyl benzene?



811.



The IUPAC name of the compound is

- a) Propionic anhydride                      b) Dipropanoic anhydride  
c) Ethoxy propanoic acid                      d) Propanoic anhydride


812. A mixture of benzene and aniline can be separated by:

- a) Alcohol                      b) Dil. HCl                      c) Dil. NaOH                      d) Hot water

813. The correct IUPAC name of the complex  $\text{Fe}(\text{C}_5\text{H}_5)_2$  is

- a) Cyclopentadienyl iron (II)                      b) Bis (Cyclopentadienyl)iron (II)  
c) Dicyclo pentadienyl ferrate (II)                      d) Ferrocane

814.



The electrophile involved in the above reaction is:

- a) dichloromethyl cation ( $\text{CHCl}_2^{\oplus}$ )  
 b) Dichlorocarbene ( $:\text{CCl}_2$ )  
 c) Trichloromethyl anion ( $\text{CCl}_3^{\ominus}$ )  
 d) Formyl cation ( $\text{CHO}^{\oplus}$ )
815. Benzoyl Chloride is prepared from benzoic acid by:  
 a)  $\text{Cl}_2, h\nu$                       b)  $\text{SO}_2\text{Cl}_2$                       c)  $\text{SOCl}_2$                       d)  $\text{Cl}_2, \text{H}_2\text{O}$
816. Which of the following ions forms most stable complex compound?  
 a)  $\text{Fe}^{3+}$                       b)  $\text{Mn}^{2+}$                       c)  $\text{Ni}^{2+}$                       d)  $\text{Cu}^{2+}$
817. Which one of the following cyano complexes would exhibit the lowest value of paramagnetic behaviour? (Atomic no. Cr=24, Mn=25, Fe=26, Co=27)  
 a)  $[\text{Co}(\text{CN})_6]^{3-}$                       b)  $[\text{Fe}(\text{CN})_6]^{3-}$                       c)  $[\text{Mn}(\text{CN})_6]^{3-}$                       d)  $[\text{Cr}(\text{CN})_6]^{3-}$
818. Which of the following statements is not correct?  
 a) The complexes  $[\text{NiCl}_4]^{2-}$  and  $[\text{Ni}(\text{CN})_4]^{2-}$  differ in the state of hybridisation of nickel.  
 b) The complexes  $[\text{NiCl}_4]^{2-}$  and  $[\text{Ni}(\text{CN})_4]^{2-}$  differ in the magnetic properties.  
 c) The complexes  $[\text{NiCl}_4]^{2-}$  and  $[\text{Ni}(\text{CN})_4]^{2-}$  differ in geometry.  
 d) The complexes  $[\text{NiCl}_4]^{2-}$  and  $[\text{Ni}(\text{CN})_4]^{2-}$  differ in primary valencies of nickel.
819. In the complexes  $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$ ,  $[\text{Fe}(\text{CN})_6]^{3-}$ ,  $[\text{Fe}(\text{C}_2\text{O}_4)_3]^{3-}$  and  $[\text{FeCl}_6]^{3-}$ , more stability is shown by:  
 a)  $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$                       b)  $[\text{Fe}(\text{CN})_6]^{3-}$                       c)  $[\text{Fe}(\text{C}_2\text{O}_4)_3]^{3-}$                       d)  $[\text{FeCl}_6]^{3-}$
820. In the reaction,
- $$\text{C}_8\text{H}_6\text{O}_4 \xrightarrow{\Delta} \text{X} \xrightarrow{\text{NH}_3} \text{C}_6\text{H}_4(\text{CONH}_2)(\text{COOH})$$
- the intermediate 'X' is:  
 a) Phthalic anhydride                      b) Phthalic acid                      c) *o*-xylene                      d) Benzoic acid
821. Which of the following is  $\pi$  complex?  
 a) Trimethyl aluminium                      b) Ferrocene                      c) Diethyl zinc                      d) Nickel carbonyl
822. When phenol is reacted with chloroform and an alkali like NaOH, the compound formed is salicylaldehyde. If we use pyrene in place of chloroform the product obtained is:  
 a) Salicylaldehyde                      b) Phenolphthalein                      c) Salicylic acid                      d) Cyclohexanol
823. Among the properties (a) reducing (b) oxidizing (c) complexing, the set of properties shown by  $\text{CN}^-$  ion towards metal species is  
 a) B, c                      b) A, b, c                      c) C, a                      d) A, b
824. Which of the following is most powerful *meta* directing group?  
 a)  $-\text{NO}_2$                       b)  $-\text{SO}_3\text{H}$                       c)  $-\text{CHO}$                       d)  $-\text{COOH}$
825. Which among the following compounds will show metamerism?  
 a)  $\text{CH}_3\text{COC}_3\text{H}_7$                       b)  $\text{CH}_3\text{OC}_2\text{H}_5$                       c)  $\text{CH}_3\text{SC}_2\text{H}_5$                       d)  $\text{CH}_3\text{OCH}_3$
826. The hybridization of  $[\text{PtCl}_6]^{2-}$  ion is:  
 a)  $d^2sp^3$                       b)  $sp^2d^3$                       c)  $sp^3d$                       d)  $sp^3d^2$
827. The correct name of  $[\text{Pt}(\text{NH}_3)_4\text{Cl}_2][\text{PtCl}_4]$  is  
 a) Tetrachloro platinum (II) dichloro tetrammine platinate  
 b) Dichloro tetrammine platinum (IV) tetrachloro platinate(II)  
 c) Tetrammine dichloro platinum (IV) tetrachloro platinate (II)  
 d) Tetrachloro platinum (II) tetrammine platinate (IV)
828. The oxidation state of iron in  $\text{K}_4[\text{Fe}(\text{CN})_6]$  is  
 a) 1                      b) 4                      c) 3                      d) 2
829. Formation of complex compound can be detected by:  
 a) Change in colour                      b) Change in solubility                      c) Change in pH                      d) All are correct
830. The complex that violates the EAN:



844. From the equation,  $3\text{C}_2\text{H}_2 \rightarrow \text{C}_6\text{H}_6$ , find the volume of acetylene (NTP) for the manufacture of 3 mole of benzene:
- a) 67.2 litre                      b) 134.4 litre                      c) 201.6 litre                      d) 33.8 litre
845. According to IUPAC nomenclature sodium nitroprusside is named as
- a) Sodium pentacyanonitrosyl ferrate(II)                      b) Sodium pentacyanonitrosyl ferrate(III)  
c) Sodium nitroferrocyanide                      d) Sodium nitroferrocyanide
846. Among  $[\text{Ni}(\text{CO})_4]$ ,  $[\text{Ni}(\text{CN})_4]^{2-}$  and  $[\text{NiCl}_4]^{2-}$  species the hybridisation states of Ni atom are respectively:
- a)  $sp^3, dsp^2, dsp^2$                       b)  $sp^3, dsp^2, sp^3$                       c)  $sp^3, sp^3, dsp^2$                       d)  $dsp^2, sp^3, sp^3$
847. The chemical name of DDT is:
- a) Dichloro dinitro toluene  
b) Dichloro dimethyl toluene  
c) *p, p'*-dichloro diphenyl trichloroethane  
d) None of the above
848. The stability of complexes of  $\text{Cu}^{2+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Co}^{2+}$  and  $\text{Fe}^{2+}$  varies in the order
- a)  $\text{Cu}^{2+} > \text{Ni}^{2+} > \text{Co}^{2+} > \text{Fe}^{2+}$                       b)  $\text{Cu}^{2+} > \text{Fe}^{2+} > \text{Ni}^{2+} > \text{Co}^{2+}$   
c)  $\text{Ni}^{2+} > \text{Co}^{2+} > \text{Fe}^{2+} > \text{Cu}^{2+}$                       d)  $\text{Cu}^{2+} < \text{Ni}^{2+} < \text{Co}^{2+} < \text{Fe}^{2+}$
849. The number of unpaired electrons in  $\text{Ni}(\text{CO})_4$  is
- a) 0                      b) 1                      c) 3                      d) 4
850. In sodium tetrafluorooxochromate(...),  $\text{Na}_3[\text{Cr}(\text{O})\text{F}_4]$  the left out place should be filled with which of the following roman numerals?
- a) VI                      b) III                      c) IV                      d) None of these
851. The IUPAC name of compound  
 $\text{CN} - \text{CH}_2 - \underset{\substack{| \\ \text{OCH}_3}}{\text{CH}} - \text{CH}_2 - \text{COOCH}_3$  is
- a) 3-methoxy-4-cyano methyl butanoate                      b) Methyl-4-cyano-3-methoxy butanoate  
c) 4-cyano-3-methoxy methyl butanoate                      d) Methyl-3-methoxy-4-cyano butanoate
852. Cumene is:
- a) *o*-methyl phenol                      b) *p*-cresol                      c) Isopropyl benzene                      d) Phenyl *n*-propane
853. In Etard's reaction toluene is oxidised to benzaldehyde using:
- a)  $\text{H}_2\text{O}_2$   
b)  $\text{Cl}_2$   
c) Chromium trioxide or  $\text{CrO}_2\text{Cl}_2$   
d)  $\text{KMnO}_4$
854. Which of the following will exhibit geometrical isomerism?
- a) Propene                      b) Butene-2  
c) Butene-1                      d) 1, 1-dichloro butane
855. Ferrocene is:
- a)  $\text{Fe}(\eta^5 - \text{C}_5\text{H}_5)_2$                       b)  $\text{Fe}(\eta^2 - \text{C}_5\text{H}_5)_2$                       c)  $\text{Cr}(\eta^5 - \text{C}_5\text{H}_5)_5$                       d)  $\text{Os}(\eta^5 - \text{C}_5\text{H}_5)_2$
856. Which one is an outer orbital complex?
- a)  $[\text{Ni}(\text{NH}_3)_6]^{2+}$                       b)  $[\text{Mn}(\text{CN})_6]^{4-}$                       c)  $[\text{Co}(\text{NH}_3)_6]^{3+}$                       d)  $[\text{Fe}(\text{CN})_6]^{4-}$
857. The pair of  $[\text{Co}(\text{SO}_4)(\text{NH}_3)_5]\text{Cl}$  and  $[\text{CoCl}(\text{NH}_3)_5]\text{SO}_4$  constitutes
- a) Optical isomers                      b) Linkage isomers                      c) Coordination isomers                      d) Ionisation isomers
858. The IUPAC name of  $\text{K}_2[\text{Cr}(\text{CN})_2\text{O}_2(\text{O})_2(\text{NH}_3)]$  is
- a) Potassiumammine dicyanodioxoperoxochromate (VI)  
b) Potassiumammine cyanoperoxodioxochrometic (IV)  
c) Potassiumammine dicyanodioxoperoxochromium (IV)  
d) Potassiumammine dicyanodioxoperoxochromium (IV)
859. In spectrochemical series chlorine is above than water *i.e.*,  $\text{Cl} > \text{H}_2\text{O}$ , this is due to
- a) Good  $\pi$ -acceptor properties of Cl

- b) Strong  $\sigma$  –donor and good  $\pi$ -acceptor properties of Cl  
 c) Good  $\pi$  –donor properties of Cl  
 d) Larger size of Cl than H<sub>2</sub>O
860. The type of isomerism shown by [Co(en)<sub>2</sub>(NCS)<sub>2</sub>]Cl and [Co(en)<sub>2</sub>(NCS)Cl]NCS is:  
 a) Coordination                      b) Ionization                      c) Linkage                      d) All of these
861. Which ion shows only the coordination number 4 in complexes?  
 a) Pt<sup>2+</sup>                      b) Cr<sup>3+</sup>                      c) Fe<sup>3+</sup>                      d) Pt<sup>4+</sup>
862. The spin magnetic moment of cobalt in Hg[Co(SCN)<sub>4</sub>] is :  
 a)  $\sqrt{3}$                       b)  $\sqrt{8}$                       c)  $\sqrt{15}$                       d)  $\sqrt{24}$
863. Which of the following is not an isomer of but-1-yne?  
 a) But-2-yne  
 b) Buta-1-3-diene  
 c) Methyl cyclopropene  
 d) But-2-ene
864. How many unpaired electrons are present in the central metal ion of [CoCl<sub>4</sub>]<sup>2-</sup>?  
 a) 2                      b) 3                      c) 4                      d) 5
865. The brown ring complex compound is formulated as [Fe(H<sub>2</sub>O)<sub>5</sub>NO]SO<sub>4</sub>. The oxidation state of Fe is:  
 a) +1                      b) +2                      c) +3                      d) Zero
866. Correct IUPAC name of
- is
- a) Gammexane                      b) Dichloro diphenyl trichloroethane  
 c) Diparachlorophenyl trichloroethane                      d) 1,1,1-trichloro-2,2-bis (4-chlorophenyl) ethane
867. IUPAC name of
- is
- a) Cumene                      b) 2-phenyl propane  
 c) Phenyl propane                      d) 1-(2-propyl) benzene
868. Which of the following gives violet colour with an alcoholic solution of FeCl<sub>3</sub>?  
 a) Benzoic acid                      b) Toluene                      c) Salicylic acid                      d) Nitrobenzene
869. Which of the following is wrong statements?  
 a) Ni(CO)<sub>4</sub>, has zero oxidation number for Ni                      b) Ni(CO)<sub>4</sub>, has oxidation number +4 for Ni  
 c) Ni is metal                      d) CO is gas
870. Which of the following represents a chelating ligand?  
 a) H<sub>2</sub>O                      b) Cl<sup>-</sup>                      c) OH<sup>-</sup>                      d) DMG
871. The correct order of reactivity of PhMgBr with;
- (I)

(II)

(III)
- is:  
 a) I > II > III  
 b) III > I > II  
 c) II > III > I  
 d) II > I > III
872. Which of the following will give maximum number of isomers?  
 a) [Co(NH<sub>3</sub>)<sub>4</sub>Cl<sub>2</sub>]                      b) [Ni(en)(NH<sub>3</sub>)<sub>4</sub>]<sup>2+</sup>                      c) [Ni(C<sub>2</sub>O<sub>4</sub>)(en)<sub>2</sub>]                      d) [Cr(SCN)<sub>2</sub>(NH<sub>3</sub>)<sub>4</sub>]<sup>+</sup>
873. CuCl reacts with KCN solution forming a complex. Coordination number of copper in the complex is:

- a) 2                                      b) 3                                      c) 4                                      d) 6

874. The terms stereoisomers, enantiomers and diastereomers will refer

- a) Only to configurational isomers including geometric isomers  
 b) Only to configurational isomers  
 c) To both configurational as well as conformational isomers  
 d) To neither configuration nor conformational isomers

875. Aniline was acetylated. The product on nitration followed by alkaline hydrolysis gave:

- a) *o*-nitroacetanilide              b) *o*- and *p*-nitroaniline              c) *m*-nitroaniline              d) Acetanilide

876. The IUPAC name of the compound  $[\text{CuCl}_2(\text{CH}_3\text{NH}_2)_2]$  is

- a) Dichloro bis (dimethyl amine) copper(II)              b) Dichloro bis (methyl amine) copper(II)  
 c) Dimethyl amine copper (II) chloride              d) Bis (dimethyl amine ) copper (II) chloride

877. Which is the structure of compound 2-(1-cyclobutenyl)-1-hexene?



878. On explosion TNT gives:

- a)  $\text{CO} + \text{N}_2 + \text{H}_2 + \text{CH}_4 + \text{CO}_2$   
 b)  $\text{CO} + \text{N}_2 + \text{H}_2$   
 c)  $\text{CO}_2 + \text{N}_2 + \text{H}_2\text{O}$   
 d)  $\text{CO} + \text{N}_2 + \text{H}_2\text{O}$


879. Hexafluoroferrate(III) ion is an outer orbital complex. The number of unpaired electrons present in it is:

- a) 1                                      b) 5                                      c) 4                                      d) Unpredictable

880. The EAN of Fe in  $\text{K}_3[\text{Fe}(\text{CN})_6]$  is:

- a) 36                                      b) 37                                      c) 38                                      d) 35

881. The IUPAC name of the compound



- a) 4-methyl cyclopent-1-en-3-ol                                      b) 5-methyl cyclopent-2-en-1-ol  
 c) 2-methyl cyclopent-4-en-1-ol                                      d) 3-methyl cyclopent-1-en-2-ol

882. Which one amongst the following, exhibit geometrical isomerism?

- a)  $[\text{Co}^{\text{III}}(\text{NH}_3)_5\text{Br}]\text{SO}_4$               b)  $\text{Co}^{\text{III}}[\text{EDTA}]^{1-}$               c)  $[\text{Cr}^{\text{III}}(\text{SCN})_6]^{3-}$               d)  $[\text{Pt}^{\text{II}}(\text{NH}_3)_2\text{Cl}_2]$

883. Chiral molecules are those which are

- a) Superimposable on their mirror images                                      b) Non-superimposable on their mirror images  
 c) Unstable molecules                                      d) Capable of showing geometrical isomerism

884. At room temperature the eclipsed and the staggered forms of ethane cannot be isolated because

- a) Both the conformers are equally stable                                      b) They interconvert rapidly  
 c) There is a large energy barrier of rotation about the  $\sigma$ -bond                                      d) The energy difference between the conformers is large

885. A group of atoms can function as a ligand only when

- a) It is a small molecule                                      b) It has an unshared electron pair  
 c) It is a negatively charged ion                                      d) It is a positively charged ion


886. The IUPAC name of  $\text{Ni}(\text{CO})_4$  is:

- a) Tetracarbonyl nickelate(0)

- b) Tetracarbonyl nickelate(II)  
 c) Tetracarbonyl nickel(0)  
 d) Tetracarbonyl nickel(II)
887. 2-methyl phenol is:  
 a) *o*-cresol                      b) Catechol                      c) *p*-cresol                      d) *m*-cresol
888.  $\text{NH}_2 \cdot \text{NH}_2$  serves as:  
 a) Monodentate ligand      b) Chelating ligand      c) Bridging ligand      d) Both (a) and (c)
889. For blasting purpose TNT is mixed with:  
 a)  $\text{NH}_4\text{Cl}$                       b)  $\text{NH}_4\text{NO}_3$                       c)  $\text{NH}_4\text{NO}_2$                       d)  $(\text{NH}_4)_2\text{SO}_4$
890. During the debromination of *meso*-dibromobutane, the major compound formed will be  
 a) *cis*-2-butene                      b) 1-butene                      c) *n*-butane                      d) *trans*-2-butene
891. The IUPAC name of  $\text{K}_2[\text{Cr}(\text{CN})_2\text{O}_2(\text{O})_2(\text{NH}_3)]$  is  
 a) Potassium ammine dicyano dioxoperoxochromate (VI)      b) Potassium ammine cyano peroxodioxochromium(VI)  
 c) Potassium ammine cyano peroxodioxochromium(V)      d) Potassium ammine cyano peroxodioxochromatic (IV)
892. Benzene on reaction with a mixture of  $\text{HNO}_3$  and  $\text{H}_2\text{SO}_4$  followed by reaction of  $\text{Cl}_2/\text{FeCl}_3$  gives:  
 a) 3-chloro-1-nitrobenzene  
 b) 2-chloro-1-nitrobenzene  
 c) 4-chloro-1-nitrobenzene  
 d) A mixture of 2-chloro and 4-chloro-1-nitrobenzene
893. The number of isomeric forms in which  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+$  ion can occur is:  
 a) 2                      b) 3                      c) 4                      d) 1
894. Nitration of benzene is:  
 a) Nucleophilic substitution  
 b) Electrophilic substitution  
 c) Electrophilic addition  
 d) Nucleophilic addition
895. Reimer-Tiemann reaction involves a:  
 a) Carbonium ion intermediate  
 b) Carbene intermediate  
 c) Carbanion intermediate  
 d) Free radical intermediate
896. Which does not have a carboxyl group?  
 a) Picric acid                      b) Ethanoic acid                      c) Aspirin                      d) Benzoic acid
897. In Cannizaro's reaction given below:  

$$2\text{PhCHO} \xrightarrow{\text{:}\ddot{\text{O}}\text{H}} \text{PhCH}_2\text{OH} + \text{PhC}\ddot{\text{O}}_2^-$$
 the slowest step is:  
 a) The transfer of hydride to the carbonyl group  
 b) The abstraction of proton from the carboxylic group  
 c) The deprotonation of  $\text{PhCH}_2\text{OH}$   
 d) The attack of  $\text{:}\ddot{\text{O}}\text{H}$  at the carboxyl group
898. The oxidation state of Ag in Tollens' reagent is:  
 a) Zero                      b) +1                      c) +2                      d) +1.5
899. Hybridization of Fe in  $[\text{K}_3\text{Fe}(\text{CN})_6]$  is  
 a)  $sp^3$                       b)  $d^2sp^3$                       c)  $sp^3d^2$                       d)  $dsp^3$
900. Which of the following is not isomeric with diethyl ether?  
 a) Methyl *n*-propyl ether                      b) Butan-1-ol  
 c) 2-methyl propan-2-ol                      d) Butan-2-one

901. In the given conformation  $C_2$  is rotated about  $C_2 - C_3$  bond anticlockwise by an angle of  $120^\circ$  then the conformation obtained is





- a) Fully eclipsed conformation  
 b) Partially eclipsed conformation  
 c) Gauche conformation  
 d) Staggered conformation
902. Crystal field stabilization energy for high spin  $d^4$  octahedral complex is:  
 a)  $-1.8 \Delta_0$   
 b)  $-1.6 \Delta_0 + P$   
 c)  $-1.2 \Delta_0$   
 d)  $-0.6 \Delta_0$
903. Which kind of isomerism is exhibited by octahedral  $[\text{Co}(\text{NH}_3)_4\text{Br}_2]\text{Cl}$ ?  
 a) Geometrical and ionisation  
 b) Geometrical and optical  
 c) Optical and ionisation  
 d) Geometrical only
904. The IUPAC name of the following compound is
- 
- a) 5-cyclopropyl pent-2-en-1-oic acid  
 b) 6-cyclopropyl pent-2-en-1-oic acid  
 c) 5-cyclopropyl pent-1-en carboxylic acid  
 d) 6-cyclopropyl pent-1-en carboxylic acid
905. Which of the following compounds will show a negative test with phenyl hydrazine?  
 a) Glucose  
 b) Ethyl alcohol  
 c) A cetaldehyde  
 d) Benzophenone
906. Friedel-Craft's reaction is not possible in:  
 a)  $\text{C}_6\text{H}_5\text{OH}$   
 b)  $\text{C}_6\text{H}_5\text{C}_2\text{H}_5$   
 c)  $\text{C}_6\text{H}_5\text{NO}_2$   
 d)  $\text{C}_6\text{H}_5\text{CH}_3$
907. The geometry of  $\text{Ni}(\text{CO})_4$  and  $\text{Ni}(\text{PPh}_3)_2\text{Cl}_2$  are  
 a) Both square planar  
 b) Tetrahedral and square planar respectively  
 c) Both tetrahedral  
 d) Square planar and tetrahedral respectively
908. The number of isomers possible for square planar complex  $\text{K}_2[\text{PdClBr}_2\text{SCN}]$  is:  
 a) 2  
 b) 3  
 c) 4  
 d) 6
909. The correct order for the wavelength of absorption in the visible region is  
 a)  $[\text{Ni}(\text{NO}_2)_6]^{4-} < [\text{Ni}(\text{NH}_3)_6]^{2+} < [\text{Ni}(\text{H}_2\text{O})_6]^{2+}$   
 b)  $[\text{Ni}(\text{NH}_3)_6]^{2+} < [\text{Ni}(\text{H}_2\text{O})_6]^{2+} < [\text{Ni}(\text{NO}_2)_6]^{4-}$   
 c)  $[\text{Ni}(\text{H}_2\text{O})_6]^{2+} < [\text{Ni}(\text{NH}_3)_6]^{2+} < [\text{Ni}(\text{NO}_2)_6]^{4-}$   
 d)  $[\text{Ni}(\text{NO}_2)_6]^{4-} < [\text{Ni}(\text{H}_2\text{O})_6]^{2+} < [\text{Ni}(\text{NH}_3)_6]^{2+}$
910. The IUPAC name of  $\text{CCl}_3\text{CH}_2\text{CHO}$  is  
 a) Chloral  
 b) 1,1,1-trichloropropanol  
 c) 2,2,2-trichloropropanol  
 d) 3,3,3-trichloropropanol
911. The coordination number of Cu in  $[\text{Cu}(\text{H}_2\text{O})_4]^{2+}$  complex is  
 a) 2  
 b) 1  
 c) 3  
 d) 4
912. Among the following, the correct statement is  
 a) Prefixes are written before the name of compound  
 b) Suffixes are written after the name of compound  
 c) The IUPAC name is always written as a single word  
 d) All of the above
913. In which of the following  $p$ -electrons of the halogens are not involved in delocalisation?  
 a) Chlorobenzene  
 b) Bromobenzene  
 c) Allyl chloride  
 d) Vinyl chloride
914. Which of the following does not have optical isomer?  
 a)  $[\text{Co}(\text{en})(\text{NH}_3)_2\text{Cl}_2]\text{Cl}$   
 b)  $[\text{Co}(\text{en})_2\text{Cl}_2]\text{Cl}$   
 c)  $[\text{Co}(\text{NH}_3)_3\text{Cl}]$   
 d)  $[\text{Co}(\text{en})_3]\text{Cl}_3$
915. Ethylene diamine is an example of

- a) Monodentate ligand    b) Bidentate ligand    c) Tridentate ligand    d) Polydentate ligand

916. In chlorobenzene, the —Cl group:

- a) Activates the benzene ring more via resonance effect than deactivating it via inductive effect  
 b) Deactivates the benzene ring more via inductive effect than activating it via resonance effect  
 c) Activates the benzene ring via resonance effect and deactivates it via inductive effect. Both these effects are more evenly matched  
 d) None of the above

917. The *R*-isomer among the following are




- a) (i) and (ii)                      b) (ii) and (iii)                      c) (iii) and (iv)                      d) (i) and (iii)

918. Which possesses tetrahedral shape ( $sp^3$ -hybridization of central atom)?

- a)  $[\text{Zn}(\text{NH}_3)_4]^{2+}$                       b)  $[\text{Ni}(\text{CO})_4]$                       c)  $[\text{Cd}(\text{NH}_3)_4]^{2+}$                       d) All are correct

919. The reaction,



- a) Benzoin condensation  
 b) Claisen condensation  
 c) Perkin's reaction  
 d) Cannizaro's reaction

920. Complexation is shown by:

- a) Ag                      b) Au                      c) Cu                      d) All of these


921. AgO in Ag(II) complex which is:




- a) Diamagnetic                      b) Paramagnetic                      c) Ferromagnetic                      d) Neutral

922. Acylation of benzene to produce aliphatic aromatic ketones is called:

- a) Benzoin condensation  
 b) Hydroformylation  
 c) Friedel-Crafts reaction  
 d) None of these

923. The structure of the major product formed in the given reaction





924. Chlorobenzene is prepared commercially by:

- a) Grignard reaction      b) Raschig process      c) Wurtz - Fittig reaction      d) Friedel-Crafts reaction

925. An aqueous solution of  $\text{CoCl}_2$  on addition of excess of concentrated  $\text{HCl}$  turns blue to formation of

- a)  $[\text{CoCl}_4]^{2-}$       b)  $[\text{Co}(\text{H}_2\text{O})_2\text{Cl}_4]^{2-}$       c)  $[\text{Co}(\text{H}_2\text{O})_22\text{Cl}_4]^{2-}$       d)  $[\text{Co}(\text{H}_2\text{O})_4\text{Cl}_2]$


926. Which one of the following will not show geometrical isomerism?

- a)  $[\text{Cr}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$       b)  $[\text{Co}(\text{en})_2\text{Cl}_2]\text{Cl}$       c)  $[\text{Co}(\text{NH}_3)_5\text{NO}_2]\text{Cl}_2$       d)  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$

927. When ethyl benzoate is hydrolysed with aqueous alkali, the products present in the medium are:

- a)  $\text{C}_6\text{H}_5\text{COOH}$ ,  $\text{C}_2\text{H}_5\text{O}^-$       b)  $\text{C}_6\text{H}_5\text{COO}^-$ ,  $\text{C}_6\text{H}_5\text{OH}$       c)  $\text{C}_2\text{H}_5\text{OH}$ ,  $\text{C}_6\text{H}_5\text{COOH}$       d)  $\text{C}_6\text{H}_5\text{COO}^-$ ,  $\text{C}_2\text{H}_5\text{O}^-$

928. The IUPAC name of






- a) 2-carbamoyl hexanal      b) 2-carbamoyl hex-3-en-1-al  
c) 6-keto-2-methylhexanamide      d) 5-formyl-2-methylpent-3-en-1-amide



929. Which of the following is more basic than aniline?

- a) *p*-Nitroaniline      b) Benzylamine      c) Diphenylamine      d) Triphenylamine

930. Name of some compounds are given below. Which one is not in IUPAC system?



2-ethyl-3-methyl-1-butene



3-methyl-4-ethylheptane

931. For which transition metal ions are low spin complexes possible?

- a)  $\text{Rh}^{3+}$       b)  $\text{Mn}^{3+}$       c)  $\text{Ru}^{2+}$       d) All are correct

932. Which one is monodentate ligand?

- a)  $\text{F}^-$       b)  $\text{NO}_2^-$       c)  $\text{H}_2\text{O}$       d) All are correct

933. Cyclic hydrocarbon molecule A has all the carbons and hydrogens in a single plane. All the carbon-carbon bonds are of same length and less than  $1.54 \text{ \AA}$  and more than  $1.34 \text{ \AA}$ . The C—C—C bond angle will be:

- a)  $120^\circ$       b)  $180^\circ$       c)  $100^\circ$       d)  $109^\circ 28'$

934. Chlorine reacts with benzaldehyde to give:

- a) Benzyl chloride      b) Benzal chloride      c) Benzoyl chloride      d) Chlorobenzene
935. Phenol is:
- a) A base weaker than  $\text{NH}_3$   
 b) An acid stronger than carbonic acid  
 c) An acid weaker than carbonic acid  
 d) Neutral

936. Which one is example of octahedral complex?

- a)  $\text{Cu}(\text{NH}_3)_4^{2+}$       b)  $\text{FeF}_6^{3-}$       c)  $\text{Zn}(\text{NH}_3)_4^{2+}$       d)  $\text{Ni}(\text{CN})_4^{2-}$

937. Which one of the following statement is correct?

- a) Ferric ions give a deep green precipitate on adding potassium ferrocyanide solution.  
 b) On boiling a solution having  $\text{K}^+$ ,  $\text{Ca}^{2+}$  and  $\text{HCO}_3^-$  ions, we get a precipitate of  $\text{K}_2\text{Ca}(\text{CO}_3)_2$   
 c) Manganese salt give a violet vortex test in reducing flame  
 d) From a mixed precipitate of  $\text{AgCl}$  and  $\text{AgI}$ , ammonia solution dissolves only  $\text{AgCl}$

938. Which of the following fractions obtained in fractional distillation of coal-tar contains benzene and toluene?

- a) Light oil  
 b) Heavy oil  
 c) Middle oil  
 d) Green oil

939. The tetrahedral complexes have coordination number

- a) 3      b) 6      c) 4      d) 8


940. The C—C bond length in benzene is .....than C—C bond length in alkenes.

- a) Less      b) More      c) Equal      d) None of these

941. Which are generally used for preparing derivative of aldehydes and ketones?

- a) Hydroxylamine hydrochloride  
 b) 2,4-dinitrophenylhydrazine  
 c) Phenylhydrazinehydrochloride  
 d) All of the above

942. In the reaction,





The compounds (A), (B) and (C) are the following:




- a) Benzene, nitrobenzene and aniline  
 b) Benzene, dinitrobenzene and *m*-nitroaniline  
 c) Toluene, *m*-nitrobenzene and *m*-toluidine  
 d) Benzene, nitrobenzene and hydrazobenzene

943. En is an example of a:





- a) Monodentate ligand      b) Bidentate ligand      c) Tridentate ligand      d) Hexadentate ligand

944. The major product obtained when  $\text{Br}_2/\text{Fe}$  is treated with





945. Phenol on treatment with dil.  $\text{HNO}_3$  at room temperature gives:



946. In an octahedral structure, the pair of  $d$ -orbitals involved in  $d^2 sp^3$  hybridisation is

- a)  $d_{x^2-y^2}, d_{z^2}$       b)  $d_{xz}, d_{x^2-y^2}$       c)  $d_{z^2}, d_{xz}$       d)  $d_{xy}, d_{yz}$

947. In which of the following ions has the metal atom EAN as 36?

- a)  $[\text{Fe}(\text{CN})_6]^{4-}$       b)  $[\text{Fe}(\text{CN})_6]^{3-}$       c)  $[\text{PbCl}_4]^{2-}$       d)  $[\text{Pd}(\text{CN})_6]^{2-}$

948. The number of ions given by  $\text{K}[\text{Pt}(\text{NH}_3)_5\text{Cl}_5]$  in aqueous solution is:

- a) 2      b) 3      c) 4      d) 1

949.  $\text{CuCl}$  is sparingly soluble in  $\text{H}_2\text{O}$  but it dissolves in  $\text{KCl}$  solution due to the formation of:


- a)  $\text{K}_2(\text{CuCl}_4)$       b)  $\text{K}_3(\text{CuCl}_4)$       c)  $\text{K}(\text{CuCl}_2)$       d) None of these

950. A characteristics group test for phenolic gp. is:

- a) Libermann's nitroso reaction  
b) Coupling with diazonium salt  
c)  $aq. \text{FeCl}_3$

d) All of the above

951. Write the IUPAC name of the compound




a) 5, 6-dimethyl bicyclo [2,2,1] heptane

b) 2, 3-dimethyl bicyclo [2,2,1] heptane

c) 2, 3-dimethyl bicyclo [1,2,2] heptane

d) 3, 4-dimethyl bicyclo [2,1,2] heptane

952. Choose the correct statement from the ones given below for two anilium in:



a) II is not an acceptable canonical structure because carbonium ions are less stable than ammonium ions

b) II is not an acceptable canonical structure because it is non-aromatic

c) II is not an acceptable canonical structure because the nitrogen has 10 valence electrons

d) II is an acceptable canonical structure

953. Which of the following statements is/are incorrect?

a) Metamerism belongs to the category of structural isomerism

b) Tautomeric structures are the resonating structures of a molecule

c) The violet colouration produced by a molecule with neutral ferric chloride solution indicates the presence of an enolic group in the molecule

d) Geometrical isomerism is not shown by alkenes

954. Given are (i) cyclohexanol; (ii) acetic acid; (iii) 2, 4, 6-trinitrophenol; and (iv) phenol. In these the order of decreasing acidic character will be:

a) (iii) > (ii) > (iv) > (i)

b) (ii) > (iii) > (i) > (iv)

c) (ii) > (iii) > (iv) > (i)

d) (iii) > (iv) > (ii) > (i)

955. Phenol and benzoic acid can be distinguished by:

a) Aqueous NaHCO<sub>3</sub>

b) Aqueous NaNO<sub>3</sub>

c) Aqueous NaOH

d) Conc. H<sub>2</sub>SO<sub>4</sub>

956. The functional groups -OH, -COOH, -CHO, -OCH<sub>3</sub> attached to a chiral carbon is in the preference order

a) OH > COOH > CHO > OCH<sub>3</sub>

b) OCH<sub>3</sub> > OH > CHO > COOH

c) OCH<sub>3</sub> > OH > COOH > CHO

d) OCH<sub>3</sub> > COOH > CHO > OH

957. The hypothetical complex chloro diaquatrimmine cobalt(II) chloride can be represented as:

a) [CoCl(NH<sub>3</sub>)<sub>3</sub>(H<sub>2</sub>O)<sub>2</sub>]Cl<sub>2</sub>

b) [Co(NH<sub>3</sub>)<sub>3</sub>(H<sub>2</sub>O)Cl<sub>3</sub>]

c) [Co(NH<sub>3</sub>)<sub>3</sub>(H<sub>2</sub>O)<sub>2</sub>]Cl

d) [Co(NH<sub>3</sub>)<sub>3</sub>(H<sub>2</sub>O)<sub>3</sub>]Cl<sub>3</sub>

958. Which is expected to be paramagnetic?


a) [Ni(H<sub>2</sub>O)<sub>6</sub>]<sup>2+</sup>

b) [Ni(CO<sub>4</sub>)]

c) [Zn(NH<sub>3</sub>)<sub>4</sub>]<sup>2+</sup>

d) [Co(NH<sub>3</sub>)<sub>6</sub>]<sup>3+</sup>

959. The molecular formula of diphenyl methane



How many structural isomers are possible when one of the hydrogen is replaced by a chlorine atom?

a) 8

b) 7

c) 6

d) 4

960. Among the properties (A) reducing, (B) oxidising (C) complexing, the set of properties shown by CN<sup>-</sup> ion towards metal species is


a) A, B

b) B, C

c) C, A

d) A, B, C

961.




In the double bonds are

a) cis, cis

b) cis, trans


c) trans, cis

d) trans, trans

962. The reaction of toluene with  $\text{Cl}_2$  in presence of  $\text{FeCl}_3$  gives 'X' and the reaction in presence of light gives 'Y'. Thus, 'X' and 'Y' are:
- $X$  = benzal chloride;  $Y$  = *o*-chlorotoluene
  - $X$  = *m*-chlorotoluene;  $Y$  = *p*-chlorotoluene
  - $X$  = *o*- and *p*-chlorotoluene;  $Y$  = trichloro methyl benzene
  - $X$  = benzal chloride;  $Y$  = *m*-chlorotoluene
963. Among the following four compounds:
- Phenol
  - Methyl phenol
  - meta*-nitrophenol
  - para*-nitrophenol
964. Which gives phthalic anhydride on reaction with hot, conc.  $\text{H}_2\text{SO}_4$  in presence of Hg?
- Naphthalene
  - Phenol
  - p*-xylene
  - m*-xylene
965. *Cis-trans*-isomerism is found in square planar complexes of the molecular formula: (*a* and *b* are monodentate ligands)
- $\text{Ma}_4$
  - $\text{Ma}_3\text{b}$
  - $\text{Ma}_2\text{b}_2$
  - $\text{Mab}_3$
966. Which ion produces a small crystal field splitting (a weak ligand field)?
- $\text{I}^-$
  - $\text{Cl}^-$
  - $\text{F}^-$
  - All of these
967. Benzene undergoes substitution reaction more easily than addition because:
- It has a cyclic structure
  - It has three double bonds
  - It has six hydrogen atoms
  - Of resonance
968. Isomers have essentially identical
- Structural formula
  - Chemical properties
  - Physical properties
  - Molecular formula
969. Which of the following pair is not correctly matched?
- Absorption peak for  $[\text{Cr}^{\text{III}}(\text{NH}_3)_6]^{3+} = 21680 \text{ cm}^{-1}$
  - Effective atomic no. of Pt in  $[\text{PtCl}_6]^{2-} = 84$
  - Crystal field stabilization energy of  $d^2$  in weak ligand field =  $(-)\text{0.8 } \Delta_0$
  - Example of weak ligand field for  $d^5$  configuration =  $[\text{Mn}^{\text{II}}\text{F}_6]^{4-}$
970. Aspirin (or acetyl salicylic acid) is obtained by action of  $\text{CH}_3\text{COCl}$  with:
- Salicylic acid
  - Phenol
  - Benzaldehyde
  - Aniline
971.  $\text{CuCl}$  dissolves in ammonia forming a complex. The coordination number of copper in the complex is:
- 1
  - 2
  - 4
  - 6
972. IUPAC name of the following cycloalkane is
- 
- 8-methyl bicyclo [4,3,0] nonane
  - 1-methyl bicyclo [4,3,0] nonane
  - 3-methyl bicyclo [4,3,0] nonane
  - 4-methyl bicyclo [4,3,0] nonane
973. Schiff's bases are formed when aniline is condensed with:
- Phenols
  - Aromatic aldehydes
  - Aryl chlorides
  - Aliphatic alcohols
974. Which of the following is not an organometallic compound?
- Zeise's salt
  - TEL
  - Sodium ethoxide
  - Ferrocene
975. Molecular formula  $\text{C}_5\text{H}_{12}\text{O}$  will show
- Position
  - Optical isomerism
  - Functional isomerism
  - All of these
976. Both  $\text{Co}^{3+}$  and  $\text{Pt}^{4+}$  have a coordination number of six. Which of the following pairs of complexes will show approximately the same electrical conductance for their 0.001 M aqueous solutions?
- $\text{CoCl}_2 \cdot 4\text{NH}_3$  and  $\text{PtCl}_4 \cdot 4\text{NH}_3$
  - $\text{CoCl}_3 \cdot 3\text{NH}_3$  and  $\text{PtCl}_4 \cdot 5\text{NH}_3$
  - $\text{CoCl}_3 \cdot 6\text{NH}_3$  and  $\text{PtCl}_4 \cdot 5\text{NH}_3$
  - $\text{CoCl}_3 \cdot 6\text{NH}_3$  and  $\text{PtCl}_4 \cdot 3\text{NH}_3$



990.



The IUPAC name of the given structure

is

- a) Diisohexane                      b) Isohexane                      c) 2, 2-dimethylbutane                      d) 2, 3-dimethylbutane

991. Aniline on treating with phosgene gives:

- a) Phenyl isocyanate                      b) A secondary base                      c) A neutral substance                      d) A tertiary base

992. On boiling with conc. hydrobromic acid, phenylethylether will yield:

- a) Phenol and ethyl bromide  
b) Bromobenzene and ethanol  
c) Phenol and ethane  
d) Bromobenzene and ethane





993. Ammonia gas does not evolve from the complex  $\text{FeCl}_3 \cdot 4\text{NH}_3$  but it gives white precipitate with aqueous solution of  $\text{AgNO}_3$ . Coordination number of central metal ion in above complex is six. Give IUPAC name of the complex.

- a) Ammonium trichloro triammine ferrum(III)                      b) Tetra ammine ferrum (III) chloride  
c) Dichloro tetraammine ferrate (II) chloride                      d) Dichloro tetraammine ferrum (III) chloride






994. Nickel ( $Z=28$ ) combines with a uninegative monodentate ligand  $X^-$  to form a paramagnetic complex  $[\text{NiX}_4]^{2-}$ . The number of unpaired electron ( $s$ ) in the nickel and geometry of this complex ion are respectively

- a) One, tetrahedral                      b) Two, tetrahedral                      c) One, square planar                      d) Two, square planar

995. Amongst the compounds given, the one that would form a brilliant coloured dye on treatment with  $\text{NaNO}_2$  in dil. HCl followed by addition to an alkaline solution of  $\beta$ -naphthol is:



996. Identify 'Z' in the reaction given below;



997. A solution containing 2.675 g of  $\text{CoCl}_3 \cdot 6\text{NH}_3$  (molar mass = 267.5 g mol<sup>-1</sup>) is passed through a cation exchanger. The chloride ions obtained in solution were treated with excess of  $\text{AgNO}_3$  to give 4.78 g of  $\text{AgCl}$  (molar mass = 143.5 g mol<sup>-1</sup>). The formula of the complex is (Atomic mass of Ag = 108 u)

998.  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$  ion has *d*-electrons equal to:
- a) 2                      b) 3                      c) 4                      d) 5

999. Enol form is more stable in

- a)  $\text{CH}_3\text{CHO}$                       b)  $\text{CH}_3\text{COCH}_3$   
 c)  $\text{CH}_3\text{COCH}_2\text{COOC}_2\text{H}_5$                       d) Cyclohexanone


100 The coordination number of cobalt in  $[\text{Co}(\text{en})_2\text{Br}_2]\text{Cl}_2$  is:





0. a) 2                      b) 4                      c) 6                      d) 8

100 Which one readily accepts a proton?

1. a) Acetylene                      b) Nitrobenzene                      c) Aniline                      d) Phenol

100 Identify 'Z' in the reaction;







- a)   
 b)   
 c)   
 d) 

100 The number of  $\sigma$  and  $\pi$ -bonds in a molecule of benzene is:


3. a)  $6\sigma$  and  $9\pi$                       b)  $9\sigma$  and  $3\pi$                       c)  $12\sigma$  and  $3\pi$                       d)  $6\sigma$  and  $6\pi$

100 The phenomenon of optical activity will be shown by:

4. a)   
 b)   
 c)   
 d) 

100

5.



The correct name of the compound

- a) 1,3,4-trimethyldecaline                      b) 1,3,9-trimethyldecaline  
c) 1,8,10-trimethyldecaline                  d) 1,3,10-trimethyldecaline

100 If  $\text{NH}_4\text{OH}$  is added to the  $(\text{PtCl}_4)^{2-}$  ion, the complex formed represents:

6.

- a) Zero dipole                      b) Finite dipole                      c) Infinite dipole                      d) All of these

100 Which one of the following will be able to show *cis-trans*-isomerism?

7.

- a)  $M_{A_3B}$   
b)  $M_{(AA')_2}$   
c)  $M_{A_2BCD}$   
d)  $M_{A_4}$   
( $AA'$  is unsymmetrical bidentate ligand,  $ABCD$  are unidentate ligands.)

100 The coordination number of a metal in coordination compound is

8.

- a) Same as primary valency                      b) Sum of primary and secondary valencies  
c) Same as secondary valency                      d) None of the above

100 The IUPAC name of  $\text{K}_4[\text{Ni}(\text{CN})_4]$  is

9.

- a) Tetrapotassium tetracyanonickelate (II)                      b) Potassium tetracyanonickel (II)  
c) Potassium tetracyanonickelate (0)                      d) Potassium tetracyanonickelate (II)

101 Which of the following compounds shows optical isomerism?

0.

- a)  $[\text{Co}(\text{CN})_6]^{3-}$                       b)  $[\text{Cr}(\text{C}_2\text{O}_4)_3]^{3-}$                       c)  $[\text{ZnCl}_4]^{2-}$                       d)  $[\text{Cu}(\text{NH}_3)_4]^{2+}$

101  $[\text{C}_6\text{H}_5]_2\text{Pd}(\text{SCN})_2$  and  $[(\text{C}_6\text{H}_5)_2\text{Pd}(\text{NCS})_2]$  are:

1.

- a) Linkage isomers                      b) Coordination isomers                      c) Ionization isomers                      d) Geometrical isomers


101 Mark the correct statement

2.


- a) Ethane has two conformations of which staggered conformation is more stable than the eclipsed conformation  
b) Ethane has an infinite number of conformations of which eclipsed conformation is more stable than the staggered conformation  
c) Ethane has an infinite number of conformation of which staggered conformation has the maximum energy  
d) Ethane has an infinite number of conformation of which the staggered conformation is possessed by majority of the molecules at room temperature

101

3.



Of the following on heating with alkali followed with acid hydrolysis?



101 Which among the following will be named as dibromidobis (ethylene diamine) chromium (III) bromide?

4.

- a)  $[\text{Cr}(\text{en})_2\text{Br}_2]\text{Br}$       b)  $[\text{Cr}(\text{en})\text{Br}_4]^-$       c)  $[\text{Cr}(\text{en})\text{Br}_2]\text{Br}$       d)  $[\text{Cr}(\text{en})_3]\text{Br}_3$

101 Which one of the following complex is an outer orbital complex?

5. (Atomic no. Mn=25, Fe=24, Co=27, Ni=28)

- a)  $[\text{Fe}(\text{CN})_6]^{4-}$       b)  $[\text{Mn}(\text{CN})_6]^{4-}$       c)  $[\text{Co}(\text{NH}_3)_6]^{3+}$       d)  $[\text{Ni}(\text{NH}_3)_6]^{2+}$

101 Benzene can be directly obtained from:

6.

- a)  $\text{CH}\equiv\text{CH}$   
 b)  $\text{CH}_2=\text{CH}_2$  and butadiene  
 c) Chlorobenzene  
 d) All of the above





101 Chlorobenzene on treatment with Raney nickel or Al in presence of alkali gives:

7.

- a) Benzene      b) Chlorophenol      c) Phenol      d) None of these

101 The compound that undergoes decarboxylation most readily under mild condition is:

8.



101 Which ion is paramagnetic?

9.

- a)  $[\text{Ni}(\text{NH}_3)_4]^{2+}$       b)  $[\text{Ni}(\text{CO})_4]$       c)  $[\text{Co}(\text{NH}_3)_6]^{3+}$       d)  $[\text{Ni}(\text{CN})_4]^{2-}$

102 Which kind of isomerism is exhibited by octahedral  $[\text{Co}(\text{NH}_3)_4\text{Br}_2\text{Cl}]$ ?

0.

- a) Geometrical and ionization      b) Geometrical only  
 c) Geometrical and optical      d) Optical and ionisation

102 Resorcinol and conc.  $\text{H}_2\text{SO}_4$  in presence of phthalic anhydride produce a compound which is:

1.

- a) A dye      b) An antiseptic      c) An indicator      d) A detergent

102 Which of the following compounds shows optical isomerism?


2.

- a)  $[\text{Cr}(\text{C}_2\text{O}_4)_3]^{3-}$       b)  $[\text{Cu}(\text{NH}_3)_4]^{2+}$       c)  $[\text{Co}(\text{CN})_6]^{3-}$       d)  $[\text{ZnCl}_4]^{2-}$

102 The IUPAC name of  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$  is


3.  
 a) Hexamine cobalt (II) chloride      b) Triammine cobalt (III) trichloride  
 c) Hexamine cobalt (III) chloride      d) None of the above

102 In the following compounds, the order of acidity is:



- a) III > IV > I > II      b) I > IV > III > II      c) II > I > III > IV      d) IV > III > I > II

102 Consider the following structure and choose the correct statements




- a) I and II have *R*-configuration      b) I and III have *R*-configuration  
 c) Only III has *S*-configuration      d) Both (a) and (c) are correct

102 Benzaldehyde, when heated with concentrated KOH solution, gives:

6.  
 a)  $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$   
 b)  $\text{C}_6\text{H}_5\text{COOH}$   
 c)  $\text{C}_6\text{H}_5\text{COOK}$   
 d) Mixture of  $\text{C}_6\text{H}_5\text{COOK}$  and  $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$

102 Write the IUPAC name of the compound



- a) Bicyclo-[2.2.2] octane-2,6-dione      b) Bicyclo-[2.2.2] octane-3,5-dione  
 c) Bicyclo-[2.2] octane 2,6-dione      d) Bicyclo [2,2] octane-3,5-dione

102 3-chloro-4-methyl benzene sulphonic acid on steam distillation gives:

8.  
 a) Toluene      b) *m*-chloro benzene sulphonic acid      c) *p*-methyl benzene sulphonic acid      d) *o*-chloro toluene


102 The oxidation number of platinum in  $[\text{Pt}(\text{NH}_3)_5\text{Cl}]\text{Cl}_3$  is



9.  
 a) 2      b) 3      c) 4      d) 6

103 Which of the following is not an organometallic compound?

0.  
 a)  $\text{C}_2\text{H}_5\text{ONa}$       b)  $\text{CH}_3\text{MgI}$       c) Tetraethyl tin      d)  $\text{KC}_4\text{H}_9$

103 Which of the following pairs of compounds are enantiomers?





103 Which complex has square planar shape  $dsp^2$ -hybridization?

2. a)  $[\text{Ni}(\text{CN})_4]^{2-}$       b)  $[\text{Cu}(\text{NH}_3)_4]^{2-}$       c)  $[\text{PtCl}_4]^{2-}$       d) All of these

103 The complex used as an anticancer agent is

3. a)  $\text{cis}-[\text{PtCl}_2(\text{NH}_3)_2]$       b)  $\text{Na}_2\text{CO}_3$   
c)  $\text{trans}-[\text{Co}(\text{NH}_3)_3\text{Cl}_3]$       d)  $\text{cis}-\text{K}_2[\text{PtCl}_2\text{Br}_2]$

103 Dyes are formed when diazonium salts react with:

4. a) Phenols  
b) Aldehydes  
c) Ketones  
d) Alcohols

103 Potassium ferrocyanide is a

5. a) Complex salt      b) Double salt      c) Normal salt      d) Mixed salt

103 The primary and secondary valencies of chromium in the complex ion, dichlorodioxalatochromium (III), are respectively.

6. a) 3,4      b) 4,3      c) 3,6      d) 6,3

103 The reaction,  $\text{C}_6\text{H}_6 + \text{CH}_3\text{Cl} \xrightarrow[\text{AlCl}_3]{\text{Anhydrous}} \text{C}_6\text{H}_5\text{CH}_3 + \text{HCl}$  is an example of:

7. a) Friedel-Craft's reaction  
b) Kolbe's synthesis  
c) Wurtz's reaction  
d) Grignard synthesis

103 The correct statement related to IUPAC nomenclature is

8. a) If 2 or more chains of equal length are seen in the compound then the chain with minimum number of side chains will be preferred  
b) If double and triple bonds are at symmetrical positions in a compound then triple bond gets lower preference  
c) Correct IUPAC name of  $\text{CH}_3\text{COC}_2\text{H}_5$  is ethyl methyl ketone  
d) As far as possible, the IUPAC name of a compound is written as a single word


103 Which of the following isomerism is shown by ethyl acetoacetate?





9. a) Geometrical isomerism      b) Keto-enol tautomerism  
c) Enantiomerism      d) Diastereoisomerism

104 The number of moles of ions given on complete ionisation of one mole of  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$  is/are

0. a) 4      b) 3      c) 2      d) 1

104 The major products (P, Q) in the given reaction are:





104 *p*-nitrophenol is stronger acid than phenol because nitro group is:

2. a) Electron withdrawing    b) Electron donating    c) Basic    d) Acidic

104 Among the following group, which deactivates benzene ring for electrophilic substitution:

3. a) Methyl    b) Amino    c) Hydroxyl    d) Chlorine

104 Iron has lowest oxidation state in:

4. a)  $\text{Fe}(\text{CO})_5$   
 b)  $\text{Fe}_2\text{O}_3$   
 c)  $\text{K}_2\text{FeO}_4$   
 d)  $\text{FeSO}_4 \cdot (\text{NH}_4)_2\text{SO}_4 \cdot 6\text{H}_2\text{O}$

104 The dihedral angle between the two methyl groups in Gauch conformation of *n* butane is

5. a)  $120^\circ$     b)  $180^\circ$     c)  $45^\circ$     d)  $60^\circ$

104 Increasing order of acid strength among *p*-methoxyphenol, *p*-methylphenol and *p*-nitrophenol is:

6. a) *p*-methylphenol < *p*-methoxyphenol < *p*-nitrophenol  
 b) *p*-methoxyphenol < *p*-methylphenol < *p*-nitrophenol  
 c) *p*-nitrophenol < *p*-methoxyphenol < *p*-methylphenol  
 d) *p*-nitrophenol < *p*-methylphenol < *p*-methoxyphenol

104 Total number of isomeric alcohols with formula  $\text{C}_4\text{H}_{10}\text{O}$  are

7. a) 1    b) 2    c) 3    d) 4

104 What is the IUPAC name of  $\text{Na}_2[\text{Fe}(\text{CN})_5\text{NO}]$ ?


8. a) Pentacyanonitroso sodium ferrate    b) Pentacyanonitroso sodium ferrate(II)  
 c) Sodium pentacyanonitroso ferrate(II)    d) Sodium pentacyanonitroso ferrate

104 Which of the following cations does not form an ammine complex with excess of ammonia?

9. a)  $\text{Ag}^+$     b)  $\text{Cu}^{2+}$     c)  $\text{Cd}^{2+}$     d)  $\text{Na}^+$

105 In the complex  $\text{K}_2[\text{Fe}(\text{CN})_6]$

- 0.
- a) The complex is high spin complex  
 c) The coordination number of iron is 4
- b) Both Fe atoms are in the same oxidation state  
 d) Both Fe atoms are in different oxidation state
- 105 The number of chiral carbon atoms present in the molecule




- a) 3  
 b) 4  
 c) 2  
 d) 1
- 105 The complex that doesn't give a precipitate with  $\text{AgNO}_3$  solution
- 2.
- a)  $[\text{Co}(\text{NH}_3)_3\text{Cl}_3]$   
 b)  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$   
 c)  $[\text{Ag}(\text{NH}_3)_2]\text{Cl}$   
 d)  $[\text{Cr}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$
- 105 The IUPAC name of the given compound  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$  is

- 3.
- a) Penta amino cobalt chloride chlorate  
 c) Pentamine chloro cobalt (III) chloride.
- b) Cobalt penta ammine chloro chloride  
 d) Penta amino cobalt (III) chlorate

- 105 Amongst  $\text{Ni}(\text{CO})_4$ ,  $[\text{Ni}(\text{CN})_4]^{2-}$  and  $[\text{NiCl}_4]^{2-}$
- 4.

- a)  $\text{Ni}(\text{CO})_4$  is diamagnetic,  $[\text{NiCl}_4]^{2-}$  and  $[\text{Ni}(\text{CN})_4]^{2-}$  are paramagnetic  
 b)  $\text{Ni}(\text{CO})_4$  and  $[\text{NiCl}_4]^{2-}$  are diamagnetic and  $[\text{Ni}(\text{CN})_4]^{2-}$  is paramagnetic  
 c)  $\text{Ni}(\text{CO})_4$  and  $[\text{Ni}(\text{CN})_4]^{2-}$  are diamagnetic and  $[\text{NiCl}_4]^{2-}$  is paramagnetic  
 d)  $[\text{NiCl}_4]^{2-}$  and  $[\text{Ni}(\text{CN})_4]^{2-}$  are diamagnetic and  $\text{Ni}(\text{CO})_4$  is paramagnetic


- 105 Which aromatic acid among the following is weaker than simple benzoic acid?




- 105 Which statement is incorrect?

- 6.
- a)  $\text{Ni}(\text{CO})_4$ -tetrahedral, paramagnetic  
 b)  $[\text{Ni}(\text{CN})_4]^{2-}$ -square planar, diamagnetic  
 c)  $\text{Ni}(\text{CO})_4$ -tetrahedral, diamagnetic  
 d)  $[\text{NiCl}_4]^{2-}$ -tetrahedral, paramagnetic

- 105 Which of the following has asymmetric C-atom?



- 105 The IUPAC name of



- a) 1-phenyl-3-propanol  
 c) 1-hydroxy-3-phenyl-propane
- b) 3-phenyl-1-propanol  
 d) None of the above
- 105 The complexes  $[\text{Co}(\text{NH}_3)_6][\text{Cr}(\text{CN})_6]$  and  $[\text{Cr}(\text{NH}_3)_6][\text{Co}(\text{CN})_6]$  are the examples of which type of

9. isomerism?  
 a) Geometrical isomerism  
 b) Linkage isomerism  
 c) Ionization isomerism  
 d) Coordination isomerism
- 106 Racemic tartaric acid is optically inactive due to  
 0.  
 a) External compensation  
 b) Internal compensation  
 c) Presence of plane of symmetry  
 d) All of the above
- 106 Nitration of aniline is done in:  
 1.  
 a) Acidic medium  
 b) Alkaline medium  
 c) Neutral medium  
 d) In acidic medium by first converting it into acetanilide before nitration
- 106 A bridging ligand possesses:  
 2.  
 a) Polydentate or monodentate nature  
 b) Two or more donor centres  
 c) The tendency to get itself attached to two metal ions  
 d) All of the above
- 106 What is the neutralization equivalent of benzoic acid?  
 3.  
 a) 122                                      b) 61                                      c) 244                                      d) 488
- 106 *m*-chlorobenzaldehyde on reaction with conc. KOH at room temperature gives:  
 4.  
 a) Potassium *m*-chlorobenzoate and *m*-hydroxy benzaldehyde  
 b) *m*-hydroxybenzaldehyde and *m*-chlorobenzylalcohol  
 c) *m*-chlorobenzylalcohol and *m*-hydroxy benzylalcohol  
 d) Potassium *m*-chlorobenzoate and *m*-chlorobenzyl alcohol
- 106 The oxidation number of Fe in brown ring  $[\text{Fe}(\text{H}_2\text{O})_5\text{NO}]^{2+}$  is  
 5.  
 a) 0                                      b) +1                                      c) +2                                      d) +3
- 106  $[\text{Cr}(\text{H}_2\text{O})_6]\text{Cl}_3$  (at. No. of Cr=24) has a magnetic moment of 3.83 BM. The correct distribution of 3d-  
 6. electrons in the chromium of the complex:  
 a)  $3d_{xy}^1, 3d_{yz}^1, 3d_{xz}^1$                       b)  $3d_{xy}^1, 3d_{yz}^1, 3d_{z^2}^1$                       c)  $(3d_{x^2-y^2}^1), 3d_{z^2}^1, 3d_{xz}^1$                       d)  $3d_{xy}^1, (3d_{x^2-y^2}^1), 3d_{yz}^1$
- 106 Excess of silver nitrate solution is added to 100 mL of 0.01 M pentaqua chloro chromium (III) chloride  
 7. solution. The mass of silver chloride obtained in grams is [Atomic mass of silver is 108].  
 a)  $287 \times 10^{-3}$                       b)  $143.5 \times 10^{-3}$                       c)  $143.5 \times 10^{-2}$                       d)  $287 \times 10^{-2}$
- 106 The total number of possible structural isomers of the compound  $[\text{Cu}^{\text{II}}(\text{NH}_3)_4][\text{Pt}^{\text{II}}\text{Cl}_4]$  are:  
 8.  
 a) 3                                      b) 5                                      c) 4                                      d) 6
- 106 A similarity between optical and geometrical isomerism is that :  
 9.  
 a) Each gives equal number of isomers for a given compound  
 b) If in a compound one is present then so is the other  
 c) Both are included in stereoisomerism  
 d) They have no similarity
- 107 In  $[\text{Ni}(\text{NH}_3)_4]\text{SO}_4$ , the valency and coordination number of Ni will be respectively  
 0.

- a) 3 and 6                      b) 4 and 4                      c) 4 and 2                      d) 2 and 4
- 107  $C_6H_5CHO$  is different from aliphatic aldehyde in its reaction towards:
1. a) Tollen's reagent              b) Schiff's reagent              c)  $NaHSO_3$               d) Fehling's solution

107 Oxidation of naphthalene by acidic  $KMnO_4$  gives:

2. a) Toluene                      b) Benzaldehyde              c) Phthalic acid              d) Benzoic acid

107 The number of possible theoretical conformations of *n*-butane are

3. a) Two                      b) Three                      c) Four                      d) Infinite

107 Which is correct order for acidic nature of following acids?

4. (I)  $PhCOOH$                       (II)  $o-NO_2C_6H_4COOH$   
 (III)  $p-NO_2C_6H_4COOH$               (IV)  $m-NO_2C_6H_4COOH$
- a)  $II > III > IV > I$               b)  $II > IV > III > I$               c)  $II > IV > I > III$               d)  $I > II > III > IV$

107 Salicylic acid when treated with zinc dust gives:

5. a) Phenol                      b) Salicylaldehyde              c) Benzene                      d) Benzoic acid


107 Action of  $PCl_5$  on salicylic acid produces:

6. a) *o*-chlorobenzoyl  
 b) *o*-hydroxybenzoyl chloride  
 c) *o*-chlorobenzoic acid  
 d) None of the above

107 Which of the following species is most stable?

7. a)  $p-O_2N-C_6H_4-\overset{+}{C}H_2$   
 b)  $C_6H_5-\overset{+}{C}H_2$   
 c)  $p-Cl-C_6H_4-\overset{+}{C}H_2$   
 d)  $p-CH_3O-C_6H_4-\overset{+}{C}H_2$

107 Give the IUPAC name of the following



- a) 5-ethyl-4, 4-dimethyloctane                      b) 4-ethyl-5, 5-dimetyloctane  
 c) 3-ethyl-2-methyl-2-propyl hexane                      d) 4-ethyl-5-methyl, 5-propyl hexane

107 Which of the following reacts with KCN to form benzoin?

9. a)  $C_6H_5CHO$                       b)  $C_6H_5Cl$                       c)  $C_2H_5Cl$                       d)  $C_6H_5CH_3$

108 Which one is an organometallic compound in the following?



0. a)  $C_2H_5ONa$                       b)  $C_2H_5-S-S-C_2H_5$   
 c)  $Al_2(CH_3)_6$                       d)  $Al(C_6H_5S)_3$

108 The formula of picramide is:

1.



- 2.
- To absorb water
  - To absorb hydrochloric acid
  - To produce an electrophile
  - To produce nucleophile
- 109 Coordination isomerism is caused by interchange of ligands between the
- 3.
- Complex cation and complex anion
  - Inner sphere and outer sphere
  - Low oxidation and higher oxidation states
  - cis* and *trans* structure
- 109 Which aldehyde is used in the manufacture of perfumes?
- 4.
- Cinnamaldehyde
  - Salicylaldehyde
  - Benzaldehyde
  - None of these
- 109 Which of the following statements is not correct?
- 5.
- A *meso* compound has chiral centres but exhibits no optical activity
  - A *meso* compound has no chiral centres and thus are optically inactive
  - A *meso* compound has molecules in which one half of molecule is superimposable on the other even through chiral centre is present in them
  - A *meso* compound is optically inactive because the rotation caused by one half of molecule is cancelled by the rotation produced by another half
- 109 The volume (in mL) of 0.1 M  $\text{AgNO}_3$  required for complete precipitation of chloride ions present in 30 mL of 0.01 M solution of  $[\text{Cr}(\text{H}_2\text{O})_5\text{Cl}]\text{Cl}_2$ , as silver chloride is close to
- 3
  - 4
  - 5
  - 6
- 109 Benzene is a resonance hybrid mainly of two Kekule structures. Hence:
- 7.
- Half of the molecules correspond to one structure, and half to the second structure
  - At low temperatures benzene can be separated into two structures
  - Two structures make equal contribution to resonance hybrid
  - An individual benzene molecule changes back and forth between two structures
- 109 Keto form is more stable in
- 8.
- $\text{CH}_3\text{COCH}_2\text{COOC}_2\text{H}_5$
  - $\text{CH}_3\text{COCH}_2\text{COCH}_3$
  - $\text{CH}_3\text{COCH}_3$
  - $\text{CH}_3\text{COCH}_2\text{COC}_2\text{H}_5$
- 109 The oxidation state and effective atomic number (EAN) of cobalt ( $\text{CoF}_6$ )<sup>2-</sup> are respectively
- 9.
- 3 and 36
  - 4 and 35
  - 4 and 37
  - 2 and 35
- 110 Benzamide on reaction with  $\text{POCl}_3$  gives:
- 10.
- Aniline
  - Chlorobenzene
  - Benzylamine
  - Benzonitrile
- 110 Which pair of carbon skeleton is an example of isomerism?
- 1.
- $$\begin{array}{c} \text{C} \\ | \\ \text{C} - \text{C} - \text{C} - \text{C} \end{array}$$
and
$$\begin{array}{c} \text{C} \\ | \\ \text{C} - \text{C} - \text{C} \\ | \\ \text{C} \end{array}$$
  - $$\begin{array}{c} \text{C} \\ | \\ \text{C} - \text{C} - \text{C} \\ | \quad | \\ \text{C} \quad \text{C} \end{array}$$
and
$$\text{C} - \text{C} - \text{C}$$



110 Electrolytic reduction of nitrobenzene in weak acidic medium gives:

2.

- a) Aniline
- b) *p*-hydroxy aniline
- c) Nitrobenzene
- d) *N*-phenyl hydroxyl amine

110 In complexes, metal atom acts as:

3.

- a) Lewis base
- b) Bronsted acid
- c) Bronsted base
- d) Lewis acid

110 When benzene is treated with concentrated  $\text{HNO}_3$  at room temperature it will give:

4.

- a)  $\text{CO}_2$  and  $\text{H}_2\text{O}$
- b) Nitrochlorobenzene
- c) Dark red colour
- d) Dinitrobenzene


110 Which of the following compounds exhibit linkage isomerism?

5.

- a)  $[\text{Co}(\text{en})_3]\text{Cl}_3$
- b)  $[\text{Co}(\text{NH}_3)_6][\text{Cr}(\text{CN})_6]$
- c)  $[\text{Co}(\text{en})_2\text{NO}_2\text{Cl}]\text{Br}$
- d)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Br}_2$

110 The compound

6.



have IUPAC name as

- a) Tricyclopropyl
- b) Tricyclopropane
- c) 1,1', 2', 1''-tercyclo propane
- d) None of the above

110 The most stable conformation of chlorohydrin at room temperature is

7.

- a) Fully eclipsed
- b) Partially eclipsed
- c) Gauche
- d) Staggered

110 Among  $[\text{Ni}(\text{CO})_4]$ ,  $[\text{Ni}(\text{CN})_4]^{2-}$  and  $[\text{Ni}(\text{Cl})_4]^{2-}$ :

8.

- a)  $[\text{Ni}(\text{CO})_4]$ ,  $[\text{NiCl}_4]^{2-}$  are diamagnetic and  $[\text{Ni}(\text{CN})_4]^{2-}$  is paramagnetic
- b)  $[\text{NiCl}_4]^{2-}$ ,  $[\text{Ni}(\text{CN})_4]^{2-}$  are diamagnetic and  $[\text{Ni}(\text{CO})_4]$  is paramagnetic
- c)  $[\text{Ni}(\text{CO})_4]$ ,  $[\text{Ni}(\text{CN})_4]^{2-}$  are diamagnetic and  $[\text{NiCl}_4]^{2-}$  is paramagnetic
- d)  $[\text{Ni}(\text{CO})_4]$  is diamagnetic and  $[\text{NiCl}_4]^{2-}$ ,  $[\text{Ni}(\text{CN})_4]^{2-}$  are paramagnetic

110 The complex  $\text{Hg}[\text{Co}(\text{CNS})_4]$  is correctly named as:

9.

- a) Mercury tetrathiocyanatocobaltate(II)
- b) Mercury cobalt tetrasulphocyno(II)
- c) Mercury tetrasulphocyanidecobaltate(II)
- d) Mercury sulphocyanatocobalt(II)

111 Which of the following compounds is not coloured?

10.

- a)  $\text{Na}_2[\text{CuCl}_4]$
- b)  $\text{Na}_2[\text{CdCl}_4]$
- c)  $\text{K}_4[\text{Fe}(\text{CN})_6]$
- d)  $\text{K}_3[\text{Fe}(\text{CN})_6]$

111 Which one has square planar geometry?

1. a)  $[\text{CoCl}_4]^{2-}$                       b)  $[\text{FeCl}_4]^{2-}$                       c)  $[\text{NiCl}_4]^{2-}$                       d)  $[\text{PtCl}_4]^{2-}$   
111 Which exhibits highest molar conductivity?
2. a)  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$                       b)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$                       c)  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$                       d)  $[\text{Co}(\text{NH}_3)_3\text{Cl}_3]$   
111 Coordination compounds have great importance in biological systems. In this context which of the following statement is incorrect?
3. a) Chlorophyll is green pigment in plants and contain calcium.  
b) Haemoglobin is the red pigment of blood and contains iron.  
c) Cyanocobalamin is vitamin  $\text{B}_{12}$  and contains cobalt.  
d) Carboxypeptidase-A is an enzyme and contains zinc.
- 111 Complex salt can be made by the combination of  $[\text{Co}^{\text{III}}(\text{NH}_3)_5\text{Cl}]^x$  with
4. a)  $\text{Cl}^-$                       b)  $2\text{Cl}^-$                       c)  $\text{PO}_4^{3-}$                       d)  $2\text{K}^+$   
111 Which of the following pairs represents linkage isomers?
5. a)  $[\text{Cu}(\text{NH}_3)_4][\text{PtCl}_4]$  and  $[\text{Pt}(\text{NH}_3)_4][\text{CuCl}_4]$                       b)  $[\text{Pd}(\text{PPh}_3)_2(\text{NCS})_2]$  and  $[\text{Pd}(\text{PPh}_3)_2(\text{SCN})_2]$   
c)  $[\text{Co}(\text{NH}_3)_5]\text{NO}_3\text{SO}_4$  and  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{NO}_3$                       d)  $[\text{PtCl}_2(\text{NH}_3)_4]\text{Br}_2$  and  $[\text{PtBr}_2(\text{NH}_3)_4]\text{Cl}_2$   
111 The reaction products of  $\text{C}_6\text{H}_5\text{OCH}_3 + \text{HI} \xrightarrow{\Delta}$  is:
6. a)  $\text{C}_6\text{H}_5\text{OH} + \text{CH}_3\text{I}$                       b)  $\text{C}_6\text{H}_5\text{I} + \text{CH}_3\text{OH}$                       c)  $\text{C}_6\text{H}_5\text{CH}_3 + \text{HOI}$                       d)  $\text{C}_6\text{H}_6 + \text{CH}_3\text{OI}$   
111 An aromatic amine (A) was treated with alcoholic potash and another compound (Y) when a foul smelling gas was formed with formula  $\text{C}_6\text{H}_5\text{NC}$ . (Y) was formed by reacting a compound (Z) with  $\text{Cl}_2$  in the presence of slaked lime. Compound (Z) is:
7. a)  $\text{C}_6\text{H}_5\text{NH}_2$                       b)  $\text{CH}_3\text{OH}$                       c)  $\text{CH}_3\text{COCH}_3$                       d)  $\text{CHCl}_3$   
111 Chlorine is most reactive in:
8. a)  $\text{CH}_3\text{Cl}$                       b)  $\text{CH}_2=\text{CHCl}$                       c)  $\text{C}_6\text{H}_5\text{Cl}$                       d)  $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$   
111 The C—C bond order in benzene is close to:
9. a) 1.5                      b) 2.5                      c) 3.0                      d) 6.0  
112 Mixture X = 0.02 mole of  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Br}$  and 0.02 mole of  $[\text{Co}(\text{NH}_3)_5\text{Br}]\text{SO}_4$  was prepared in 2 L of solution  
1 L of mixture X + excess  $\text{AgNO}_3 \rightarrow Y$   
1 L of mixture X + excess  $\text{BaCl}_2 \rightarrow Z$   
Number of moles of Y and Z are
- a) 0.01, 0.01                      b) 0.01, 0.02                      c) 0.02, 0.01                      d) 0.02, 0.02  
112 Phenol can be converted into salicylic acid by:
1. a) Etard's reaction  
b) Kolbe's reaction  
c) Reimer-Tiemann reaction  
d) Both (b) and (c)  
112  $\text{Fe}_2(\text{CO})_9$  is diamagnetic. Which of the following reasons is correct?
2. a) Presence of one CO as bridge group  
b) Presence of monodentate ligand  
c) Metal-metal (Fe-Fe) bond in molecule  
d) Resonance hybridization of CO  
112 The formula of dichlorobis (urea) copper(II) is:

3.

- a)  $[\text{CuO} = \text{C}(\text{NH}_2)_2]\text{Cl}_2$
- b)  $[\text{CuCl}_2\{\text{O} = \text{C}(\text{NH}_2)_2\}]$
- c)  $[\text{Cu}\{\text{O} = \text{C}(\text{NH}_2)_2\}\text{Cl}]\text{Cl}$
- d)  $[\text{CuCl}_2][\text{O} = \text{C}(\text{NH}_2)_2]\text{H}_2$






112 Which of the following facts about the complex  $[\text{Cr}(\text{NH}_3)_6]\text{Cl}_3$  is wrong?

4.

- a) The complex involves  $d^2sp^3$  hybridisation and is b) The complex is paramagnetic. octahedral in shape.
- c) The complex is an outer orbital complex. d) The complex gives white precipitate with silver nitrate solution.

112 The compounds P, Q and S were separately subjected to nitration using  $\text{HNO}_3/\text{H}_2\text{SO}_4$  mixture. The major

5. product formed in each case respectively, is:



112 Aromaticity of benzene is due to:

6.

- a) Ring
- b) Three double bonds
- c) Delocalisation of  $\pi$ -electrons



- 113 C<sub>6</sub>H<sub>6</sub> is a very good industrial solvent for:  
8.
- a) Oil                                      b) Fat                                      c) Rubber                                      d) All of these
- 113 Salol is used as:  
9.
- a) Antiseptic                                      b) Antipyretic                                      c) Both (a) and (b)                                      d) None of these
- 114 Presence of nitro gp. in benzene ring:  
0.
- a) Deactivates the ring for S<sub>E</sub> reaction  
b) Activates the ring for S<sub>E</sub> reactions  
c) Renders the ring basic  
d) Deactivates the ring for S<sub>N</sub> reaction
- 114 Which of the following complexes will show geometrical as well as optical isomerism? (en = ethylene diamine)  
1.
- a) [Pt(NH<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub>]                                      b) [Pt(NH<sub>3</sub>)<sub>2</sub>Cl<sub>4</sub>]                                      c) [Pt(en)<sub>3</sub>]<sup>4+</sup>                                      d) [Pt(en)<sub>2</sub>Cl<sub>2</sub>]
- 114 The huge number of organic compounds is due to the fact that  
2.
- a) Tetravalency of carbon                                      b) Carbon possesses property of catenation  
c) Carbon compounds exhibits polymerisation                                      d) Both (b) and (c)
- 114 When nitrobenzene is treated with Br<sub>2</sub> in presence of FeBr<sub>3</sub>, the major product formed is *m*-bromonitrobenzene. Statements which are related to obtain the *m*-isomer are:  
3.
- a) The relative electron density on *meta* carbon is more than that of *ortho* and *para* positions  
b) Loss of aromaticity when Br<sup>+</sup> attacks at the *ortho* and *para* positions and not at *meta* position  
c) Easier loss of H<sup>+</sup> to regain aromaticity from the *meta* position than from *ortho* and *para* positions  
d) None of the above
- 114 Which one of the following compounds when dissolved in water, gives a solution with pH more than 7?  
4.
- a) C<sub>6</sub>H<sub>5</sub>NH<sub>2</sub>                                      b) C<sub>6</sub>H<sub>5</sub>OH                                      c) C<sub>2</sub>H<sub>5</sub>OH                                      d) CH<sub>3</sub>COCH<sub>3</sub>
- 114 Formula of ferrocene is:  
5.
- a) [Fe(CN)<sub>6</sub>]<sup>4-</sup>                                      b) [Fe(CN)<sub>6</sub>]<sup>3+</sup>                                      c) [Fe(CO)<sub>5</sub>]                                      d) [(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>Fe]
- 114 What is the EAN of nickel in Ni(CO)<sub>4</sub>?  
6.
- a) 38                                      b) 30                                      c) 36                                      d) 32
- 114 One mole of the complex compound Co(NH<sub>3</sub>)<sub>5</sub>Cl<sub>3</sub>, gives 3 moles of ions on dissolution in water. One mole of the same complex reacts with two moles of AgNO<sub>3</sub> solution to yield two moles of AgCl(s). The structure of the complex is  
7.
- a) [Co(NH<sub>3</sub>)<sub>5</sub>Cl]Cl<sub>2</sub>                                      b) [Co(NH<sub>3</sub>)<sub>3</sub>Cl<sub>2</sub>].2NH<sub>3</sub>  
c) [Co(NH<sub>3</sub>)<sub>4</sub>Cl<sub>2</sub>]Cl.NH<sub>3</sub>                                      d) [Co(NH<sub>3</sub>)<sub>4</sub>Cl]Cl<sub>2</sub>.NH<sub>3</sub>
- 114 Which one of the following has largest number of isomers?  
8. (R=alkyl group, en=ethylenediamine)
- a) [Ru(NH<sub>3</sub>)<sub>4</sub>Cl<sub>2</sub>]<sup>+</sup>                                      b) [Co(NH<sub>3</sub>)<sub>5</sub>Cl]<sup>2+</sup>                                      c) [Ir(PR<sub>3</sub>)<sub>2</sub>H(CO)]<sup>2+</sup>                                      d) [Co(en)<sub>2</sub>Cl<sub>2</sub>]<sup>+</sup>
- 114 Which complex is likely to show optical activity?  
9.
- a) *Trans*-[Co(NH<sub>3</sub>)<sub>4</sub>Cl<sub>2</sub>]<sup>+</sup>.  
b) [Cr(H<sub>2</sub>O)<sub>6</sub>]<sup>3+</sup>  
c) *Cis*-[Co(NH<sub>3</sub>)<sub>2</sub>(en)<sub>2</sub>]<sup>3+</sup>  
d) *Trans*-[Co(NH<sub>3</sub>)<sub>2</sub>(en)<sub>2</sub>]<sup>3+</sup>
- 115 A square planar complex is formed by hybridization of which atomic orbitals?  
0.

a)  $s, p_x, p_y, d_{yz}$

b)  $s, p_x, p_y, d_{x^2-y^2}$

c)  $s, p_x, p_y, d_{z^2}$

d)  $s, p_x, p_y, d_{xy}$

115 The IUPAC name of the compound

1.  $\text{CH}_2 - \text{CH} - \text{CH} - \text{CH}_2\text{COCl}$  is



a) 1, 2, 3, 4-butanetetrachlorocarbonyl

b) 1, 2, 3, 4-butanetetrachloroformyl

c) 1, 2, 4-butanetricarboxylic acid

d) None of the above

115 Nitrobenzene can be prepared from benzene by using a mixture of conc.  $\text{HNO}_3$  and conc.  $\text{H}_2\text{SO}_4$ . In the

2. nitrating mixture  $\text{HNO}_3$  acts as a:

a) Base

b) Acid

c) Reducing agent

d) Catalyst

115 In the compound lithium tetrahydroaluminate, the ligand is

3.

a) H

b)  $\text{H}^+$

c)  $\text{H}^-$

d) None of these

SMART ACHIEVERS LEARNING PVT. LTD.

# COORDINATION COMPOUNDS

## CHEMISTRY

### : ANSWER KEY :


1)	b	2)	b	3)	c	4)	c	177)	c	178)	b	179)	a	180)	b
5)	b	6)	b	7)	c	8)	a	181)	c	182)	b	183)	a	184)	b
9)	c	10)	b	11)	c	12)	c	185)	b	186)	d	187)	c	188)	d
13)	d	14)	d	15)	c	16)	c	189)	d	190)	b	191)	b	192)	a
17)	b	18)	d	19)	b	20)	b	193)	d	194)	d	195)	a	196)	d
21)	a	22)	c	23)	d	24)	d	197)	b	198)	b	199)	c	200)	c
25)	d	26)	d	27)	d	28)	d	201)	b	202)	b	203)	d	204)	a
29)	b	30)	a	31)	d	32)	d	205)	b	206)	d	207)	d	208)	c
33)	d	34)	c	35)	a	36)	b	209)	b	210)	c	211)	c	212)	b
37)	d	38)	c	39)	c	40)	d	213)	c	214)	c	215)	b	216)	a
41)	a	42)	c	43)	c	44)	d	217)	a	218)	d	219)	b	220)	a
45)	b	46)	d	47)	b	48)	d	221)	d	222)	c	223)	c	224)	d
49)	b	50)	d	51)	d	52)	d	225)	a	226)	b	227)	c	228)	c
53)	a	54)	b	55)	b	56)	c	229)	b	230)	d	231)	c	232)	a
57)	a	58)	c	59)	a	60)	b	233)	b	234)	b	235)	d	236)	d
61)	c	62)	c	63)	a	64)	d	237)	b	238)	d	239)	a	240)	b
65)	c	66)	d	67)	c	68)	a	241)	a	242)	a	243)	b	244)	b
69)	d	70)	d	71)	c	72)	a	245)	a	246)	c	247)	b	248)	a
73)	a	74)	b	75)	d	76)	b	249)	c	250)	d	251)	d	252)	c
77)	c	78)	b	79)	c	80)	d	253)	c	254)	a	255)	a	256)	b
81)	d	82)	d	83)	c	84)	d	257)	b	258)	c	259)	d	260)	d
85)	b	86)	b	87)	c	88)	c	261)	d	262)	d	263)	c	264)	a
89)	d	90)	a	91)	c	92)	a	265)	b	266)	d	267)	b	268)	b
93)	c	94)	b	95)	a	96)	c	269)	b	270)	c	271)	d	272)	a
97)	b	98)	c	99)	b	100)	c	273)	d	274)	b	275)	c	276)	d
101)	d	102)	c	103)	d	104)	b	277)	d	278)	c	279)	c	280)	b
105)	d	106)	a	107)	b	108)	c	281)	a	282)	c	283)	b	284)	b
109)	a	110)	a	111)	b	112)	c	285)	b	286)	c	287)	b	288)	d
113)	b	114)	a	115)	d	116)	c	289)	c	290)	b	291)	d	292)	b
117)	c	118)	a	119)	b	120)	b	293)	b	294)	c	295)	c	296)	a
121)	a	122)	d	123)	d	124)	d	297)	c	298)	b	299)	d	300)	b
125)	d	126)	b	127)	a	128)	d	301)	a	302)	b	303)	b	304)	b
129)	b	130)	c	131)	d	132)	b	305)	a	306)	b	307)	b	308)	b
133)	c	134)	d	135)	d	136)	d	309)	b	310)	c	311)	b	312)	c
137)	c	138)	d	139)	b	140)	a	313)	c	314)	a	315)	a	316)	c
141)	b	142)	a	143)	a	144)	c	317)	b	318)	b	319)	d	320)	d
145)	a	146)	d	147)	d	148)	a	321)	b	322)	d	323)	d	324)	b
149)	a	150)	c	151)	a	152)	b	325)	d	326)	c	327)	b	328)	d
153)	b	154)	c	155)	a	156)	b	329)	b	330)	d	331)	d	332)	a
157)	d	158)	b	159)	b	160)	c	333)	d	334)	c	335)	b	336)	d
161)	b	162)	c	163)	d	164)	a	337)	d	338)	d	339)	b	340)	d
165)	a	166)	a	167)	b	168)	b	341)	b	342)	c	343)	a	344)	c
169)	b	170)	b	171)	c	172)	b	345)	b	346)	c	347)	a	348)	b
173)	c	174)	a	175)	c	176)	c	349)	c	350)	b	351)	a	352)	b

353) d	354) b	355) b	356) d	557) a	558) d	559) c	560) c
357) b	358) d	359) d	360) d	561) a	562) b	563) c	564) a
361) d	362) b	363) b	364) a	565) c	566) c	567) c	568) a
365) a	366) a	367) c	368) d	569) c	570) c	571) b	572) d
369) a	370) d	371) c	372) a	573) b	574) c	575) d	576) a
373) d	374) d	375) c	376) c	577) a	578) b	579) b	580) b
377) c	378) a	379) d	380) c	581) d	582) b	583) c	584) d
381) d	382) a	383) d	384) c	585) b	586) c	587) b	588) d
385) d	386) c	387) a	388) c	589) a	590) c	591) b	592) c
389) c	390) c	391) b	392) a	593) c	594) d	595) b	596) b
393) c	394) a	395) c	396) d	597) c	598) a	599) b	600) c
397) a	398) d	399) b	400) d	601) a	602) b	603) d	604) b
401) a	402) b	403) c	404) a	605) d	606) d	607) d	608) b
405) b	406) d	407) b	408) b	609) c	610) d	611) b	612) b
409) a	410) b	411) c	412) d	613) a	614) c	615) a	616) a
413) b	414) a	415) a	416) b	617) d	618) c	619) c	620) c
417) d	418) c	419) c	420) c	621) c	622) c	623) a	624) c
421) c	422) a	423) c	424) b	625) a	626) b	627) d	628) c
425) d	426) c	427) a	428) b	629) a	630) c	631) a	632) b
429) d	430) d	431) a	432) a	633) c	634) b	635) a	636) a
433) c	434) c	435) a	436) b	637) c	638) c	639) c	640) b
437) a	438) b	439) a	440) b	641) d	642) b	643) a	644) c
441) b	442) c	443) b	444) d	645) c	646) a	647) c	648) d
445) a	446) a	447) a	448) a	649) a	650) c	651) a	652) d
449) b	450) d	451) b	452) c	653) d	654) a	655) d	656) d
453) c	454) d	455) a	456) d	657) b	658) a	659) b	660) c
457) a	458) a	459) c	460) c	661) d	662) c	663) b	664) b
461) c	462) b	463) b	464) d	665) c	666) c	667) c	668) c
465) a	466) b	467) a	468) d	669) d	670) a	671) c	672) a
469) c	470) c	471) c	472) d	673) a	674) b	675) b	676) a
473) d	474) c	475) b	476) a	677) c	678) c	679) c	680) b
477) b	478) b	479) c	480) c	681) a	682) a	683) c	684) b
481) b	482) a	483) a	484) c	685) c	686) d	687) b	688) b
485) a	486) a	487) d	488) c	689) b	690) b	691) a	692) c
489) b	490) a	491) c	492) a	693) a	694) a	695) d	696) a
493) d	494) d	495) a	496) a	697) a	698) b	699) b	700) d
497) c	498) c	499) a	500) d	701) d	702) c	703) b	704) a
501) a	502) d	503) b	504) d	705) d	706) d	707) c	708) d
505) b	506) d	507) a	508) d	709) b	710) b	711) b	712) c
509) b	510) d	511) b	512) a	713) a	714) b	715) d	716) d
513) d	514) c	515) b	516) a	717) c	718) b	719) c	720) c
517) a	518) c	519) a	520) a	721) b	722) c	723) c	724) b
521) b	522) c	523) d	524) a	725) a	726) c	727) c	728) b
525) a	526) d	527) d	528) d	729) b	730) c	731) a	732) c
529) d	530) c	531) a	532) b	733) b	734) a	735) a	736) b
533) b	534) a	535) b	536) d	737) b	738) a	739) b	740) d
537) a	538) a	539) b	540) c	741) b	742) d	743) a	744) d
541) c	542) d	543) a	544) b	745) d	746) d	747) b	748) c
545) d	546) c	547) d	548) c	749) c	750) c	751) b	752) a
549) a	550) b	551) b	552) b	753) b	754) b	755) d	756) c
553) d	554) b	555) a	556) d	757) b	758) b	759) b	760) b

761) d	762) b	763) a	764) d	961) c	962) c	963) a	964) a
765) c	766) a	767) b	768) c	965) c	966) d	967) d	968) d
769) b	770) d	771) a	772) b	969) b	970) a	971) b	972) a
773) a	774) d	775) d	776) b	973) b	974) c	975) d	976) c
777) b	778) c	779) b	780) d	977) a	978) d	979) b	980) d
781) d	782) a	783) c	784) d	981) a	982) b	983) b	984) b
785) c	786) b	787) a	788) c	985) d	986) c	987) d	988) a
789) b	790) c	791) c	792) c	989) c	990) d	991) a	992) a
793) b	794) b	795) c	796) d	993) d	994) b	995) c	996) c
797) b	798) d	799) b	800) d	997) a	998) b	999) c	1000) c
801) c	802) b	803) c	804) a	1001) c	1002) c	1003) c	1004) b
805) c	806) d	807) a	808) c	1005) b	1006) b	1007) c	1008) c
809) b	810) d	811) d	812) b	1009) c	1010) b	1011) a	1012) d
813) b	814) b	815) c	816) b	1013) b	1014) a	1015) d	1016) d
817) a	818) d	819) c	820) a	1017) a	1018) b	1019) a	1020) a
821) b	822) c	823) c	824) a	1021) a	1022) a	1023) c	1024) d
825) a	826) d	827) c	828) d	1025) a	1026) d	1027) a	1028) d
829) d	830) b	831) a	832) b	1029) c	1030) a	1031) b	1032) d
833) a	834) c	835) b	836) d	1033) a	1034) a	1035) a	1036) c
837) d	838) a	839) d	840) c	1037) a	1038) d	1039) b	1040) a
841) c	842) b	843) b	844) c	1041) c	1042) a	1043) d	1044) a
845) b	846) b	847) a	848) d	1045) d	1046) b	1047) d	1048) c
849) a	850) b	851) b	852) c	1049) d	1050) b	1051) c	1052) a
853) c	854) b	855) a	856) a	1053) c	1054) c	1055) b	1056) a
857) d	858) a	859) b	860) b	1057) d	1058) b	1059) d	1060) a
861) a	862) c	863) d	864) b	1061) d	1062) d	1063) a	1064) d
865) a	866) d	867) d	868) c	1065) b	1066) a	1067) a	1068) c
869) b	870) d	871) c	872) d	1069) c	1070) d	1071) d	1072) c
873) c	874) a	875) b	876) b	1073) d	1074) a	1075) d	1076) a
877) b	878) a	879) b	880) d	1077) d	1078) a	1079) a	1080) c
881) a	882) d	883) b	884) b	1081) c	1082) d	1083) d	1084) a
885) b	886) c	887) a	888) d	1085) d	1086) d	1087) b	1088) d
889) b	890) d	891) a	892) a	1089) c	1090) a	1091) c	1092) c
893) a	894) b	895) b	896) a	1093) a	1094) c	1095) b	1096) d
897) a	898) b	899) b	900) d	1097) c	1098) c	1099) b	1100) d
901) c	902) d	903) a	904) a	1101) a	1102) a	1103) d	1104) c
905) b	906) c	907) c	908) c	1105) c	1106) c	1107) c	1108) c
909) a	910) d	911) d	912) c	1109) a	1110) b	1111) d	1112) a
913) c	914) c	915) b	916) b	1113) a	1114) b	1115) b	1116) a
917) a	918) d	919) b	920) d	1117) c	1118) d	1119) a	1120) a
921) a	922) c	923) d	924) b	1121) d	1122) c	1123) b	1124) c
925) a	926) c	927) b	928) d	1125) c	1126) c	1127) c	1128) d
929) b	930) d	931) d	932) d	1129) d	1130) c	1131) c	1132) b
933) a	934) c	935) c	936) b	1133) a	1134) c	1135) a	1136) d
937) d	938) a	939) c	940) b	1137) d	1138) d	1139) a	1140) a
941) d	942) d	943) b	944) c	1141) d	1142) d	1143) a	1144) a
945) b	946) a	947) a	948) a	1145) d	1146) c	1147) a	1148) d
949) b	950) d	951) b	952) c	1149) c	1150) b	1151) a	1152) a
953) b	954) a	955) a	956) c	1153) c			
957) a	958) a	959) d	960) c				




$S_E$  reactions




13 (d)

In both  $[\text{Co}(\text{NH}_3)_6]^{3+}$  and  $[\text{CoF}_6]^{3+}$ , Co is present as  $\text{Co}^{3+}$ .


Thus, the electronic configuration of Co is



In case of  $[\text{Co}(\text{NH}_3)_6]^{3+}$ ,  $\text{NH}_3$  is a strong field ligand, so pairing of electrons in  $3d$ -orbital takes place.



In  $[\text{CoF}_6]^{3+}$ , F is a weak field ligand, thus doesn't cause pairing. Hence,



18 (d)

It is a test for primary amines. No doubt 2,4-dimethylaniline is also primary amine but it does not give test due to steric hindrance.

19 (b)

$\text{CN}^-$  is strong field ligand because it is an example of pseudohalide. Pseudohalide ions are stronger coordinating ligands and they have the ability to form  $\sigma$  bond and  $\pi$ -bond.

20 (b)

Higher the charge and smaller the size of ligand, more stable is the complex formed

21 (a)

Trinitrobenzene is an explosive compound formed during nitration of  $\text{C}_6\text{H}_6$  with fuming  $\text{HNO}_3$ .

22 (c)

A ligand is a species that is capable of donating an electron pair(s) to the central metal ion. The substances which are capable of donating an electron pair are called Lewis base, so a ligand is also a Lewis base.

23 (d)


In  $\text{Ni}(\text{CO})_4$ , Ni is in zero oxidation state. It has tetrahedral geometry but is diamagnetic. In

$[\text{Ni}(\text{CN})_4]^{2-}$ , Ni is in +2 oxidation state. It is  $dsp^2$  hybridised and have square planar shape. The compound is diamagnetic.

24 (d)

$[\text{Co}(\text{CN})_6]^{3-}$  has  $d^2sp^3$ -hybridisation and six  $d$ -electrons are paired due to strong field ligand. Thus no unpaired electron.

25 (d)




Ether reacts with acid to give protonated ether. The next step involves nucleophilic attack by halide ion with the displacement of weakly basic alcohol molecule.

26 (d)

Octahedral complex should have six hybridized orbitals.

27 (d)



Kolbe-Schmidt's reaction.

29 (b)

The pair of electron present with nitrogen will not be available to be donated as  $\text{H}^+$  will consume that one.

30 (a)

It provides maximum number of ions (five) on ionization.

31 (d)


Follow Vorlander's rule.

32 (d)

Organometallic compounds are those in which metal is linked directly with carbon.  $\text{CH}_3\text{Li}$ , methyl lithium due to the presence of metal-carbon bond, is an organometallic compound.

33 (d)

The directive influence order is:



34 (c)

Hybridisation	Shape
---------------	-------

$dsp^2$	Square planar
$sp^3$	Tetrahedral
$sp^2$	Trigonal planar

Hence, in tetrahedral complexes metal atom is  $sp^3$  hybridised.

36 (b)

The number of ligands attached to the central metal ion is called the coordination number.

So, coordination numbers of Fe in  $[Fe(CN)_6]^{4-}$ ,  $[Fe(CN)_6]^{3-}$  and  $[FeCl_4]^-$  are 6, 6 and 4 respectively.

37 (d)

Tautomers may or may not be metamers

38 (c)

EAN = (Atomic number - O.S + 2 × C.N.)

Hence, EAN of Ni in  $[Ni(CN)_4]^{2-}$  =  $(28 - 2 + 2 \times 4) = 34$

39 (c)

Electron repelling nature of methoxy gp. facilitate the protonation of alcohol.


40 (d)

$[Ni(Cl)_4]^{2-}$  oxidation state of Ni is +2

So, configuration of  $Ni^{2+}$  =

$1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^8$

In  $Ni^{2+}$



In  $[NiCl_4]^{2-}$



$sp^3$  - hybridisation

Thus, due to  $sp^3$ -hybridisation of  $Ni^{2+}$  in  $[NiCl_4]^{2-}$ , the shape of  $[NiCl_4]^{2-}$  is tetrahedral.

41 (a)



This is Sandmeyer's reaction.

42 (c)

*p*-nitrophenols are more acidic.

43 (c)

Benzoylacetato beryllium exhibit optical isomerism as follows



bis (benzoylacetato) beryllium (II) complex

44 (d)

$Cl^-$  is a weak ligand but  $Cl^-$  cause the pairing of electron with large  $Pt^{2+}$  and consequently give  $dsp^2$  hybridisation and square planar geometry.

45 (b)

It is a double salt;

$FeSO_4 \cdot (NH_4)_2SO_4 \cdot 6H_2O$



46 (d)

Potassium ferrocyanide  $K_4[Fe(CN)_6]$  will ionize as



So, it will give five ions in solution

47 (b)

*cis*-platin is not a organometallic compound because it has no carbon- metal bonding

48 (d)

Follow mechanism of Reimer-Tiemann reaction.

49 (b)

When  $n$  = even number then for two identical ends, number of geometrical isomers

$$= 2^{n-1} + 2^{n/2-1}$$

$$= 2^1 + 2^0$$

$$= 3$$

50 (d)

The characteristics of coordination number.


51 (d)

Aliphatic amines are more basic than aromatic amines as the later are more stabilised due to resonance.


52 (d)

Aromatic hydrocarbons are called arenes with general formula  $C_nH_{2n-6y}$ , where  $n < 6$  and  $y$  is no. of cyclic rings. Benzene has one ring and  $n = 6$ , i. e., no. of carbon atoms. Thus, general formula is  $C_6H_6$ . All other aromatic hydrocarbons are derivative of benzene.


- 53 (a)  $\text{Al}(\text{OC}_2\text{H}_5)_3$  doesn't have metal-carbon bond. (i.e., it is not an example of organometallic compound).




- 54 (b) In  $[\text{Co}(\text{en})_2\text{Cl}_2]$ , four isomers are possible, two geometrical isomers and two optical isomers.



Now, *cis*-isomer also show optical isomerism. *Cis* isomer exists in enantiomeric form as it is unsymmetrical.



- 55 (b) A carbon atom which is attached by four different group is called an asymmetric carbon atom or chiral centre.  $\text{HOOC}(\text{CHOH})_2\text{COOH}$  has two asymmetric carbon atom



- 56 (c) Each  $\pi$ -electron is delocalised on each C-atom.


- 57 (a) An orange-red dye is formed with  $\text{C}_6\text{H}_5\text{NH}_2$ .

- 59 (a) Thiophene reacts more readily with  $\text{H}_2\text{SO}_4$  than  $\text{C}_6\text{H}_6$  giving thiophene sulphonic acid which is water soluble and thus, can be separated from  $\text{C}_6\text{H}_6$ . This can not be made by fractional

distillation because thiophene and  $\text{C}_6\text{H}_6$  both have nearly same b.p.

- 60 (b) As cobalt is present as  $\text{Co}^{3+}$  and coordination number of cobalt is 6, the molecular formula of compound should be  $\text{CoCl}_3 \cdot y\text{NH}_3$ . Now, as it gives a total of three ions when dissolved in water, its structural formula must be  $[\text{CoCl}(\text{NH}_3)_5]\text{Cl}_2$   
 $[\text{CoCl}(\text{NH}_3)_5]\text{Cl}_2 \rightleftharpoons [\text{CoCl}(\text{NH}_3)_5]^{2+} + 2\text{Cl}^-$   
 Thus, only one  $\text{Cl}^-$  ion is satisfying both primary and secondary valency of  $\text{Co}^{3+}$  in this compound.

- 61 (c) The structure of alcohol is



2,4-dimethyl-3-(1-methyl) ethyl pentan-3-ol

- 62 (c) The transition metal cations during complex formation show *d-d* transition to give coloured ions.

- 63 (a)  $-\text{CH}_3$  gp. Shows +ve inductive effect and  $-\text{OH}$  gp. shows resonance effect which increases the electron density on  $\text{C}_6\text{H}_6$  ring.

- 64 (d) It produces least number of ions in solution.


- 66 (d) The process is known as aromatisation or cyclization.

- 67 (c)
- $$\begin{array}{c} \text{CH}_3 \\ | \\ \text{CH}_3 - \text{C} - \text{CH}_3 \\ | \\ \text{CH}_3 \end{array}$$
- neo*-pentane

The structure shows that all the hydrogen atoms are attached to primary C-atoms hence these are primary hydrogens

- 68 (a) Follow IUPAC rules.

- 69 (d)



has no  $\alpha$ -hydrogen. Hence, it will not show tautomerism

70 (d)

Both  $\text{CN}^-$  and  $\text{NO}_2^-$  are strong field ligands.

71 (c)

Prussian blue is

$\text{Fe}_4^{\text{III}}[\text{Fe}^{\text{II}}(\text{CN})_6]_3$  or  $\text{M}^{\text{I}}\text{Fe}^{\text{III}}[\text{Fe}^{\text{II}}(\text{CN})_6]$ , where M<sup>I</sup> is Na, K, Rb, Li, Cs.

73 (a)

$\text{Co}^{3+}$ ,  $\text{Fe}^{3+}$  and  $\text{Cr}^{3+}$  have 6d-electrons, 5d-electrons and 3d-electrons respectively.  $\text{Mn}^{7+}$  has no d-electron.

74 (b)

All complexes of Co(III) have six ligands or coordination number of six and thus, are octahedral in shape.

75 (d)

$\text{NH}_3$  is weak as well as strong field ligand.

77 (c)

$[\text{Pt}(\text{NH}_3)_3\text{Br}(\text{NO}_2)\text{Cl}]\text{Cl}$

Triamminebromochloronitro platinum (IV) chloride.

78 (b)

Both the carbon attached to O are part of aromatic system.

79 (c)


Phenol is weak acid.

80 (d)

$[\text{EDTA}]^{4-}$  is a hexadentate ligand because it donates six pairs of electrons to central metal atom in a complex.

82 (d)

Ferrocene of bi-(cyclopentadienyl) iron is an orange-crystalline solid. It is  $\text{Fe}(\eta^5\text{-C}_5\text{H}_5)_2$ . The structure of ferrocene is regarded as sandwich structure, in which the iron atom is sandwiched between two  $\text{C}_5\text{H}_5$  organic rings. The planes of the rings are parallel so that all the carbon atoms are at the same distance from the iron atom. It is a  $\pi$ -bonded complex. Its structure is as



83 (c)

$[\text{Cu}(\text{NH}_3)_4]\text{SO}_4$

Oxidation number of

$$\text{Cu} \Rightarrow x + 4 \times 0 - 2 = 0$$

$$x - 2 = 0$$

$$x = +2$$

O.N of Cu = +2

O.N of pt in  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$

$$x + 2 \times 0 + 2 \times -1 = 0$$

$$x - 2 = 0$$

$$x = +2$$

O.N of Ni in  $[\text{Ni}(\text{CO})_4]$

$$x + 4 \times 0 = 0$$

$$x = 0$$

O.N of Fe in  $\text{K}_3[\text{Fe}(\text{CN})_6]$

$$3 \times (+1) + x + 6 \times -1 = 0$$





$$3 + x - 6 = 0$$

$$x = +3$$

$\therefore [\text{Ni}(\text{CO})_4]$  is zero valent compound.

84 (d)

The compounds given have following structures




Out of these the (a) and (b) contain 5 C-atoms in their longest possible chains hence, these could not be the correct options for 2, 2, 3-trimethylhexane. Out of (c) and (d), the (c) is 2, 2, 5-trimethyl hexane and (d) is 2, 3, 3-trimethyl hexane

85 (b)

Phenoxy benzene is diphenyl ether.

86 (b)

Ziegler-Natta catalyst is an organometallic compound containing titanium. It is  $\text{TiCl}_4$  and  $(\text{C}_2\text{H}_5)_3\text{Al}$ . It is used in the preparation of polyethylene.




polyethylene

87 (c)  $\text{Al}_2(\text{C}_2\text{H}_5)_6 + \text{TiCl}_4$  is Zeigler Natta catalyst.


88 (c) Transition metals have empty or half filled  $d$ -orbitals to accept electron pairs.

89 (d) The number of atom of the ligand that are directly bound to the central metal atom or ion by coordinate bonds is known as the coordinate number of the metal or ion. It is actually the number of chemical bonds which the ligand form with the central metal atom or ion




90 (a) Acyl chlorides or acid amhydrides are used in acylation.

92 (a) Zeise's salt,  $\text{K}[\text{PtCl}_3(\text{C}_2\text{H}_4)]$  is a  $\pi$ -bonded organometallic compound. Its structure is as




95 (a) Follow IUPAC rules.


96 (c) Since the complexes  $[\text{PtCl}_2(\text{NH}_3)_4]\text{Br}_2$  and  $[\text{PtBr}_2(\text{NH}_3)_4]\text{Cl}_2$  have the same molecular formula but on ionisation they give different ions, they exhibit ionisation isomerism.




97 (b)  $\text{Ni}^{2+} + 4\text{CN}^- \rightarrow [\text{Ni}(\text{CN})_4]^{2-}$   
Here  $\text{Ni}^{2+}$  has  $d^8$ -configuration with  $\text{CN}^-$  as strong ligand.




$d^8$ -configuration in strong ligand field gives  $dsp^2$  hybridisation, hence square planar geometry.



Here  $\text{Ni}^{2+}$  has  $d^8$ -configuration with  $\text{Cl}^-$  as weak ligand.



$d^8$ -configuration in weak ligand field gives  $sp^3$  hybridisation, hence tetrahedral geometry.  $\text{Ni}^{2+}$  with  $\text{H}_2\text{O}$  forms  $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$  complex and  $\text{H}_2\text{O}$  is a weak ligand.



Therefore,  $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$  has octahedral geometry.

98 (c) Benzene ring is activates for  $S_E$  reaction by the  $+I$  effect as well as hyperconjugation of  $\text{CH}_3$  group  $-\text{Cl}$  deactivates as  $-I$  effect predominates over  $+M$  effect.  $-\text{NO}_2$  group deactivates ring by  $-I$  effect and  $-M$  effect.

100 (c) Alcohols are neutral.

101 (d)  $-\text{OH}$  is  $o$ - and  $p$ -directing gp.

102 (c)  $[\text{Fe}(\eta^5 - \text{C}_2\text{H}_5)_2]$  is the organometallic compound which has  $\sigma$  and  $\pi$  bonds present

103 (d) BHC is  $\text{C}_6\text{H}_6\text{Cl}_6$  a saturated cyclic molecule.

104 (b) The complexes can be written as follows  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$ ,  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$ ,  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$  Hence, no. of primary valencies are 3, 2 and 1 respectively


105 (d)  $[\text{Cr}(\text{NH}_3)_5\text{NO}_2]\text{Cl}_2$  compound shows linkage isomerism because it has  $\text{NO}_2$  group which is ambidentate ligand. It can be linked *via* N atom ( $-\text{NO}_2$ ) or *via* O atom ( $-\text{ONO}$ ) to form two different isomers.

106 (a)

In  $[Sc(H_2O)_6]^{3+}$ ,

Oxidation state of Sc is +3.

Sc (ground state)



$Sc^{3+}$



$\therefore Sc^{3+}$  has no unpaired electron.


$\therefore [Sc(H_2O)_6]^{3+}$  is diamagnetic and colourless.

107 (b)


$[MA_5B]$  due to absence of symmetry of 'B' ligand cannot exist in the form of *cis-trans* isomer.

108 (c)

Out of the 3 functional groups attached




will be the principal functional group and rest as the substituents



2-amino-3-hydroxy propanoic acid

109 (a)

A transition metal complex absorbs visible light only when it has unpaired electron.  $Ni^{2+}$  in strong field ligand has configuration as



110 (a)

The directive influence order and tendency to release electron for *o*- and *p*-directing group is,



111 (b)

(a) In  $MnO_2$ ,  $FeCl_3$  oxidation states of Mn and Fe are +4 and +3 respectively.

(b) In  $(MnO_4)^-$ ,  $CrO_2Cl_2$  oxidation states of Mn and Cr are +7 and +6 respectively.


(c) In  $[Fe(CN)_6]^{3-}$ ,  $[Co(CN)_3]$  oxidation states of Fe and Co are +3 and +3 respectively.

(d)  $[NiCl_4]^{2-}$ ,  $[CoCl_4]^-$  oxidation states of Ni and Co are +2 and +3 respectively.

112 (c)


$[M(abcd)]$  complex is square planar so will have three geometrical isomers.

127 (a)



113 (b)

This is carbylamines reaction.



114 (a)

An experimental fact depending upon the ability of the ligand to cause crystal field splitting (*i.e.*, strength of ligand).

115 (d)

$K_3[Fe(CN)_5NO]$

Potassium pentacyanonitrosyl ferrate (II).

116 (c)

The *d-d* excitation is responsible for colour of  $Ti(H_2O)_6^{3+}$  which has one unpaired electron.

117 (c)

The oxidation number of Fe in  $K_4[Fe(CN)_6]$  is +2.

119 (b)


Both Ag and Au are extracted by complex formation method.

120 (b)

$EAN = 24 - 3 + 2 \times (6) = 33$ .

121 (a)

According to Werner's theory, only those ions are precipitated which are attached to the metal atoms with ionic bonds and are present outside the coordination sphere.



122 (d)

It is a fact.

123 (d)

In acidic solution, proton coordinate with ammonia to form  $NH_4^+$ .  $NH_4^+$  does not act as ligand because nitrogen atom has no lone pair of electrons which it can donate to metal atom


124 (d)

Disubstituted cyclic compounds and disubstituted alkenes show geometrical isomerism

125 (d)


$Ag(NH_3)_2^+$  has *sp*-hybridization and linear complex.

The replacement of Cl is due to the formation of stable benzyl carbocation. Alternatively Cl is present in side chain and thus replaced whereas Br is attached in benzene nucleus.




128 (d)


The electronic configuration of Ni in  $[\text{Ni}(\text{CN})_4]^{2-}$ ,  $[\text{Ni}(\text{Cl})_4]^{2-}$  and  $\text{Ni}(\text{CO})_4$  are:  
 $\text{Ni}^{2+}$  in  $[\text{Ni}(\text{CN})_4]^{2-}$ :



$\text{Ni}^{2+}$  in  $[\text{Ni}(\text{Cl})_4]^{2-}$ :



Ni in  $[\text{Ni}(\text{CO})_4]$ :




129 (b)

Replacement of  $\text{N}_2\text{Cl}$  by halogen atom of  $\text{CuX}$  -  $\text{HX}$  from benzene diazonium chloride is called Sandmeyer's reaction.

134 (d)


(i)  $-\text{NO}_2$  can show linkage



130 (c)

Optical isomerism is shown by the type  $[\text{M}(\text{AA})\text{X}_2\text{Y}_2]$ ,  $[\text{M}(\text{AA})_3]$ ,  $[\text{M}(\text{AA})_2\text{X}_2]$


131 (d)



1,2-dimethyl cyclobut-1-ene

132 (b)





(iii) Also  $[\text{Co}(\text{NH}_3)_4(\text{NO}_2)_2]\text{Cl}$  has its ionisation isomer as  $[\text{Co}(\text{NH}_3)_4\text{NO}_2\text{Cl}]\text{NO}_2$ .

135 (d)

Complex	Hybridization
$[\text{Ni}(\text{CO})_4]$	$sp^3$
$[\text{Ni}(\text{CO})_4]^{2-}$	$dsp^2$
$[\text{CoF}_6]^{3-}$	$sp^3d^2$
$[\text{Fe}(\text{CN})_6]^{3-}$	$d^2sp^3$

136 (d)


2, 4, 6-trinitrophenol is known as picric acid, an explosive.

137 (c)

It is a fact.


139 (b)

OH gp., an *o*- and *p*-directing group activates ring for reactions. The presence of *m*-directing groups in benzene nucleus simply decreases electron density at *o*- and *p*-, whereas no change in electron density at *m*-position is noticed.



On the contrary *o*- and *p*-directing groups in nucleus increases the electron density at *o*- and *p*-position.

Thus, presence of *o*- and *p*-directing groups provide seats for  $S_E$  reactions or activates the ring, whereas presence of *m*-directing groups does not activate the ring and thus, deactivate the ring for  $S_E$  reactions



140 (a)

If magnetic moment is zero the species should not have unpaired electrons.

141 (b)

Chlorophyll is a complex having Mg-atom.





142 (a)

Primary valency of metal is satisfied only by the anion. It is simply ionic valency. While secondary valency is satisfied by ligands (which can give a lone pair of electron). The ligands satisfying secondary valency, are always written in coordination sphere. This concept was given by Werner.

In  $\text{K}_3[\text{Fe}(\text{CN})_6]$ , the  $\text{CN}^-$  ions satisfy both the primary as well as secondary valency of  $\text{Fe}^{3+}$  ion.

143 (a)

The following isomers the alkene have




145 (a)

According to postulates of Werner's theory for coordination compounds, metal atoms exhibit two types of valencies *i.e.*, primary valency and secondary valency. The primary valency is ionisable whereas the secondary valency is non-ionisable.


146 (d)

$[\text{Fe}(\text{CN})_6]^{4-}$  is **diamagnetic** ( $\text{Fe}^{2+}$  has  $3d^6$  configuration and the 6 electron pairs up in three  $d$ -orbitals followed by  $d^2sp^3$ -hybridisation).  $[\text{Cr}(\text{NH}_3)_6]^{3+}$  is **paramagnetic** ( $\text{Cr}^{3+}$  has  $3d^3$  configuration. Hybridisation is  $d^2sp^3$ . Due to 3 unpaired electrons it is **paramagnetic**)



25):  $[\text{Ar}]^{18}4s^1, 3d^5$ .

The one  $4s$ -electron pairs up with five  $3d$ -electrons in three  $d$ -orbitals. This is followed by  $d^2sp^3$ -hybridisation to give octahedral complex. No unpaired electron and hence complex is **diamagnetic**.



26):  $[\text{Ar}]^{18}4s^2, 3d^6$ .


The six electrons in  $d$ -subshell pairs up in three  $d$ -orbitals. This is followed by  $d^2sp^3$ -hybridisation to give octahedral geometry with one vacant hybridised orbital. The resulting shape of the complex is square based pyramid. As there is no unpaired electron, the complex is **diamagnetic**.

147 (d)

A modified or extended Friedel-Crafts reaction.

148 (a)

$\text{cis}[\text{Co}(\text{en})_2\text{Cl}_2]\text{Cl}$  is optically active hence, it will give a pair



149 (a)

$\Delta_t$  is roughly  $4/9$  times to  $\Delta_o$ .


150 (c)

Follow IUPAC rules.

151 (a)

Alkanes having less than four carbon atoms in basic chain will not show chain isomerism

152 (b)



3 mole of  $\text{Br}_2$  are needed.

153 (b)

Diamethyl glyoxime forms a colour complex with nickel

154 (c)


$\text{BF}_4^-$  has  $sp^3$ -hybridisation and tetrahedral.

155 (a)

Oxidation state of iron in haemoglobin is +2.

156 (b)

- Geometrical isomers have same structural formula but differ in spatial arrangement of groups.
- Different arrangement of atoms or groups in three dimensional space results in two optical isomers which are image of each other.



Therefore, number of geometrical isomers, optical isomers and total number of isomers are 2, 2 and 3 respectively.

157 (d)

Hetero aromatics show aromatic nature due to  $4n + 2\pi$  electrons.

158 (b)

$\text{CN}^-$  ligand has strong ligand field because of higher value of  $\Delta$ .

159 (b)

% Enantiomeric excess

$$= \frac{\text{observed specific rotation}}{\text{specific rotation of pure enantiomer}} \times 100$$

$$\text{Observed specific rotation} = \frac{3/4}{100} \times (+16^\circ) \times 100 = +12^\circ$$

160 (c)

Follow IUPAC rules.

161 (b)

$[(\text{C}_6\text{H}_5)_3\text{P}]_3\text{RhCl}$  or  $[(\text{Ph}_3\text{P})_3\text{RhCl}]$  is a Wilkinson's catalyst, the most widely used of all catalysts for homogeneous hydrogenation.


162 (c)

Halogens attack double bond of  $\text{C}_6\text{H}_6$  in presence

of light. In absence of light as well as in presence of only  $\text{AlCl}_3$ ,  $\text{S}_E$  reactions are noticed.

163 (d)

$[\text{Pt}(\text{NH}_3)_6]\text{Cl}_4$  complex gives five ions in the solution.



164 (a)

The EAN for Cu in  $[\text{Cu}(\text{NH}_3)_4]^{2+}$  is 35 and not 36, the next inert gas at. No.

165 (a)

$$1 \times 3 + a + 6 \times (-1) = 0, \therefore a = +3$$


166 (a)

In  $\text{NaOC}_2\text{H}_5$ , Na is attached to O-atom.

167 (b)


In  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$ , Mn is present as  $\text{Mn}^{2+}$  or Mn (II), so its electronic configuration

$$= 1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^5$$



In  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$ , the coordination number of Mn is six, but in presence of weak field ligand, there will be no pairing of electrons in  $3d$ . So, it will form high spin complex due to presence of five unpaired electron.

In  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$




170 (b)

Due to aromatic nature;  $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$  is exception and does not burn with sooty flame.

171 (c)

EDTA (Ethylenediaminetetraacetic acid)



hexadentate (6 electron pairs)

that's why for octahedral complex only one EDTA is required.

173 (c)

It is Friedel-Crafts reaction.

174 (a)

Resonance in phenoxide ion makes it more stable. More stable is ion less stable is phenol or more is acidic nature.


175 (c)

Triethylenediamine cobalt(III) chloride is


$[\text{Co}(\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2)_3]\text{Cl}_3$ ;  $\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2$  is bidentate ligand and thus, coordination no. =  $3 \times 2 = 6$ .

177 (c)

CO is a neutral ligand, so the oxidation state of metal in metal carbonyls is always zero.




178 (b)



179 (a)

$[\text{Ni}(\text{CN})_4]^{2-}$  has  $dsp^2$ -hybridization while  $[\text{Ni}(\text{Cl}_4)^{2-}]$  and  $[\text{Ni}(\text{CO})_4]$  have  $sp^3$ -hybridization.

180 (b)




182 (b)

$\text{Cr}^{2+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Fe}^{2+}$  and  $\text{Ni}^{2+}$  have 4, 5, 4 and 2 unpaired electrons respectively.

183 (a)

It is a reason for the fact.

186 (d)



187 (c)

Non-polar part  $\text{C}_6\text{H}_5$  — shows more hydrophobic nature.

189 (d)

All involve  $d^2sp^3$ -hybridization.

191 (b)

Aromatic amines are less basic than aliphatic amines. Also presence of electron attracting group decreases the basic character of aromatic amines.

192 (a)


Follow IUPAC rules.

193 (d)

All are weak field ligands and thus, give high spin complex.

194 (d)

Tartaric acid is



2,3-dihydroxybutane-1,4-dioic acid

195 (a)

$\beta_4$  for  $[\text{ML}_4]^{2-}$  can be written as


$$\beta_4 = \frac{[\text{ML}_4]^{2-}}{[\text{M}^{2+}][\text{L}^-]^4} = 2.5 \times 10^{13}$$

The overall formation equilibrium constant can be written as

$$k = \frac{[\text{ML}_4]^{2-}}{[\text{M}^{2+}][\text{L}^-]^4}$$

$$k = \beta_4 = 2.5 \times 10^{13}$$

196 (d)



Let oxidation state of Cr =  $x$

$$\text{NH}_3 = 0$$

$$\text{Cl} = -1$$

Net charge = +1

$$\therefore [\text{Cr}(\text{NH}_3)_4\text{Cl}_2]^+$$

$$x + 4 \times 0 + 2(-1) = +1$$

$$\therefore x = +3$$

197 (b)

Phenols are acidic; alcohols are neutral.

198 (b)

$$2 \times a + 4 \times (-2) + 2 \times 0 + 2 \times 0 = -2, \therefore a = +3$$

199 (c)

$\text{CH}_3\text{MgI}$  (Grignard reagent) is an organometallic compound due to C—Mg bond.

200 (c)

Effective atomic number = electrons in  $\text{Cr}^{3+}$  + electrons from  $6\text{NH}_3$  ligands.  
 $= 21 + 6 \times 2 = 33$


203 (d)

Hückel rule for aromaticity suggests that an aromatic compound must possess  $(4n + 2)\pi$ -electrons, where  $n = 0, 1, 2, \dots$ , etc., as well as  $\pi$ -electrons cloud should embrace all the carbon atoms of the cyclic systems.

204 (a)




205 (b)



N-bromo-N-chloro ethanamide

206 (d)


It is condensation reaction.



207 (d)

Due to more canonical forms.

208 (c)




209 (b)

Isomeric substances that differ only in the arrangement of carbon atoms forming the base chain are known as chain isomers



and



If the compound with the same molecular formula differ in the position of the same functional group on the identical base chain the compounds are called position isomers

210 (c)

In  $\text{K}_3[\text{Fe}(\text{CN})_6]$ , the ligands are negative which is present in coordination spheres shows a dual behavior. It may satisfied both primary and secondary valencies while, neutral ligand satisfied only secondary valencies


211 (c)

A number of transition metals form polymetallic carbonyls.

212 (b)



Since, Cl is a weak field ligand, it doesn't cause pairing of electron.



Number of unpaired electrons,  $n=2$


$$\mu = \sqrt{n(n+2)} = \sqrt{2(2+2)} \text{ BM} = \sqrt{8} \text{ BM} = 2.82 \text{ BM}$$

213 (c)

Follow IUPAC rules.

214 (c)


The reaction occurs via., electrophilic addition following Markownikoff's rule.,



215 (b)

$\text{C}_6\text{H}_5\text{OH}$  is also called carboic acid.

216 (a)



It is easier to do nucleophilic substitution on alkyl halides than on aryl halides.

217 (a)

Oxidation state of Co in  $\text{K}[\text{Co}(\text{CO})_4]$  is

$$+1 + x + 4 \times 0 = 0$$


$$x = -1 (\text{For co})$$

218 (d)

Bakelite is formed as a result of condensation of HCHO and phenol.

219 (b)

Based on spectrochemical series, ligands arranged in increasing order of crystal field strength are as



222 (c)

Follow mechanism of sulphonation on xylene.

224 (d)

All these are used to explain *o*-, *p*- directing nature of  $-\text{CH}_3$  gp.

225 (a)

The separation of racemic mixture back into *d* and *l* isomers is known as resolution. It can be

done by

(I) mechanical method

(II) bio-chemical method using enzymes

(III) chemical method (salt formation)

227 (c)

Alkanes are not dissolved in  $\text{H}_2\text{SO}_4$ .

228 (c)

Monomeric form of iron carbonyl is  $\text{Fe}(\text{CO})_5$ .

229 (b)

Molecular formula of naphthalene is  $\text{C}_{10}\text{H}_8$ .

230 (d)

It is clear from the chemical formulae that Ag is central metal atom and ligands are 2 ammonia molecule

Hence, compound is  $[\text{Ag}(\text{NH}_3)_2]\text{Cl}$

231 (c)

For  $[Mabcd]$  square planar complex, the number of possible geometrical isomers is three which is obtained by fixing the position of one of the ligands say *a* while the other ligands *b*, *c* and *d* are placed *trans* to it.

232 (a)

Follow exceptions of Vorlander's rule.

233 (b)


Follow IUPAC rules.

234 (b)

Coordination number is equal to total number of ligands in a complex

235 (d)


Chair and boat conformations of cyclohexane differ in energy by 44 kJ/mol



236 (d)

Ligands form coordinate bond with central atom or ion and donate electron pair.

238 (d)




This is Gattermann-Koch reaction -CHO gp. in  $\text{C}_6\text{H}_6$  nucleus.

239 (a)


$[\text{Cr}(\text{NH}_3)_6]^{3+}$  has three unpaired electrons.

Electronic configuration of  $\text{Cr}^{3+}$  in  $\text{Cr}(\text{NH}_3)_6$  is:



×× Electron pair donated by NH<sub>3</sub>.

240 (b)




The above compound has chiral centre  
Hence, it can exhibit optical isomerism while geometrical isomerism is not possible due to presence of identical groups on double bonded carbon atom

241 (a)

It has no unpaired electron.

242 (a)



Meso tartaric acid is optically inactive due to the presence of molecular symmetry. It is optically inactive due to internal compensation, i.e., the effect of one half of the molecule is neutralized by other

243 (b)


Smaller is cation, more is effective nuclear charge, more is the tendency to attract electron pair from ligands.

245 (a)


Presence of *o*-, *p*-directing gp. facilitates the SE reactions.

246 (c)

K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> gives quinol; KMnO<sub>4</sub> gives mesotartaric acid.



247 (b)




The compound is substituted octane, it has branches at carbon-3, carbon-4, carbon-5. The name is

4-sec-butyl-5-ethyl-3-methyl octane

248 (a)


The presence of *m*-directing groups in benzene

nucleus simply decreases electron density at *o*- and *p*-, whereas no change in electron density at *m*-position is noticed.



On the contrary *o*- and *p*-directing groups in nucleus increases the electron density at *o*- and *p*-position.

Thus, presence of *o*- and *p*-directing groups provide seats for S<sub>E</sub> reactions or activates the ring, whereas presence of *m*-directing groups does not activate the ring and thus deactivate the ring for S<sub>E</sub> reactions



249 (c)

CuF<sub>2</sub> is blue coloured crystalline solid.

250 (d)

Different ionization gives different colour.

251 (d)

The complex has coordination number of six which is found in octahedral complex.

252 (c)




BF<sub>3</sub> has incomplete octet and is Lewis acid; it cannot donate electron pair.

253 (c)

Methyl thiomethyl group is inserted at *ortho* position by heating phenol with dimethyl sulphoxide and pyridine -SO<sub>2</sub>/(CH<sub>3</sub>CO)<sub>2</sub>O.



254 (a)

Ni(CN)<sub>4</sub><sup>2-</sup>





Ni(CN)<sub>4</sub><sup>2-</sup>

Ni is in +2 oxidation state.







Oxidation state of Ni is zero



255 (a)

259 (d)

Aspirin is acetyl salicylic acid.



Thus, only aspirin has carboxylic group.

260 (d)

Replacement of H-atom of ring usually takes place following  $S_E$  reaction mechanism.

261 (d)



262 (d)

Because there is direct bonding of metal ion with carbon

263 (c)

$\text{CuCl}$  in  $\text{NH}_4\text{OH}$  absorbs  $\text{CO}$ .

265 (b)

It is a fact.

267 (b)

$-\text{COOH}$  gp. is *meta* directing gp.

268 (b)



269 (b)


4-methyl benzene sulphonic acid is stronger than acetic acid thus it will release acid from sodium acetate.

256 (b)

Phthalein test is characteristics of phenols.

257 (b)

$[\text{Cr}(\text{NH}_3)_6][\text{Co}(\text{CN})_6]$  is isomer to  $[\text{Cr}(\text{CN})_6][\text{Co}(\text{NH}_3)_6]$ , i.e., ligands are partially changes in complex anion and complex cation.



Compound has one chiral carbon atom thus, it has two geometrical (*cis* and *trans*) and two optical isomers

270 (c)


The directive influence order and tendency to release electron for *o*- and *p*-directing group is,  
 $\text{O}_2^- > \text{NR}_2 > \text{NHR} > \text{NH}_2 > \text{OH} > \text{OCH}_3$   
 $\approx \text{NHCOCH}_3 > \text{CH}_3 > \text{X}$

271 (d)

In organometallic compound, carbon atom is directly bonded to metal atom. Methyl lithium ( $\text{CH}_3\text{Li}$ ) is an organometallic compound.

272 (a)

This is crossed Cannizzaro's reaction in which HCHO is oxidized.



273 (d)


Gammexane is  $\text{C}_6\text{H}_6\text{Cl}_6$ .

274 (b)


Each ligand donates one electron pair.

275 (c)

7. is isomer of  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$  which is used as an anticancer drug for treating several types of malignant tumours.




276 (d)



277 (d)

Petroleum and coal are main sources of aromatic compounds.


279 (c)



Butane-1,2,4-tricarbonitrile

280 (b)

Fac-mer isomerism is associated with  $[\text{MA}_3\text{B}_3]$  type complexes.



282 (c)


Intramolecular H-bonding gives rises to lower m.p.

283 (b)



Mn does not form mononuclear carbonyl.

284 (b)

Ethylenediamine tetraacetic acid is a hexadentate ligand because it has six donor centres.



285 (b)




Both are position isomers

286 (c)

Hybridisation	Geometry of complex
$sp^3$	Tetrahedral
$dsp^2$	Square planar
$d^2sp^3$	Octahedral
$sp^2d^2$	Not possible

287 (b)



288 (d)

Waxes are not obtained by destructive distillation of wood or coal.

289 (c)

Formaldehyde and benzophenone are also obtained.

290 (b)

It ionizes to  $\text{Fe}^{3+}$  and  $\text{SO}_4^{2-}$ .

291 (d)

These are the concepts of Werner's theory.

293 (b)

In  $\text{K}_3[\text{Co}(\text{CO}_3)_3]$ , cobalt shows the +3 oxidation state i.e., ( $d^6$ ) ion. Hence, Co (+3) has four unpaired electrons so, it is paramagnetic. The magnetic moment of Co(+3)

$$\begin{aligned} \text{In } \text{K}_3[\text{Co}(\text{CO}_3)_3] &= \sqrt{n(n+2)} \text{ BM} \\ &= \sqrt{4(4+2)} \text{ BM} = 4.9 \text{ BM} \end{aligned}$$

Where,  $n$  = number of unpaired electrons


$\text{CO}_3^{2-}$  is a weak field bidentate ligand, so  $3\text{CO}_3^{2-}$  ligands occupy six orbitals, thus it shows  $sp^3 d^2$  hybridisation and octahedral in shape.

296 (a)


Some heterocyclic compounds (hetero aromatics) possess aromatic nature. Follow Hückel rule.

297 (c)

Nickel reacts with dimethylglyoxime to give red ppt. of nickel-dimethyl glyoxime complex.



298 (b)



299 (d)

$-\text{CH}_3$  gp. is *o*- and *p*-directing  $\text{FeCl}_3$  is halogen carrier.

300 (b)

It has coordination number of six and thus, should have six hybridized orbitals, *i.e.*,  $d^2sp^3$ - hybridization.

301 (a)

For  $\text{K}_4[\text{Fe}(\text{CN})_6]$ , the EAN of  $\text{Fe}^{2+}$  ion =  $(26 - 2 + 12) = 36$ .

Hence it follows EAN rule, as its EAN is equal to number of electrons of Kr (inert gas), *i.e.*, 36.

302 (b)

$\text{OCH}_3$  gp. an *o*- and *p*-directing group activates ring for  $S_E$  reactions.

303 (b)

Anilium hydrochloride gives white ppt. with  $\text{AgNO}_3$ .

305 (a)

This is a fact for given statement.

307 (b)

*Ortho*, *meta* and *para*.

308 (b)

Follow IUPAC rules.

309 (b)

8.  $\text{Mn}^+ = 3d^5, 4s^1$ . In presence of CO effective configuration  $3d^6, 4s^0$ .

Three lone pair of back bonding with vacant orbital of C in CO.

9.  $\text{Fe}^0 = 3d^6, 4s^2$ . In presence of CO effective configuration  $3d^8$ .

Four lone pair for back bonding with CO.

10.  $\text{Cr}^0 = 3d^4, 4s^2$ . Effective configuration =

$3d^6$ .

Three lone pair for back bonding with CO.

11.  $\text{V}^- = 3d^4, 4s^2$ . Effective configuration =  $3d^6$ .

Three lone pair for back bonding with CO.

Maximum back bonding in  $\text{Fe}(\text{CO})_5$ , therefore CO bond order is lowest here.

311 (b)

$d^4$ : Forms outer complex in high spin and forms inner complex in low spin. It cannot form octahedral complex.

$d^6$ : In low spin it forms inner octahedral complex and in high spin forms outer octahedral complex.

$d^8$ : Forms only outer spin octahedral complex.

312 (c)

Bromo group, *o*- and *p*-directing.

313 (c)

The given statement represents only ether.

314 (a)

Due to asymmetric carbon atom.

315 (a)

This is a fact for the given statement.

316 (c)

$\text{CN}^-$  and  $\text{OH}^-$  are strong nucleophiles.  $[\text{Fe}(\text{OH})_5]^{3-}$  is not formed.

317 (b)

The prefixes erythro and threo are used in systems containing two asymmetric carbons when two of the groups are the same and the third is different. The erythro-isomer has identical groups as the same side when a drawn in Fischer projection and threo-isomer has them on opposite side


318 (b)

Main fractions of coal-tar and the compounds present there in are:

	Main fraction	Temp. range	Chief constituents

1.	Light oil or crude	80-170°C	Benzene, toluene, xylenes, etc.
2.	naphtha	170-230°C	Phenol, naphthalene, pyridine, etc.
3.	Middle oil or carbolic acid	230-270°C	Cresols, naphthalene quinolone, etc.
4.	Heavy oil or creosote oil	270-360°C	Anthracene, phenanthrene, etc.
5.	Green oil or anthracene oil	Residue	90-94% of carbon
	Pitch		

Tris-(ethylenediamine) cobalt(III) bromide ( $[\text{Co}(\text{en})_3]\text{Br}_3$ ) exhibits optical isomerism.



319 (d)

The order of reactivity depends on the stability of intermediate carbocation formed due to heterolytic cleavage of C—X bond.

320 (d)

Greater is the number of chelate rings, greater is stability of the chelate. Hence, five fused cyclic system is most stable for a chelate.

321 (b)

$\text{HNO}_3 + \text{H}_2\text{SO}_4$  acts as nitration mixture.

323 (d)

The negative charge density on V-atom favours easy electron pair donation.

324 (b)

Due to acidic nature.

325 (d)

If a substance rotates the plane polarised light in clockwise direction it is dextrorotatory (+). If it rotates the plane polarised light in anticlockwise direction then it is laevorotatory (–)

326 (c)



327 (b)

EAN of Fe in  $\text{K}_4\text{Fe}(\text{CN})_6 = 26 - 2 + 2 \times 6 = 36$ ; the at. no. of next inert gas.

329 (b)

Both  $\text{Ni}(\text{CO})_4$  and  $\text{Ni}(\text{PPh}_3)_2\text{Cl}_2$  have  $sp^3$ -hybridisation

331 (d)



All are examples of strong ligand field, because all have greater value of  $\Delta$ ;  $\Delta$  represents the strength of ligand field.

332 (a)

Due to rearrangement because  $2^\circ$  carbon is more stable than  $1^\circ$  carbon.

333 (d)

334 (c)



Number of unpaired electrons=5

$$\text{Magnetic moment} = \sqrt{n(n+2)} = \sqrt{s(s+2)}$$

$$= \sqrt{35} = 5.91 \text{ BM}$$

335 (b)

$$\text{EAN of Cr} = 24 - 3 + 6 \times 2 = 33.$$


336 (d)

Paramagnetic character  $\propto$  number of unpaired electrons.

$^{25}\text{Mn}^{2+}$  ion has maximum unpaired (five unpaired electrons)

electrons. So,  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$  is most paramagnetic.

337 (d)




**Note :** The reaction gives 2, 4, 6-trinitrophenol.

Choice is not given. Only option left is *o*-nitrophenol, which is not formed in this course of reaction.

338 (d)

$\text{NH}_2$  in aniline is highly susceptible to oxidant and therefore nitration of aniline is carried out by protecting it against oxidation by acetyl chloride.



339 (b)

—COOH is *meta*-directing group.

340 (d)



Pentaammine nitrito -N- cobalt (III) chloride.

341 (b)

The ionisation isomer of


$[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}(\text{NO}_2)]\text{Cl}$  is  $[\text{Cr}(\text{H}_2\text{O})_4\text{Cl}_2](\text{NO}_2)$   
because of exchanging of ligand and counter ions.

342 (c)

All can be prepared from phenol.

343 (a)

Only iodobenzene gives Ulmann's reaction,



344 (c)

Fischer projection can be manipulated by rotating a group of any three ligands in clockwise (D) or anticlockwise (L) direction, the fourth ligand does not change its position

345 (b)


A strong field ligand produces low spin complexes.

347 (a)


Ni in  $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$  has two unpaired electrons in it.

348 (b)

Compounds in which a chiral centre is part of a ring are handed in an analogous fashion




351 (a)



352 (b)

The formula of hexamine copper (II) sulphate is  $[\text{Cu}(\text{NH}_3)_6]\text{SO}_4$ . It dissolves in water as




354 (b)

Complex of type  $[\text{M}(\text{AA})_3]$  show optical isomerism.


355 (b)

Electronic configuration of  $\text{Fe}^{2+}$  in  $\text{K}_4\text{Fe}(\text{CN})_6$  is:



$\times\times$  Electron pair donated by  $\text{CN}^-$ .

356 (d)



357 (b)

Oxidation state of Ni in  $\text{K}_2\text{NiF}_6$  is +4; the highest among all.

359 (d)

Substituents always get higher number than the principal functional group while, numbering the longest possible chain

360 (d)


Draw different isomers.

362 (b)

Metal carbonyl organometallic compounds possess both  $\sigma$ - and  $\pi$ -characters.

363 (b)

$\text{F}^-$  is a weak field ligand and  $[\text{FeF}_6]^{3-}$  is an outer-orbital complex





only for  $K^+$  ions and not for  $Fe^{2+}$  ions.



366 (a)

Only primary valencies are ionized.



0.01 mole of each by 0.01 mole of reactants.

367 (c)

Wilkinson's catalyst,  $(Ph_3P)_3RhCl$



ie,  $dsp^2$  hybridisation

Rh atom in Wilkinson's catalyst is  $dsp^2$  hybridised giving a square planar shape to the molecule

368 (d)

$H_2O$  is weak field ligand, thus  $Co^{2+}$  has only 3 unpaired electrons.


369 (a)

If an enantiomerically pure acid is treated with racemic mixture of an alcohol having a chiral carbon, the product formed will be optically active mixture

370 (d)

Cyclopropane is most strained since it has a maximum angle strain of  $24^\circ - 44'$


371 (c)



this is Gattermann-Koch reaction to introduce  $-CHO$  gp. in  $C_6H_6$  nucleus.

372 (a)

Mn in  $Mn(CN)_6^{4-}$  has configuration:



373 (d)

It does not ionize to give  $Cl^-$  ions and thus, white ppt. of  $AgCl$  will not be obtained.

375 (c)



$$x + 0 + (-1) \times 3 = 0$$

$$x + (-3) = 0$$

$$x = +3$$

377 (c)

$Ni^{2+}$  has two unpaired ( $3d^8$ ) electrons.  $CN^-$  is strong field ligand and thus all the eight electrons are paired giving  $dsp^2$ -hybridisation.

379 (d)

Coal-tar is source of all these.

380 (c)

Halogen attached to benzene nucleus is stabilized due to resonance.

382 (a)


$-NO_2$  group is reduced to  $-NH_2$  by  $Sn/HCl$ .

383 (d)

Each central atom attains the EAN equal to at. No. of next inert gas Kr, i. e., 36.

384 (c)

A bidentate ligand has two donor sites available for coordination, e. g.,



385 (d)

$-N_2Cl$  is reduced to H by either of these reducing agents.

386 (c)

Fe is present in the form of complex ion, i. e.,  $[Fe(CN)_6]^{3-}$  which is not ionized to  $Fe^{3+}$  and  $CN^-$ .

387 (a)



The molecular formula of both of the above compounds is same but on ionisation they give different ions in solution, so they are called ionization isomers.

390 (c)


Phenols are weak acids and do not react with  $NaHCO_3$  (a weak base).

391 (b)

$C_6H_5CH = CHCOOH$  is cinnamic acid; it has unsaturation.

392 (a)

Magnetic moment of




$$\text{Magnetic moment} = \sqrt{n(n+2)}$$

$n$  = number of unpaired electrons present in molecule

$$1.7 = \sqrt{n(n+2)}$$

$$-n^2 + 2n - 2.89 = 0 \text{ then } n = 0.97 \text{ or } 1$$

393 (c)




This compound contains 9 carbon atoms and corresponding alkane is nonane. Three bridges

contain 5, 2 and 0 carbon atoms. Therefore, the name of the compound is bicyclo [5.2.0] nonane

395 (c)

The compound have structure as written below



bicyclo [3.2.1] octane



396 (d)

It is *m*-directing gp.

397 (a)

Carbylamines reaction.

400 (d)



401 (a)

The stability order of conformations of cyclohexane is

Chair > twist boat > boat > half chair

402 (b)

Phenolic group is susceptible for oxidation and thus, to obtain *o*- and *p*-nitrophenol dil.  $HNO_3$  is used in place of conc.  $HNO_3 + H_2SO_4$ .

403 (c)

Lower is mol. Wt. lower is b.p., also 1, 2-dihydroxy benzene show chelation and thus have lower b.p. than 1,3 and 1,4-derivatives.

Phenol	1,2-dihydroxyb	1,3-dihydroxyb
	enzene	enzene

enzene

m.p. $43^\circ C$	$< 105^\circ C$	$< 110^\circ C$
-------------------	-----------------	-----------------

$< 170^\circ C$

b. p.	$<$	$<$
-------	-----	-----

$<$

405 (b)



407 (b)

Phenols are weakly acidic due to resonance.

408 (b)

Follow IUPAC rules.

409 (a)

Compounds having coordination number six and following the general formula show geometrical and optical isomerism.

$M_{A_4B_2}$ ,  $M_{A_4BC}$ ,  $M_{A_3B_3}$  and  $M_{(AA)_2B_2}$  show geometrical isomerism and

$M_{A_2X_2Y_2}$ ,  $M_{A_2X_2YZ}$ ,  $M_{A_2XYZL}$

$M_{ABXYZL}$ ,  $M_{(AA)_3}$ ,  $M_{(AA)_2X_2}$  show optical isomerism. (AA) is bidentate ligand.

410 (b)

$[Co(NH_3)_4Cl_2]Cl$ , Its IUPAC name is tetraammine dichloro cobalt III chloride.

411 (c)

The ease of hydrolysis depends upon the magnitude of the +ve charge on the carbonyl group.

412 (d)

All are the common uses of nitrobenzene.

413 (b)



$$x + (4 \times -1) = -4$$

$$x - 4 = -4$$

$$x = 0$$

414 (a)

Presence of  $-OH$  gp. in  $C_6H_6$  nucleus increases acidic nature.

416 (b)



Halogen attached on side chain behaves as in aliphatic molecule.

417 (d)

Cyclohexane is an aliphatic cyclic compound.

418 (c)



12. In  $[Cu(NH_3)_4]^{2+}$ , Cu is present as  $Cu^{2+}$



( $NH_3$  being a strong field ligand shifts one electron from 3d-orbital to 4p-orbital.)

13. In  $[Ni(CO)_4]$ , CO is a neutral ligand

14. In  $[Fe(CN)_6]^{3-}$ , Fe is present as  $Fe^{3+}$ .




Thus, its hybridization is  $d^2sp^3$  not  $sp^3d^2$ , i.e., it is

an inner orbital complex.

15.  $[\text{Co}(\text{en})_3]^{3+}$  contains total 36 electrons, *i. e.* follows EAN rule.

420 (c)


Optical isomerism is very common in octahedral complexes having general formula



where *AA* is symmetrical bidentate ligand like  $\begin{array}{c} \text{COO}^- \\ | \\ \text{AB} \\ | \\ \text{COO}^- \end{array}$  and *AB*

is unsymmetrical bidentate ligand.

421 (c)



422 (a)

$\text{C}_6\text{H}_5\text{C}$  is benzo gp.

423 (c)

It is the reason for given fact.

424 (b)

$d^6$ -cation with low spin has electronic configuration  $t_{2g}^6 e_g^0$ .

Total energy =  $(-0.4 \Delta_0 \text{ per } e^- \times 6) + (e^- \text{ pairing energy of 3 pairs})$

$$= -2.4 \Delta_0 + 3P$$


$$= -\frac{12}{5} \Delta_0 + 3P$$

426 (c)

It is DDT, *i. e.*, *p, p'*-dichloro diphenyl trichloroethane.

428 (b)





$[\text{Cr}(\text{NO}_2)(\text{NH}_3)_5] \text{Cl}$  show linkage isomerism.



429 (d)

Cl, being a weak field ligand, does not cause pairing of *d*-electrons of the metal atom and thus, forms outer orbital complex as.


In  $[\text{Fe}(\text{Cl})_6]^{3-}$ , Fe is present as  $\text{Fe}^{3+}$



430 (d)

$pK_a$  for (a), (b), (c) and (d) are 4.17, 4.09, 3.49 and 3.43 respectively.

431 (a)



16. In this complex, four donor sites are occupied by oxygen and two donor sites are occupied by nitrogen.

17. This complex is six coordinated.

18. Complex  $[\text{MgEDTA}]^{2-}$  is colourless.


19. Increase in  $[\text{H}^+]$  decreases pH of the solution.

433 (c)

Different compounds having the same molecular formula but different properties are called isomers

434 (c)

In the compound




Numbering will be done from this end because both are side chains and  $-\text{OCH}_3$  is smaller than  $-\text{OC}_2\text{H}_5$

3-ethoxy-1-methoxy-propane

435 (a)


Each carbon in benzene is  $sp^2$ -hybridized.

436 (b)



So, gauche form stabilized by intermolecular hydrogen bonding hence, gauche is more stable than anti

437 (a)



438 (b)

Carbocyclic compounds which resemble aliphatic compounds in their properties are called alicyclic compounds

439 (a)



—NO<sub>2</sub> gp. is deactivating gp.

440 (b)

The molecule contains three chlorine atoms out of which only two are ionized.

441 (b)

[Co(en)<sub>2</sub>Cl<sub>2</sub>]<sup>+</sup> have three optical isomers which are given below.



442 (c)

The reaction carried out in alkaline pH, *ie*, 9 – 11

444 (d)

NO<sub>2</sub> gp. withdraws electrons from *o*- and *p*-position and thus, deactivates the ring. This deactivation stabilises the negatively charged intermediates formed during reaction and thus, replacement of—Cl becomes easier.

446 (a)

Ti(C<sub>2</sub>H<sub>5</sub>)<sub>4</sub> is an organometallic compound because there is direct bonding of metal ion with carbon.

447 (a)

Octahedral Co(NH<sub>3</sub>)<sub>4</sub>Br<sub>2</sub>Cl shows ionisation and geometrical isomerism.

In ionisation isomerism ligands show different coordination sphere and the anions present outside the coordination sphere.


These are exchanged with each other as follows  
Co(NH<sub>3</sub>)<sub>4</sub>Br<sub>2</sub>Cl ⇌ [Co(NH<sub>3</sub>)<sub>4</sub>Br<sub>2</sub>]<sup>+</sup> + Cl<sup>-</sup>

I

[Co(NH<sub>3</sub>)<sub>4</sub>BrCl]Br ⇌ [Co(NH<sub>3</sub>)<sub>4</sub>BrCl]<sup>+</sup> + Br<sup>-</sup>

II

In geometrical isomerism, coordination number of central atom (cobalt) is six and shape is octahedral, so it shows following geometrical isomers.



448 (a)


Aliphatic amines are stronger base than aromatic amines.

449 (b)

Follow Hückel rule.

450 (d)

⇌ sign represents oscillating structures (Kekule) for C<sub>6</sub>H<sub>6</sub> ;



451 (b)

C<sub>6</sub>H<sub>5</sub>CHCl<sub>2</sub>  $\xrightarrow{\text{HOH}}$  C<sub>6</sub>H<sub>5</sub>CHO.

453 (c)

Cr has coordination no. 6 in its carbonyl and other complexes.

454 (d)

Neutral FeCl<sub>3</sub> (*aq.*) gives violet coloured complex with phenol.

455 (a)

[CoF<sub>6</sub>]<sup>3-</sup> is an outer complex having sp<sup>3</sup>d<sup>2</sup>-hybridization.

456 (d)

[Sc(CN)<sub>6</sub>]<sup>3-</sup>


Sc=21=1s<sup>2</sup>, 2s<sup>2</sup>, 2p<sup>6</sup>, 3s<sup>2</sup>, 3p<sup>6</sup>, 4s<sup>2</sup>, 3d<sup>1</sup>

3d                      4s                      4d





No unpaired electron

Ni(CN)<sub>4</sub><sup>2-</sup>






Ni<sup>2+</sup>







It has no unpaired electron hence, diamagnetic.






(Pairing due to  $\text{CN}^-$ )



(due to  $\text{CN}^-$ )

It has one unpaired electron so, paramagnetic.

458 (a)



$\text{CN}^-$  is strong field ligand, it cause pairing while  $\text{C}_2\text{O}_4^{2-}$  and  $\text{F}^-$  are weak field ligands and don't causes pairing.

Hence,  $[\text{Fe}(\text{CN})_6]^{4-}$  due to the absence of unpaired electrons is diamagnetic.

459 (c)

$_{28}\text{Ni}^{2+}$  in  $[\text{Ni}(\text{NH}_3)_6]^{2+}$  has  $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^8$  configuration. It uses 4th orbital to show  $sp^3 d^2$  hybridisation to form outer complex with 2 unpaired electrons in 3d-orbital.


460 (c)

Moth repellent due to insecticide nature.

461 (c)

The number of unidentate ligand in the complex ion is called coordination number.


462 (b)



7 double bonds and thus, 14  $\pi$ - electrons.

463 (b)

In ethane and cyclohexane, staggered and chair forms are more stable respectively



Staggered form of ethane Chair form of cyclohexane

464 (d)

$(\text{CH}_3)_4\text{Sn}$  has no  $\pi$ -bond.

465 (a)

Halogens however *o*- and *p*-directing group but deactivate ring for  $\text{S}_\text{E}$  reactions due to electron withdrawing nature.

466 (b)

It is the reason for the fact.

467 (a)

$-\text{NO}_2$  gp. is deactivating gp.

470 (c)

The side reaction produces diphenyl ether. The yield may be increased by adding little diphenyl ether with  $\text{C}_6\text{H}_5\text{Cl} + \text{NaOH}$ .

471 (c)

Atoms or groups donating electron pair to metal are ligands.

472 (d)

Aniline is insoluble in water.

473 (d)

$\text{CrO}_2\text{Cl}_2$  has + 6 oxidation state of Cr.


475 (b)

It is a method to estimate hardness of water.

476 (a)

Both are non-polar; like gets dissolved in like.

477 (b)



There are two carbon atoms, *ie*, C-2 and C-4 are tertiary C-atoms



493 (d)


Metal carbonyl are regarded as the coordination compounds formed by the donation of lone pair of electron of CO into the suitable empty orbital of zero valent transition metals such as Ni, Fe etc.

Therefore, the  $M-C$  bond is coordinate covalent.

494 (d)

When the three ligands (with same donor atoms) are on the same triangular face of the octahedron, the isomer is called *facial* or *fac* isomer.

The octahedral complex is *facial* or *fac* isomer.



In this complex, the three ligands are on the same triangular face of the octahedron.

495 (a)

Number of unpaired electrons in  $[Fe(CN)_6]^{4-}$  is zero.

Thus, magnetic moment

$$= \sqrt{n(n+2)} = 0 \text{ BM}$$

( $n$  = unpaired electrons)

$$n \text{ in } [MnCl_4]^{2-} = 5, \sqrt{35} \text{ BM}$$

$$n \text{ in } [CoCl_4]^{2-} = 3, \sqrt{15} \text{ BM}$$

496 (a)

Orange-red dye is formed with aniline.

499 (a)

It is a fact. Follow *ortho* effect.


500 (d)

Half chair is transition state conformation between the chair and boat conformation. The energy difference between the chair and half chair conformation being  $44 \text{ kJ mol}^{-1}$ . Hence it is most unstable

501 (a)

The  $M-C$   $\pi$ -bond in metal carbonyl which is formed by the donation of an electron pair from a filled  $d$ -orbital of metal into the vacant antibonding  $\pi$ -orbital of CO, strengthens the  $M-C$   $\sigma$ -bond. This is called synergic effect and is usually observed in metal carbonyls. Thus  $[Ni(CO)_4]$  exhibits synergic effect.

503 (b)




504 (d)

Follow IUPAC rules.

505 (b)

The structure of the compound is



1,7,7-trimethyl bicyclo [2.2.1] heptan-2-one


506 (d)

$-CH_3$  gp. is *ortho* and *para* directing.

507 (a)

Estimation of calcium and magnesium is done by EDTA.

508 (d)




Due to the presence of chiral centre it shows the optical activity and its mirror image are non superimposable hence it shows one enantiomer pair




509 (b)

$C_6H_6$  and other aromatic compounds show characteristics  $S_E$  reactions.

510 (d)



511 (b)




512 (a)






The attacking species in sulphonation is  $SO_3$ .  
 $H_2SO_4 \rightarrow H_3O^+ + SO_3 + HSO_4^-$

513 (d)

$CHCl_3$  has no reaction with  $Br_2$ .

514 (c)





### $sp^3$ -hybrid orbitals, tetrahedral

$Cl^-$  is a weak ligand and thus unpaired electrons are not paired. Lone pairs from  $4Cl^-$  are accommodated in four  $sp^3$  hybrid orbitals.

$N$ =unpaired electron=2, paramagnetic

Magnetic moment (spin only)


$$= \sqrt{N(N+2)} \text{ BM} = \sqrt{8} =$$

2.828 BM

515 (b)

Pyridine shows  $S_E$  reactions at position-3 preferentially and at 2,4-positions under specific conditions.

516 (a)




517 (a)

Its coordination number will be 6 because it is bonded with three bidentate ligands

$$x + 3(-2) + 3(+1) = 0 \Rightarrow x = +3$$


518 (c)

The four ions on ionisation are possible only when three  $Cl^-$  are outside the coordination sphere.



519 (a)

In presence of  $H_2O$  which is a weak ligand no pairing occurs which results in unpaired electrons left in the compound, due to which it shows paramagnetism




520 (a)

This is a fact.

521 (b)

It is outer complex having  $sp^3 d^2 [CoF_6]^{3-}$  ion.

Electronic configuration of  $Co^{3+}$  in  $[CoF_6]^{3-}$  is:



×× Electron pair donated by  $F^-$

523 (d)

The two given compound have same composition but in solution both will give different ions. The isomerism is known as ionisation isomerism

524 (a)

Coordination number is the maximum covalency shown by a metal or metal ion. It is the maximum number of ligands attached to metal by sigma bonds or coordinate bonds.

525 (a)

$C_6H_5COOH + N_3H \rightarrow C_6H_5NH_2 + CO_2 + N_2$ ; this is Schmidt's reaction.


526 (d)

Each possess the tendency to have coordination number equal to six.



527 (d)

Hexadentate ligand donates six pair of electrons to central atom.

(a) 2, 2-dipyridyl-bidentate ligand




(b) DMG-bidentate ligand



(c) Ethylenediamine-pentadentate ligand

∴ None of the given ligand is hexadentate ligand.


528 (d)



529 (d)

20.  $d^5$  in strong field


$n$  =unpaired electron=1



$$\text{Magnetic moment} = \sqrt{n(n+2)} \text{ BM}$$

$$= \sqrt{3} \text{ BM} = 1.73 \text{ BM}$$


21.  $d^3$  in strong/weak field



$$n = 3$$

$$\text{Magnetic moment} = \sqrt{15} = 3.87 \text{ BM}$$


22.  $d^4$  in weak field



$$n = 4$$

$$\text{Magnetic moment} = \sqrt{24} = 4.90 \text{ BM}$$

23.  $d^4$  in strong field





$$n = 2$$

$$\text{Magnetic moment} = \sqrt{8} = 2.83 \text{ BM}$$

531 (a)

$[\text{Ni}(\text{CN})_4]^{2-}$  ion has  $dsp^2$  hybridisation, zero magnetic moment and square planar structure.



It has no unpaired electrons hence, its magnetic moment is zero.


534 (a)



$\text{Sc}^{3+} = [\text{Ar}]3d^0 4s^0$  no unpaired electrons in  $d$ -subshell, so it is diamagnetic and colourless.

536 (d)


In  $[\text{Co}(\text{NH}_3)_6]^{3+}$  the oxidation state of Co is +3.



$[\text{Co}(\text{NH}_3)_6]^{3+}$  does not contain unpaired electron hence, its magnetic moment is zero.


537 (a)

Presence of  $o$ -,  $p$ -directing groups in benzene nucleus activates ring for  $S_E$  reaction. Presence of  $m$ -directing deactivates ring for  $S_E$  reaction. Also halogens deactivating gp. Due to  $-IE$  inspite of  $o$ - and  $p$ -directing nature. The presence of  $m$ -directing groups in benzene nucleus simply decreases electron density at  $o$ - and  $p$ -, whereas no change in electron density at  $m$ -position is noticed.





On the contrary  $o$ - and  $p$ -directing groups in nucleus increases the electron density at  $o$ - and  $p$ -position.

Thus, presence of  $o$ - and  $p$ -directing groups provide seats for  $S_E$  reactions or activates the ring, whereas presence of  $m$ -directing groups does not activate the ring and thus, deactivate the ring for  $S_E$  reactions



539 (b)

The percentage of enolic contents of some common compounds in decreasing order will be



540 (c)

Mn forms  $Mn_2(CO)_{10}$  carbonyl.

541 (c)

Picric acid has phenolic gp.

542 (d)

(i) The sum of oxidation states of all atoms in a compound is zero.

(ii) Oxidation state of metal in carbonyl is zero.

(a)  $K_4Fe(CN)_6$

Let, oxidation state of Fe in  $K_4Fe(CN)_6 = x$

$$\therefore +4 + x + (-1 \times 6) = 0$$

$$\therefore x = +2$$

(b)  $K_2FeO_4$

Let, oxidation state of Fe in  $K_2FeO_4 = x$

$$\therefore +1 \times 2 + x + (-2 \times 4) = 0$$

$$\therefore x = +6$$

(c)  $Fe_2O_3$

Let, oxidation state of Fe in  $Fe_2O_3 = x$

$$\therefore 2x + (-2 \times 3) = 0$$

or  $2x = 6$

$$\therefore x = \frac{6}{2} = +3$$

(d)  $Fe(CO)_5$

Oxidation state of Fe in  $Fe(CO)_5 = 0$

$\therefore$  Oxidation state of Fe is least in  $Fe(CO)_5$ .

543 (a)

The name of  $[Pt(NH_3)_4Cl_2]^{2+}$ ,  $[PtCl_4]^{2-}$  is tetraamminedichloroplatinum (IV) tetrachloroplatinate (II). Since, positive ion is written first and negative ion later.

544 (b)

Resorcinol is *meta* hydroxyphenol.

545 (d)

The compounds of nickel are green coloured due to *d-d* transition in presence of ligand in  $Ni^{2+}$  cations.

546 (c)




547 (d)

Geometrical isomerism is found in compounds having coordination no. 4 (square planar and not tetrahedral shape) as well as coordination no. 6. Coordination no. 4 (square planar)




Showing geometrical isomerism  $M_{ABCD}$

Coordination no. 6 (octahedral)



Showing geometrical isomerism.



548 (c)

Follow text.

549 (a)

Presence of electron repelling gp. decreases the acidic strength.

$C_6H_5COOH$	<i>p</i> -methyl benzoic acid	<i>p</i> -chloro phenol
--------------	-------------------------------	-------------------------

$K_a$	$6.76 \times 10^{-5}$	$1.26 \times 10^{-5}$	$4.16 \times 10^{-10}$
-------	-----------------------	-----------------------	------------------------

551 (b)

It becomes brown (due to oxidation) on standing in air.

552 (b)

It is characteristics of aromatic compounds.

553 (d)

The colour of the complex  $CoCl_3 \cdot 5NH_3 \cdot H_2O$  is pink.

554 (b)


$$2 + 4 \times (-1) = x, \therefore x = -2$$

555 (a)

Halide ligands have low values of  $\Delta$ .

556 (d)

Electronic configuration of  $Co^{3+}$  in  $[CoF_6]^{3-}$  is:



$\times\times$  Electron pair donated by  $F^-$

558 (d)

Let the oxidation state of Fe in

$[Fe(H_2O)_5NO]SO_4$  is  $x$ .

$[Fe(H_2O)_5NO]^{2+}$

$$\Rightarrow x + 0 + 1 = 2$$

$$\therefore x = +1$$

Here, NO exists as nitrosyl ion ( $NO^+$ ).


559 (c)

$EAN = Z - (ON) + 2$  (C.N.)

where, O.N. = oxidation number



C.N. = coordination number

$Z$  = atomic number



561 (a)

In the complex  $\text{K}_3[\text{FeF}_6]$ , Fe is present in +3 oxidation state.



Hence, number of unpaired electrons is five as F is weak ligand.

$$\text{Magnetic moment} = \sqrt{n(n+2)}$$

$$= \sqrt{5(5+2)}$$

$$= 5.91 \text{ BM}$$

563 (c)


No doubt C—D bond cleavage is slower than C—H bond due to isotopic effect but rate of overall substitution is determined by the slow attachment of electrophile to carbocation (Morrison-Boyd 15.14P-532).

564 (a)

Due to synergic bond formation, bond order decreases and bond length increases a little.


566 (c)

It is a bicyclic compound having two common carbon atoms and three bridges. So, the IUPAC name is



5, 6-dimethyl bicyclo [2.2.1] hept-2-ene

567 (c)



568 (a)

The effective magnetic moment of a paramagnetic substance is given by the relation

$$\mu = \sqrt{n(n+2)} \text{ BM.}$$

where,  $n$  = number of unpaired electrons.

In  $[\text{FeF}_6]^{3-}$ ,  $\text{Fe}^{3+}$  has five unpaired electrons.



$$\mu \text{ of } [\text{FeF}_6]^{3-} = \sqrt{n(n+2)}$$

$$= \sqrt{5(5+2)}$$

$$= \sqrt{35} = 5.92 \text{ BM.}$$


569 (c)

$[\text{PtCl}_2 \cdot \text{P}(\text{C}_2\text{H}_5)_3]_2$  can exhibit geometrical isomerism, the geometrical isomers are



570 (c)

The minimum possible isomers of compound will be




571 (b)

$\text{C}_2\text{O}_4^{2-}$  is a bidentate group. As the complex contains three bidentate groups, the central metal ion has a coordination number of 6.

572 (d)

$[\text{Co}(\text{en})_3]^{3+}$  has *d* and *l* forms as



573 (b)

Ru forms two carbonyls with zero oxidation number. Mononuclear  $\text{Ru}(\text{CO})_5$  and trinuclear  $\text{Ru}_3(\text{CO})_{12}$ .

574 (c)

Oxidation state of nitrogen in  $(\text{N}_2\text{H}_5)_2\text{SO}_4$  is



$$4x + 10 - 2 = 0$$

$$x = -2$$

575 (d)

Linkage isomerism is exhibited by ambidentate ligands (ligands having two coordination sites). e.g.,  $\text{NO}_2^-$ .

If the bonding is through N, the ligand is named as nitro and if it is through O, it is named as nitrito.



577 (a)

Due to resonance of electron pair in aniline, nitroaniline and acetanilide, these are weaker than  $C_6H_5CH_2$  which does not involve lone pair of N in resonance. The basic order is: Benzyl amine > Aniline > Acetanilide > Nitroaniline.

578 (b)

Effective atomic no. (EAN) = at. No. of central atom - oxidation state + 2 × (no. of ligands) = 28 - 0 + 2 × 4 = 36

EAN = 78 - 4 + 2 × 6 = 86.

579 (b)

$[Cu(NH_3)_4]SO_3 \rightleftharpoons [Cu(NH_3)_4]^{2+} + SO_4^{2-}$

580 (b)

Ammonia is not an ambident legand so it can donate electron only by N-atom

582 (b)


12σ and 3π.

583 (c)

-OH gp. is activating whereas Cl- is deactivating.

-CH<sub>3</sub> gp. is less activating than OH.

584 (d)



The compound  $\begin{matrix} H & H \\ Br & Br \\ H & Br \end{matrix}$  is symmetrical with respect to centre of the molecule


585 (b)

Two *cis* and *trans* forms.

586 (c)

*p*-nitrophenol is more stronger acid than phenol.


587 (b)



588 (d)

Staggered conformation is most stable due to its minimum energy

589 (a)



590 (c)

$Fe^{2+}$ ,  $Co^{5+}$ ,  $Ti^{3+}$ , and  $V^{3+}$  have 4, 4, 1, 2 unpaired electron respectively. The pairing leads  $Fe^{2+}$  with no unpaired electron.

591 (b)


Os ( $Z=76$ ):  $[Xe] 4f^{14}, 5d^6, 6s^2$

Hence, the coordination number in an osmium complex may increase to 8.

592 (c)


Phenol has antiseptic property.

593 (c)




594 (d)

$[Co(en)_3Cl_3]$  ie,  $[Co(en)_3]^{3+}$



*cis*  $[Co(en)_2Cl_2]Cl$  ie, *cis*  $[Co(en)_2Cl_2]^+$



595 (b)

The compound in which ligands form ring with the metal are called chelate complex.

597 (c)

Benzaldehyde undergoes Cannizzaro's reaction.

600 (c)


Coordination isomerism is possible when both positive and negative ions of a salt are complex ions and the two isomers differ in distribution of ligands in the cation and the anion

601 (a)


This is bromination of acetanilide, a  $S_E$  reaction.

602 (b)

The primary valency is ionizable valency. It corresponds to oxidation state of metal. The primary valency is always satisfied by anion.




∴ Number of primary valency is 3



(B)


∴ Number of primary valency is 2



∴ Number of primary valency is 1.

603 (d)

The carbon atom which is attached to three carbon atoms is called tertiary carbon atom.  $C_6H_{14}$  has two tertiary carbons hence, its structure is as



2,3-dimethyl butane

604 (b)

The ligand  $\text{NO}_2$  has two types of linkage with central atom. In  $\text{NO}_2$ , it is the N-atom which is donor and in  $\text{O}-\text{NO}$  it is the O atom which donates electron pair.

605 (d)

Tetraethyl lead is organometallic compound.

606 (d)

$\text{C}_6\text{H}_5\text{OH} + \text{PCl}_5 \rightarrow (\text{C}_6\text{H}_5)_3\text{PO}_4$  is main product.

609 (c)

Directive influence order

$-\text{OH} > -\text{OCH}_3 > -\text{CH}_3 > -\text{NHCOCH}_3 > -\text{CH}_2\text{OH}$  of *o*-, *p*- gps. This is due to effect of +R directing influence of gp.

610 (d)


$[\text{PtCl}_4]^{2-}$  shows  $dsp^2$  hybridization because internal *d*-orbitals participate in its hybridization.

611 (b)

Dynamite, TNT, TNB, trinitroglycerine are explosive.

612 (b)

In  $\text{Hg}[\text{Co}(\text{SCN})_4]$ , Co is present as  $\text{Co}^{2+}$ . The configuration of  $\text{Co}^{2+}$  is given as following  $[\text{Ar}]3d^7 4s^0$



$$\therefore \text{Magnetic moment } (\mu) = \sqrt{n(n+2)} \\
 \sqrt{3(3+2)} = \sqrt{15} \text{ BM}$$

614 (c)

In metal carbonyls CO has ox. no. equal to zero.

615 (a)

$[\text{NiCl}_4]^{2-}$  has tetrahedral shape. In this complex, Ni is in the +2 oxidation state and  $\text{Ni}^{2+}$  ion always forms tetrahedral complexes

616 (a)

It is a differentiating point in between complex and double salt.

617 (d)

All possess lesser number of unpaired electrons.

618 (c)


Structures  $\text{K}_4[\text{Fe}(\text{CN})_6]$ ,  $\text{K}_3[\text{Co}(\text{CN})_6]$ ,  $\text{K}_2[\text{Ni}(\text{CN})_4]$  are diamagnetic.

619 (c)

Wilkinson's catalyst is used for hydrogenation of alkenes

620 (c)

Due to *ortho* effect; *ortho* benzoic acid is most acidic because its anion is highly stabilized due to strong intramolecular H-bonding.



$$K_a: 1.0 \times 10^{-3} \quad 2.9 \times 10^{-5} \quad 1.26 \times 10^{-5} \quad 6.9 \times 10^{-8}$$

623 (a)

$\text{C}_6\text{H}_5\text{COOH}$  is acid; phenol also as acid.

624 (c)

*Cis*-isomer of  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$  is used as anticancer drug.

626 (b)

$[\text{Co}(\text{en})_2\text{Cl}_2]^+$  shows geometrical as well as optical isomerism

628 (c)

Lab method for preparation of benzaldehyde.

629 (a)



The ligand at least consist one donor atom having a lone pair of electrons which it can donate to metal atom or ion

630 (c)

Aniline is steam volatile.

631 (a)

CFSE (crystal field splitting energy) for octahedral complex,  $\Delta_0$  depends on the strength of negative ligand. Spectrochemically it has been found that the strength of splitting is as follows



Therefore, magnitude of  $\Delta_0$  will be highest in case of  $[\text{Co}(\text{CN})_6]^{3-}$ .

632 (b)




Potassium hexachloroplatinate (IV).

633 (c)

The complex formed by the reaction of  $\text{NiSO}_4$ , pyridine and  $\text{NaNO}_2$  gives  $[\text{Ni}(\text{py})_4](\text{NO}_2)_2$  a

blue-coloured salt.


634 (b)



635 (a)

Only m-cresols give tribromo derivatives on treatment with  $\text{Br}_2$  water.


636 (a)



$\therefore$  Coordination number of  $E = 6$

$\therefore$  Oxidation number of  $E = 3[\text{E}^{3+} + 0 + (-2) + (-1) = 0]$

637 (c)



638

	Complex	Isomerism shown
(a)	$[\text{Co}(\text{en})]^{3+}$	Optical only
(b)	$[\text{Ni}(\text{NH}_3)_5\text{Br}]^+$	No geometrical isomer
(c)	$[\text{Co}(\text{NH}_3)_2(\text{en})_2]^3$	<i>Cis</i> and <i>trans</i>
(d)	$[\text{Cr}(\text{NH}_3)_4(\text{en})]^{3+}$	No geometrical isomer

(c)

639 (c)

Presence of  $-\text{NO}_2$  at  $p$ -position increases acidic character.

640 (b)


Alkanes are saturated hydrocarbons without any functional group, hence can show chain isomerism only

641 (d)


Both have different molecular formulae.

642 (b)

Ni in  $[\text{Ni}(\text{CO})_4]$  : ..... $3s^2, 3p^6, 3d^8, 4s^2$



Ni in  $[\text{Ni}(\text{CN})_4]^{2-}$  : ..... $3s^2, 3p^6, 3d^8, 4s^2$




643 (a)

The product ( $K$ ) is formed through simple nucleophilic substitution while major product ( $L$ ) is formed through  $\sim\text{H}^-$  shift via  $S_N1$  reaction and methoxy group stabilizes the carbocation intermediate of product( $L$ ).

645 (c)

In the Grignard reaction magnesium metal forms an organometallic bond



Grignard reagent

646 (a)


Aromatic hydrocarbons are called arenes with general formula  $\text{C}_n\text{H}_{2n-6y}$ , where  $n \neq 6$  and  $y$  is no. of cyclic rings. Benzene has one ring and  $n = 6$ , i. e., no. of carbon atoms. Thus, general formula is  $\text{C}_6\text{H}_6$ . All other aromatic hydrocarbons are derivative of benzene.

647 (c)

It is a fact.

648 (d)


Tri and tetravalent bridges derived from methane are given the prefix methyno and methyno respectively



methyno-1,1,1,1-tetracyclohexane


649 (a)

The structure of the compound is



2,4,4-trimethyl pentanal.

650 (c)



651 (a)

Complementary colours of absorbed light are seen.

652 (d)


Presence of  $-\text{SO}_3\text{H}$  gp. increases solubility of drug or dyes.

653 (d)

It is a fact.

654 (a)

The directive influence order is:




655 (d)

$\text{Ti}^{4+}$ :  $3d^0$  and  $\text{Cu}^+$ :  $3d^{10}$  can not show  $d-d$ -transition and thus colourless.

656 (d)

$\text{Fe}^{3+}$  in  $[\text{Fe}(\text{CN})_6]^{3-}$  is:



657 (b)

The IUPAC name of  $[\text{Ni}(\text{PPh}_3)_2\text{Cl}_2]^{2+}$  is dichloro bis (triphenylphosphine) nickel (II).

658 (a)

It is neutral complex as it does not ionize in solution state.

659 (b)


Higher is the stability constant of ligand, lesser is its dissociation, more is its stability.

660 (c)

It is a fact.

661 (d)

Electronic configuration of  $\text{Co}^{3+}$  in  $[\text{CoF}_6]^{3-}$  is:



662 (c)

$\text{CH}_3$  is *o*- and *p*-directing gp.


663 (b)

$\text{Ag}^+$  has two coordination number forms complex with excess of  $\text{CN}^-$ , i.e.  $\text{Ag}(\text{CN})_2$

665 (c)

$-\text{C}-\text{N}$  group is called amino

while  $-\text{C}=\text{N}$  group is called imino




666 (c)

The coordination compound  $[\text{Co}(\text{en})_2\text{Cl}_2]\text{Cl}$  doesn't show ionization isomerism.




669 (d)

Nitrogen and oxygen are common donor atoms in ligands

670 (a)




671 (c)



This is Perkin's reaction.


672 (a)



It gives **five** ions in solution.




It gives **four** ions in solution.



It gives **three** ions in solution.


673 (a)



$\mu_1$  and  $\mu_2$  both for  $\text{NO}_2$  (electron withdrawing) and  $\text{NH}_2$  (electron releasing) gp. act in some direction.

674 (b)

Electronic configuration of Fe in  $\text{Fe}(\text{CO})_5$  is:



675 (b)

Turnbull's blue is  $\text{KFe}^{\text{II}}[\text{Fe}^{\text{III}}(\text{CN})_6]$ .

679 (c)

Each carbon in  $\text{C}_6\text{H}_6$  is  $sp^2$ -hybridized and thus, C—C bond is  $sp^2$ - $sp^2$ .

680 (b)

Change in composition of coordination sphere yield ionisation isomers


681 (a)

The IUPAC name of  $\text{K}_2[\text{Ni}(\text{CN})_4]$  is Potassium tetracyanonickelate (II).



683 (c)

It is a characteristic fact.

684 (b)




The oxidation number of Fe in  $[Fe(CN)_6]^{4-}$  is +2.




Since,  $CN^-$  is a strong field ligand, pairing occurs and the hybridisation of  $[Fe(CN)_6]^{4-}$  is  $d^2sp^3$  and structure is octahedral.

685 (c)



686 (d)

All the compounds in which there should be restricted rotation about a bond in the molecule, show geometrical isomerism. Oximes of the type  $CH_3 - CH = N - OH$ ,  $C_6H_5 - CH = N - OH$ ,  $C_6H_5 - C = N - OH$  and cyclic




Compound like show geometrical isomerism

687 (b)

The oxidation of aniline by  $K_2Cr_2O_7 + H_2SO_4$  (conc.) gives *p*-benzoquinone.

688 (b)



Here, *P* and *R* represent *meso*-compound

689 (b)

Friedel-Crafts reaction involves new C—C bond.

691 (a)

Rest all show less tendency to donate electron pair due to resonance.

692 (c)




Let oxidation state of Fe in



$$(+1 \times 3) + x + (-1 \times 6) = 0$$


$$\therefore x = +3$$




Let oxidation state of Fe in  $K_2[FeO_4] = x$

$$(+1 \times 2) + x + (4 \times -2) = 0$$

$$x = +6$$



Let oxidation state of Fe in




$$\therefore x + (-2) + 2 + (-2) = 0$$

$$x = +2$$

(d)  $[Fe(CN)_6]^{3-}$

Let oxidation state of Fe in



$$\therefore x + (6 \times -1) = -3$$

$$\therefore x = +3$$

$FeSO_4 \cdot (NH_4)_2SO_4$  has Fe in lowest oxidation state.

693 (a)

CO is a strong ligand, all the six electrons of the valence shell of Cr is paired and spin only magnetic moment = 0

694 (a)

Phenol is used in carbolic soaps.

695 (d)

Werner proposed theory for complex compounds to explain the structure and isomerism in them.

696 (a)

It is p-block element and thus, has no tendency to form complex.



697 (a)

In

$[CoF_6]^{3-}$  complex ion  $Co^{3+}$  is  $sp^3 d^2$  hybridized.  $F^-$  is weak ligand and cannot pair up the *d*-electrons so, complex is high spin. Due to four unpaired electrons it is highly paramagnetic.

698 (b)

$[Mabcd]$  type complexes exist in three isomeric forms.



699 (b)

Due to H-bonding.

700 (d)

A characteristic reaction of primary amine. This is carbylamine reaction.


701 (d)

The formula of given complex are as follows:


(a) Hexammineplatinum (IV) chloride




(b) Chloropentammine platinum (IV) chloride




(c) Dichlorotetrammine platinum (IV) chloride



(d) Trichlorotriammine platinum (IV) chloride



In aqueous solution the complex ionise is



Trichlorotriammine platinum (IV) 2 ions chloride gives the minimum number of ions in the solution.

Hence, it has the minimum electrical conductivity.


702 (c)

Diastereomers have different physical properties such as m. pt, b. pt solubilities

703 (b)

The decreasing order of priority of prefix in numbering the carbon chain of an organic compound is


Bromo > Chloro > Iodo




3-bromo-2-chloro-4-iodo hexane

705 (d)

The structure of the compound 2, 2'-bipyridine is



706 (d)



4-formyl-2-oxo-cyclohexane-1-carboxylic acid

**Note :** If a compound contains two or more substituents then numbering is done in such a way that the sum of the locants is the lowest

707 (c)

$-\text{N}_2\text{Cl}$  gp. Is reduced to  $-\text{H}$  by reducing agent  $\text{C}_2\text{H}_5\text{OH}/\text{Cu}$ .

708 (d)

All are the required facts for diethyl triamine.

709 (b)






711 (b)

When ligands are exchanged between metal atoms, coordination isomerism results. Hence,  $[\text{Co}(\text{NH}_3)_6][\text{Cr}(\text{CN})_6]$  and  $[\text{Cr}(\text{NH}_3)_6][\text{Co}(\text{CN})_6]$  re

presents coordination isomerism.

712 (c)

$\text{Co}^{3+}$  and  $\text{Pt}^{4+} = 6$  coordination number



Number of ionic species are same in the solution of both complexes, therefore their equimolar solutions will show same conductance.

713 (a)

Thiocyanato-N is the name when ligand SCN has electron pair donated by N-atom to metal.


715 (d)

Any side chain is oxidised to  $-\text{COOH}$ .

716 (d)


This is another reaction.

717 (c)



is heterocyclic compound


719 (c)



tetrammine copper (II) nitrate.




720 (c)

Nitroethane exhibits tautomerism



722 (c)

The electronic configuration  $\text{Pt}=[\text{Xe}]$



$\therefore$  No unpaired electron is present in  $[\text{Pt}(\text{CN})_4]^{2-}$  ion.

723 (c)

Let the oxidation number of cobalt is  $x$  in  $[\text{Co}(\text{CO})_4]$ .

$$1 + x + 0 = 0$$

$$x = -1$$

724 (b)

The IUPAC name of  $\text{Na}_3[\text{Co}(\text{NO}_2)_6]$  is sodium hexanitrocobaltate (III).


725 (a)

- $[\text{Cu}(\text{NH}_3)_4][\text{PtCl}_4]$
- $[\text{Cu}(\text{NH}_3)_3\text{Cl}] \cdot [\text{PtCl}_3(\text{NH}_3)]$
- $[\text{Cu}(\text{NH}_3)_2] \cdot [\text{PtCl}_2(\text{NH}_3)_2]$  *cis*
- $[\text{Cu}(\text{NH}_3)_2\text{Cl}_2][\text{PtCl}_2(\text{NH}_3)_2]$  *trans*
- $[\text{Cu}(\text{NH}_3)\text{Cl}_3] \cdot [\text{Pt}(\text{Cl})(\text{NH}_3)_3]$
- $[\text{Pt}(\text{NH}_3)_4] \cdot [\text{CuCl}_4]$

727 (c)


Tautomerism and functional isomerism is not possible together

732 (c)



733 (b)

This is Cannizzaro's reaction.



734 (a)




exhibit ring chain isomerism


735 (a)

Follow Werner's theory.

736 (b)

Faraday for the first time isolated  $\text{C}_6\text{H}_6$  from coal-tar.

737 (b)



Structural isomers (position) = 2

Stereo isomers = 4

738 (a)

According to Werner's theory, the primary valency of a metal is equal to the no. of charge on complex ion, *i. e.*, 3 on  $[\text{Fe}(\text{CN})_6]^{3-}$

739 (b)

The complex which contains 18 valence electrons, follows 18-electron rule.

(a) In  $[\text{V}(\text{CO})_5]$

The number of valence electrons  
=  $5 + (2 \times 5)$

$$= 15 e^-$$

(b) In  $[\text{Fe}(\text{NH}_3)_6]^{2+}$ ,

The number of valence electrons

$$= 6 + (6 \times 2) = 6 + 12 = 18 e^-$$

(c) In  $[\text{Ni}(\text{CO})_6]$ ,

The number of valence electrons =  $10 + (2 \times 6) = 22 e^-$

(d) In  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$ ,

The number of valence electrons =  $5 + (6 \times 2) = 17 e^-$

Thus, only  $[\text{Fe}(\text{NH}_3)_6]^{2+}$  follows 18-electron rule.

740 (d)

One mole of *X* gave depression corresponding to 2 moles of particles, *i. e.*, on ionisation *X* gives 2 moles of ions, thus it contains only 1 ion outside the coordination sphere and its structural formula is  $[\text{Cr}(\text{H}_2\text{O})_4\text{Br}_2]\text{Cl} \cdot \text{H}_2\text{O}$  while *Y* gives 3 moles of ions, thus it contains two ions outside the coordination sphere and its structural formula is  $[\text{Cr}(\text{H}_2\text{O})_5\text{Cl}]\text{Br}_2$


742 (d)

Both represent only one molecule and no isomerism.

743 (a)

Haemoglobin is porphyrin complex of ferrous iron being coordinated to four nitrogen atoms and additionally coordinated to a water reversible by a molecule. The water molecule appears to be replaceable reversible by a molecule of oxygen to give oxyhaemoglobin.  $\text{Fe}^{2+}$  is diamagnetic due to strong field ligands.

746 (d)




—CN and —COOH gp. at *p*-position facilitate replacement of Cl gp. by  $\text{S}_\text{N}$  reactions to show normal  $\text{S}_\text{N}$  reactions.

747 (b)

$(\text{NH}_4)_2\text{S}_x$  brings in selective reduction of one of the two —NO<sub>2</sub> group at *m*-position.

748 (c)

If the highest priority groups on two carbon atoms of the double bond are on the opposite side, the configuration is *E*. (Entgegen)



749 (c)

Cyclopentane possess  $0^\circ 44'$  angle strain which is minimum

750 (c)

$[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$  has four unpaired electrons

751 (b)

$[\text{Pt}(\text{NH}_3)_6]\text{Cl}_4 \rightleftharpoons \text{Pt}(\text{NH}_3)_6 + 4\text{Cl}^-$


$\text{Ag}^+ + \text{Cl}^- \rightarrow \text{AgCl} \downarrow$   
White ppt.

752 (a)

$\text{CH}_3(\text{CH}_2)_3 \cdot \text{NH}_2 \xrightarrow[\text{CHCl}_3]{\text{KOH alc.}} \text{CH}_3(\text{CH}_2)_3\text{NC} + 3\text{KCl} + 3\text{H}_2\text{O}$

$\text{CH}_3 - \text{C} \equiv \text{CH} + \text{Amm} \cdot \text{AgNO}_3 \rightarrow \text{CH}_3\text{C} \equiv \text{C} \cdot \text{Ag} + \text{HNO}_3$

$\text{CH}_3 \cdot \text{CH}_2\text{COOCH}_3 + \text{NaOH} \xrightarrow{\Delta} \text{CH}_3\text{CH}_2\text{COONa} + \text{CH}_3\text{OH}$



2° alcohol

→ Cloudiness appears within 5 minute.


754 (b)

Configuration of  $\text{Mn}^{2+}$  is

$[\text{Ar}]3d^5$

According to CFSE (crystal field stabilisation energy), in excited state of  $\text{Mn}^{2+}$  ion, three electrons go in  $t_{2g}$  level ( $d_{xy}$ ,  $d_{yz}$  and  $d_{zx}$ ) and two electrons go in  $e_g$  level ( $d_{z^2}$  and  $D_{x^2-y^2}$ ).

755 (d)



756 (c)

eg,  $\text{Fe}(\text{CO})_5$ ,  $\text{Ni}(\text{CO})_4$ , etc.,

757 (b)

Follow IUPAC rules.

758 (b)

$[\text{Co}(\text{NH}_3)_6][\text{Cr}(\text{C}_2\text{O}_4)]_3$  its IUPAC name is hexa amine cobalt (III) tris (oxalato) chromate (III).



759 (b)

Gammexane is  $\text{C}_6\text{H}_6\text{Cl}_6$ .



760 (b)

1. In  $\text{Ni}(\text{CO})_4$ , nickel is  $sp^3$ -hybridised because in

it oxidation state of Ni is zero. So, configuration of  $_{28}\text{Ni} = 1s^2 2s^2 2p^6, 3s^2 3p^6 3d^8, 4s^2$





2. In  $[\text{Ni}(\text{CN})_4]^{2-}$ , nickel is present as  $\text{Ni}^{2+}$ , so its configuration =  $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^8$



is strong field ligand, hence it makes  $\text{Ni}^{2+}$  electrons to be paired up.

In  $[\text{NiCl}_4]^{2-}$  species, nickel is present as  $\text{Ni}^{2+}$ , so its configuration =  $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^8$



$\text{Cl}^-$

is weak field ligand, hence  $\text{Ni}^{2+}$  electrons are not paired.

761 (d)

$\text{Ti}^{4+} : 3d^0$


$\text{Cr}^{3+} : 3d^3$  Completely filled or empty  $d$ -orbitals are colourless.



$\text{Zn}^{2+} : 3d^{10}$

$\text{Sc}^{3+} : 3d^0$

762 (b)

Possible isomers are as follows :





Here, only (I), (II) and (III) are chain isomers

763 (a)

2,2-dinitrodiphenyl or 4,4-dinitrodiphenyl is formed.

764 (d)


We have that by breaking two bond on the chiral centre configuration changes

765 (c)

Use of oleum ( $\text{H}_2\text{SO}_4 + \text{SO}_3$ ) produces inclusion of  $-\text{SO}_3\text{H}$  gp. in  $\text{C}_6\text{H}_6$  ring.

766 (a)

$\text{NO}_2^-$  can participate in linkage isomerism because it may be bonded to metal through nitrogen or through oxygen.



768 (c)

Cuprammonium salt,  $[\text{Cu}(\text{NH}_3)_4]\text{SO}_4$   
 $[\text{Cu}(\text{NH}_3)_4]\text{SO}_4 \rightleftharpoons [\text{Cu}(\text{NH}_3)_4]^{2+} + \text{SO}_4^{2-}$   
 So, it will give two ions in water


769 (b)

(a) Shows tautomerism since aldehydes are more stable than vinyl alcohols




(b) Does not show tautomerism because it does not have hydrogens at  $\alpha$ -positions

(c) Shows tautomerism because enol form is stabilized by H-bonding



(d) Shows tautomerism because enol form is stabilized by aromatic character




771 (a)

Main fractions of coal-tar and the compounds present there in are:

	Main fraction	Temp. range	Chief constituents
1.	Light oil or crude	80-170°C	Benzene, toluene, xylenes, etc.
2.	naphtha	170-230°C	Phenol, naphthalene, pyridine, etc.
3.	Middle oil or carbolic acid	230-270°C	Cresols, naphthalene quinolone, etc.
4.	Heavy oil or creosote oil	270-360°C	Anthracene, phenanthrene, etc.
5.	Green oil or anthracene oil	Residue	90-94% of carbon
	Pitch		

772 (b)

Octahedral complexes containing three bidentate ligands shows optical isomerism. If A is a bidentate ligand then complex of type  $\text{MA}_3$  show optical isomerism.



773 (a)

Cl atom attached in side chain behaves as aliphatic in nature.

775 (d)

Due to electron deficient molecule it accepts lone pair of electron to produce electrophile.  
 $\text{AlCl}_3 + \text{Cl}_2 \rightarrow \text{AlCl}_4^- + \text{Cl}^+$

777 (b)

*o*-nitrophenol has intramolecular H-bonding.

778 (c)


IUPAC name is tetraammine nickel (II) - tetrachloronickelate (II).

779 (b)

Ligands are electron pair donor.


781 (d)

*Cis*- form of  $[\text{Co}(\text{en})_2(\text{NH}_3)_2]^{3+}$  is optically active.




784 (d)

After two interchanges at each of the two chiral carbon atoms in second structure in such a way that CH<sub>3</sub> group is held vertically upward and C<sub>2</sub>H<sub>5</sub> group vertically downward, we get first structure



Thus, the two structures are identical

785 (c)




cation anion

Oxidation state of Fe in anion = +3

Thus, it is potassium hexacyanoferrate (III).


786 (b)

In  $[\text{Zn}(\text{NH}_3)_6]^{3+}$ , Zn exists as  $\text{Zn}^{2+}$



$\text{Zn}^{2+}: 3d^{10}$ ; Thus, no unpaired electron but it is outer orbital complex.



In  $[\text{Co}(\text{NH}_3)_6]^{3+}$ , Co exists as  $\text{Co}^{3+}$



$\text{Co}^{3+}: 3d^6$ ; It is  $d^2sp^3$  inner orbital complex with 3 electron paired in  $3d$ .

787 (a)

In  $[\text{CoCl}_4]^{2-}$  ion, central metal atom i.e., cobalt is in +2 oxidation state. Hence,



Hence, number of unpaired electrons is three as Cl is weak ligand.

788 (c)

Coordination number is the number of ligands in the coordination sphere. Hence, the coordination number of cobalt ion in  $[\text{Co}(\text{H}_2\text{O})_4\text{SO}_3]\text{Cl}$  is 5.

Let the oxidation number of Co is  $x$ .


$$x + 4(0) + (-2) + (-1) = 0$$

$$x + 0 - 2 - 1 = 0$$

$$x = 3$$

Number of unpaired electrons in  $d$ -orbital are 4 because  $\text{H}_2\text{O}$  is a weak ligand and therefore, pairing of  $d$ -electrons is not possible.

789 (b)



790 (c)

$-\text{SO}_3\text{H}$  is water soluble.

791 (c)

$$+1 \times 4 + x - 1 \times 4 = 0$$

$$4 + x - 4 = 0 \Rightarrow x = 0 \text{ for Ni}$$

792 (c)

Follow IUPAC nomenclature.


793 (b)

Due to bitter almond smell. It is



795 (c)

$\text{C}_6\text{H}_5\text{COCH}_3$  acetophenone is a mixed ketone having one alkyl and other phenyl gp. attached



796 (d)

These are the facts about transition metal atoms to act as central atom.

798 (d)


Since, hybridization is  $dsp^2$  so, it is square planar

799 (b)


Metal-carbon bond in metal carbonyls has  $\sigma$  as well as  $\pi$  characters.

800 (d)


The electronic configuration of Ni in  $[\text{Ni}(\text{CN})_4]^{2-}$ ,  $[\text{Ni}(\text{Cl})_4]^{2-}$  and  $\text{Ni}(\text{CO})_4$  are:  $\text{Ni}^{2+}$  in  $[\text{Ni}(\text{CN})_4]^{2-}$ :



$\text{Ni}^{2+}$  in  $[\text{Ni}(\text{Cl})_4]^{2-}$ :




Ni in  $[\text{Ni}(\text{CO})_4]$ :





801 (c)

Both produce different ions in solution state:



803 (c)


The configuration in which  $-\text{OH}$  group is on right side, H-atom is on left side,  $-\text{CHO}$  group is on upper side and  $\text{CH}_2\text{OH}$  is on lower side found in Fischer projection known as D-configuration



804 (a)

Linkage isomerism is shown by those complexes which have an ambidentate ligand such as  $\text{NO}_2^-$ ,  $\text{CN}^-$  and  $\text{SCN}^-$  etc. In  $[\text{Fe}(\text{NO}_2)_3\text{Cl}_3]$ , N is donor atom.


805 (c)



2,6-dimethyl hepta-2,5-dienoic acid

806 (d)

$\text{NaHCO}_3$  reacts with acids to give  $\text{CO}_2$  from  $\text{HCO}_3^-$  ion.




807 (a)

The species within the coordination sphere does not ionize.

808 (c)

The reverse of enolic contents of compound is ketonic contents. Thus, the correct order of ketonic contents are



809 (b)

The anhydrous complexes of  $\text{Cu}_2^{2+}$  do not involve  $d-d$  transition and are thus, colourless.

810 (d)

The valence of C-atom of ring is 5 at two methyl gp. attachment.


812 (b)

Aniline is basic and thus, reacts with acid.

814 (b)

The Riemer-Tiemann reaction is followed by dichloro carbene mechanism.

815 (c)



816 (b)

$\text{Mn}^{2+}$  will have half filled more stable  $d^5$  configuration and without distributing it an outer orbital complex can be formed

817 (a)

	Hybridization	Unpaired electrons	Magnetic moment
1. $[\text{Co}(\text{CN})_6]^{3-}$	$d^2sp^3$	0	0
2. $[\text{Fe}(\text{CN})_6]^{3-}$	$d^2sp^3$	1	$\sqrt{3}$ BM
3. $[\text{Mn}(\text{CN})_6]^{3-}$	$d^2sp^3$	2	$\sqrt{8}$ BM
4. $[\text{Cr}(\text{CN})_6]^{3-}$	$d^2sp^3$	3	$\sqrt{15}$ BM

Thus, least paramagnetism is in (a).

818 (d)

The primary valencies of Ni, in the complexes  $[\text{Ni}(\text{Cl}_4)]^{2-}$  and  $[\text{Ni}(\text{CN})_4]^{2-}$  is same *i.e.*, (+II). Primary valencies are those valencies which a metal exhibits in the formation of its simple salt, these are non-directional. It is also referred as oxidation state of central metal atom.

819 (c)

The central metal ion is  $\text{Fe}^{3+}$  and  $\text{C}_2\text{O}_4^{2-}$  is negative bi-dentate ligand which forms more stable complex than neutral or mono-dentate ligand.

821 (b)


In  $\pi$ -complex, organic ligands use their  $\pi$  system to bond with metal, *e.g.*, ferrocene.

823 (c)

$\text{CN}^-$  ions act both as reducing agent as well as good complexing agent


824 (a)

The order of *meta* directing, gp. is:



825 (a)

$\text{CH}_3\text{COC}_3\text{H}_7$  can exhibit metamerism



826 (d)

It has coordination no. six and thus, octahedral or  $sp^3d^2$  hybridization.

828 (d)

Let the oxidation state of iron in  $K_4[Fe(CN)_6]$  is  $x$ .

$$4(+1) + x + 6(-1) = 0$$

$$4 + x - 6 = 0$$

$$x = +2$$

829 (d)

These are the methods to test complex formation.


830 (b)

EAN of Fe in  $K_3Fe(CN)_6$  is:  $26 - 3 + 2 \times 6 = 35$ , *i.e.*, not 36 the next inert gas.

833 (a)

Grignard reagent is a  $\sigma$ -bonded organometallic compound because all the bonds present in the reagent are single bonds.

835 (b)




836 (d)


Thymol is 3-hydroxy-1-isopropyl-4-methyl benzene ( $C_{10}H_{14}O$ ), a white crystalline phenol derivative, has smell of thyme, occurs in many essential oils used as fragrant material as well as mild antiseptic.

837 (d)

Ferrocyanide ion  $[Fe(CN)_6]^{4-}$  is diamagnetic in nature hence  $K_4[Fe(CN)_6]$  complex has zero magnetic moment.



$d^2 sp^3$  hybridisation



838 (a)

CO ligand has zero oxidation state, that is why  $[Ni(CO)_4]$  is a zero valent metal complex


840 (c)

A characteristic; follow ligand field theory.

842 (b)

Central ion is  $Cd^{2+}$  and ligand is  $CN^-$ .

843 (b)



In compound two chiral carbon atoms are present

844 (c)

9 mole or  $9 \times 22.4$  litre of  $C_2H_2$  are needed.

845 (b)

IUPAC name of sodium nitroprusside  $Na_2[Fe(CN)_5NO]$  is sodium pentacyanonitrosyl ferrate (III) because in it NO is neutral ligand and the oxidation number of Fe is III, which is calculated as

$$2 \times ON \text{ of } Na + ON \text{ of } Fe + 5 \times ON \text{ of } CN$$


$$+ 1 \times ON \text{ of } NO = 0$$

$$2 \times (+1) + ON \text{ of } Fe + 5 \times (-1) + 1 \times 0 = 0$$


$$\Rightarrow ON \text{ of } Fe = 5 - 2 = +3$$

846 (b)


The electronic configuration of Ni in  $[Ni(CN)_4]^{2-}$ ,  $[Ni(Cl)_4]^{2-}$  and  $Ni(CO)_4$  are :  $Ni^{2+}$  in  $[Ni(CN)_4]^{2-}$  :



$Ni^{2+}$  in  $[Ni(Cl)_4]^{2-}$  :



$Ni^{2+}$  in  $[Ni(CO)_4]$  :




848 (d)

When cations have same charge but number of  $d$ -electrons are different then the stability (or CFSE) decreases with increase in the number of  $d$ -electrons. Therefore, the correct order is  $Fe^{2+} > Co^{2+} > Ni^{2+} > Cu^{2+}$

849 (a)

Oxidation state of Ni in  $Ni(CO)_4$  is zero.



$sp^3$  - hybridisation

CO is a strong ligand. It causes pairing of electrons. Hence, there is no unpaired electrons in  $Ni(CO)_4$ .

850 (b)

Ox. no. of Cr is calculated as:


$$3 \times 1 + a + 1 \times (-2) + 4 \times (-1) = 0; \therefore a = +3$$

853 (c)

$\text{CrO}_3$  or  $\text{CrO}_2\text{Cl}_2$  and a mixture of  $\text{K}_2\text{Cr}_2\text{O}_7 + \text{H}_2\text{SO}_4 + \text{NaCl}$  can also be used.

854 (b)

Butane-2 exhibit geometrical (*cis*, *trans*) isomerism



855 (a)

Ferrocene is a  $\pi$  complex  $\text{Fe}(\eta^5 - \text{C}_5\text{H}_5)_2$ .

856 (a)

Ni in  $[\text{Ni}(\text{NH}_3)_6]^{2+}$  has  $sp^3d^2$  (outer complex) having octahedral geometry.

857 (d)

These examples are ionisation isomers because of chloride and sulphate ions.


859 (b)

When ligands are arranged in ascending order of crystal field splitting energy,  $\Delta$ , they produce a spectrochemical series.

In comparison to  $\text{H}_2\text{O}$ , Cl is strong  $\sigma$ -donor and good  $\pi$ -acceptor, therefore it is a strong ligand than  $\text{H}_2\text{O}$ . Hence, in the spectrochemical series Cl is above than water.

860 (b)

On ionization different species are formed to show ionization isomerism:



861 (a)

$\text{Pt}^{2+}$  has square planar complexes with coordination number four.

862 (c)

In  $\text{Hg}[\text{Co}(\text{SCN})_4]$ , Co exists as  $\text{Co}^{2+}$

$${}_{27}\text{Co} : 3d^7, 4s^2$$


$${}_{27}\text{Co}^{2+} : 3d^7$$

$\text{SCN}^-$  a strong field ligand provides four electron to pair to show  $sp^3$ -hybridisation in  $[\text{Co}(\text{SCN})_4]^{2-}$  and thus three unpaired electrons exists on  $\text{Co}^{2+}$ .

$$\therefore \text{Magnetic moment} = \sqrt{n(n+2)} = \sqrt{3(3+2)} = \sqrt{15}$$

863 (d)

But-2-ene and but-1-yne do not have same molecular formula, thus are not isomers



864 (b)

$${}_{27}\text{Co} \rightarrow [\text{Ar}]3d^7 4s^2$$

$$\text{Co}^{2+} \rightarrow 3d^7 4s^0$$



Number of unpaired electrons = 3

865 (a)

NO in iron complexes has ox. no. equal to one.

$$a + 5 \times 0 + 1 - 2 = 0, \therefore a = +1$$

868 (c)

Due to the presence of phenolic gp.

869 (b)


$\text{Ni}(\text{CO})_4$  has a O.N. zero for Ni

870 (d)


Because it is a polydentate ligand which binds the central atom nickel forming a ring structure

871 (c)


Carbonyl compounds react with Grignard reagent following nucleophilic addition. More +ve is charge on  $\text{C}^+$  centre of carbonyl gp., easier is nucleophile attack.



(Positive charge on  $\text{C}^+$  is dispersed due to +I.E. of  $\text{CH}_3$  gp.)



(Positive charge on  $\text{C}^+$  is dispersed more due to +I.E. of two  $\text{CH}_3$  gp.)



(Positive charge on  $\text{C}^+$  is intensified due to -I.E. of  $\text{C}_6\text{H}_5$  gps.)

But in (III) conjugation of  $>\text{CO}$  gp. with  $\pi$  system

of benzene nucleus following resonance deactivates  $\text{C}^+$  centre to attack by nucleophile.

Resonance effect overpowers over -I.E. of  $\text{C}_6\text{H}_5$  gp.

872 (d)

$[\text{Cr}(\text{SCN})_2(\text{NH}_3)_4]^+$  shows geometrical (or *cis-trans*) and linkage isomerism.

873 (c)




Thus, coordination no. of Cu is four.

876 (b)

The IUPAC name of the compound  $[\text{CuCl}_2(\text{CH}_3\text{NH}_2)_2]$  is dichloro bis-(methyl amine) copper (II).


877 (b)



2-(1-cyclobutenyl)-1-hexane

879 (b)

Electronic configuration of  $[\text{FeF}_6]^{3-}$  is:



×× Electron pair donated by  $\text{F}^-$


880 (d)

Effective atomic no. (EAN) = at. No. of central atom - oxidation state +  $2 \times$  (no. of ligands) =  $28 - 0 + 2 \times 4 = 36$

$\text{EAN} = 26 - 3 + 2 \times (6) = 35$

882 (d)

$[\text{Pt}^{\text{II}}(\text{NH}_3)_2\text{Cl}_2]$  shows geometrical isomerism.



884 (b)

Staggered and eclipsed conformers cannot be physically separated because the energy difference between them is so small that they most readily interconvert at room temperature

885 (b)

A species or group of atoms can act as ligand only when it carries an unshared pair, *i.e.*, lone pair of electrons.

886 (c)

Follow IUPAC rules.

888 (d)


$\text{NH}_2 \cdot \text{NH}_2$  serves as monodentate as well as bridging ligand because a 3-membered ring will be too strained to be stable.

889 (b)

TNT mixed with  $\text{NH}_4\text{NO}_3$  gives explosive material.


890 (d)

During debromination, *meso*-dibromobutane form *trans*-2-butene



891 (a)

The IUPAC name of compound is



Potassium ammine dicyano dioxo peroxy chromate (VI)

893 (a)

It can show ionization isomerism:  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+$  and  $[\text{Co}(\text{NH}_3)_4\text{Cl}]^{2+}\text{Cl}^-$ .

894 (b)

Replacement of H-atom of ring usually takes place following  $\text{S}_\text{E}$  reaction mechanism.

895 (b)

Follow mechanism of Reimer-Tiemann reaction.

896 (a)

It is 2,4,6-trinitrophenol.

897 (a)

Follow mechanism of cannizzaro's reaction.


898 (b)

Ag in Tollens' reagent exists as  $\text{Ag}_2\text{O}$

$$2 \times a + 1 \times (-2) = 0$$

$$\therefore a = +1$$

899 (b)




Electronic configuration of Fe =  $[\text{Ar}]3d^64s^2$

Electronic configuration of  $\text{Fe}^{3+}$  =  $[\text{Ar}]3d^5$

Number of ligand (Coordination number) = 6

Nature of ligand in strong field



Hybridisation of Fe is  $d^2sp^3$


900 (d)

Butan-2-one 0

||

$\text{CH}_3 - \text{C} - \text{CH}_2 - \text{CH}_3$  is not isomeric with diethyl ether  $\text{CH}_3\text{CH}_2 - \text{O} - \text{CH}_2 - \text{CH}_3$ . Because both are differing in molecular formula

902 (d)



$$\text{CFSE} = 3\lambda(-0.4)\Delta_0 + 0.6(\Delta_0) = 0.6 \Delta_0$$

905 (b)

Only carbonyl compounds show this test.

906 (c)

Presence of *meta* directing gp. Deactivates ring for Friedel-Crafts reaction.

907 (c)

$\text{Ni}(\text{CO})_4$  and  $\text{Ni}(\text{PPh}_3)_2\text{Cl}_2$  are tetrahedral in geometrical shape, because coordination number of Ni is four in both cases.

908 (c)

Geometrical isomers (*cis* and *trans*) and linkage isomers ( $-\text{SCN}$  and  $-\text{CNS}$ ).

909 (a)

The absorption of energy of the observation of colour in a complex transition compounds depends on the charge of the metal ion and the nature of the ligands attached. The same metal ion with different ligands shows different absorption depending upon the type of ligand. The presence of weak field ligands make the central metal ion to absorb low energies *ie*, of higher wavelength. The field strength of ligands can be obtained from spectrochemical series, *ie*,  
(weak field)  $\text{I}^- < \text{Br}^- < \text{S}^{2-} < \text{Cl}^- < \text{NO}_3^- < \text{F}^- < \text{OH}^- < \text{H}_2\text{O} < \text{NH}_3 < \text{NO}_2^- < \text{CN}^- < \text{CO}$  (strong field)

911 (d)


The total number of monodentate ligands attached to the central metal is known as coordination number. Hence, in  $[\text{Cu}(\text{H}_2\text{O})_4]^{2+}$  coordination number of Cu atoms is 4.

912 (c)

Prefixes and suffixes are written before and after the root word respectively and not before and after the compound

The IUPAC name of a compound is written as single word

913 (c)




914 (c)

$[\text{Co}(\text{NH}_3)_3\text{Cl}_3]$  does not have optical isomers

because it is of formula  $\text{MA}_3\text{B}_3$  which does not show optical isomerism

915 (b)

Ethylenediamine is a bidentate ligand.



916 (b)


Electron withdrawing nature or  $-IE$  increases the activation of ring more effectively, however resonance opposes inductive effect for attachment at *o*- and *p*-position and hence, makes less deactivation for *o*- and *p*-positions.

918 (d)

All involves  $sp^3$ -hybridization.

919 (b)

Chaisen condensation involves condensation of benzaldehyde with aliphatic aldehydes or ketones having two  $\alpha$ -H-atoms, *e. g.*,



920 (d)

All are transition elements with *d*-orbitals not filled to capacity.


921 (a)

$\text{Ag}(\text{II})$  has a  $d^9$ -configuration and must contain unpaired electron but  $\text{AgO}$  is diamagnetic because  $\text{AgO}$  does not contain  $\text{Ag}(\text{II})$  but is a mixed oxide of  $\text{Ag}^{\text{I}}\text{Ag}^{\text{III}}\text{O}_2$ .

922 (c)

Friedel-Crafts reaction involves alkylation or acylation in benzene nucleus using alkylating or acylation reagents in presence of anhy.  $\text{AlCl}_3$ .

924 (b)




925 (a)

$\text{CoCl}_2$  is a weak Lewis acid, reacting with chloride ions to produce salt containing the tetrahedral  $[\text{CoCl}_4]^{2-}$  ion.  $\text{CoCl}_2$  is blue when anhydrous, and a deep magenta colour when hydrated, for this reason it is widely used as an indicator for water

926 (c)

Octahedral complexes of the type  $\text{MA}_5\text{B}$  do not show geometrical isomerism.


927 (b)



928 (d)

In this compound  $\begin{array}{c} \text{—C—NH}_2 \\ || \\ \text{O} \end{array}$  is principal functional group

While  $\text{—CHO}$  is substituent group, hence




5-formyl-2-methyl pent-3-en-1-amide

929 (b)

$\text{C}_6\text{H}_5\text{CH}_2\text{NH}_2$  has least negative inductive effect and thus shows more basic nature.

930 (d)



4-ethyl-3-methyl heptane

**Note :** The prefix in a compound should be arranged in alphabetical order

931 (d)

All possess lesser number of unpaired electrons.


932 (d)

A monodentate ligand has one donor site available for coordination.

933 (a)

It is benzene.

934 (c)




935 (c)

Phenol is weak acid.

936 (b)

The coordination number in  $[\text{FeF}_6]^{3-}$  is 6, hence it is an octahedral complex

937 (d)



938 (a)

Main fractions of coal-tar and the compounds present there in are:

	Main fraction	Temp. range	Chief constituents
1.	Light oil or crude	80-170°C	Benzene, toluene, xylenes, etc.
2.	naphtha	170-230°C	Phenol, naphthalene, pyridine, etc.
3.	Middle oil or carbolic acid	230-270°C	Cresols, naphthalene quinolone, etc.
4.	Heavy oil or creosote oil		
5.	Green oil or anthracene		

oil Pitch	270-360°C	Anthracene, phenanthrene, etc. 90-94% of carbon
	Residue	

939 (c)


The coordination number (C.N.) of a metal atom in a complex is the total number of bonds formed by metal with ligands.

In case of tetrahedral complexes the number of bonds formed between metal and ligand is four. So, coordination number is also four.

940 (b)

Due to resonance bond length become identical and is 1.40 Å. Whereas in alkane C—C bond is 1.54 Å and in alkene it is 1.34 Å.

941 (d)



944 (c)

Electrophilic substitution occurs at electron rich centres usually at *o*- and *p*-positions. The ring attached with  $\text{—NH}$  will develop more electron density at *o*- and *p*-positions. Since *o*-position is blocked, thus electrophile will attach at *p*-position.

945 (b)

Follow text.


946 (a)

In the formation of  $d^2sp^3$  hybrid orbitals, two  $(n-1)d$  orbitals of  $e_g$  set *i.e.*,  $(n-1)d_{z^2}$  and  $(n-1)d_{x^2-y^2}$  orbitals, one  $ns$  and three  $np$  ( $np_x$ ,  $np_y$  and  $np_z$ ) orbitals combine together and form six  $d^2sp^3$  hybrid orbitals.


947 (a)

$$\text{EAN of Fe} = 26 - 2 + 6 \times 2 = 36.$$

948 (a)



949 (b)



950 (d)

All are characteristic tastes for phenol.

952 (c)

N cannot have more than 8 elements in its valence shell.

953 (b)

Tautomeric structures of a molecule are not the resonating structures of the molecule

954 (a)

2, 4, 6-trinitrophenol is strong acid than acetic acid but phenol is less acidic than acetic acid.

955 (a)

Phenol being weak acid does not react with aq.  $\text{NaHCO}_3$ .

956 (c)

If two atoms directly attached to the double bond have the same atomic number, then the relative priority of the groups is determined by a similar comparison of the atomic numbers of the next elements in the groups. Thus, the preference order of given group is  $\text{OCH}_3 > \text{OH} > \text{COOH} > \text{CHO}$

957 (a)


Follow IUPAC name.

958 (a)

It possesses  $d^8$  configuration of  $\text{Ni}^{2+}$ .

959 (d)

There are four structural isomers possible for diphenyl methane when one H-atom is replaced by a Cl-atom




960 (c)

$\text{CN}^-$  is a better complexing agent ( $C$ ) as well as a reducing agent ( $A$ )


Thus, properties ( $A$ ) and ( $C$ ) are shown.

Property ( $C$ ):  $\text{Ni}^{2+} + 4\text{CN}^- \rightarrow [\text{Ni}(\text{CN})_4]^{2-}$

Property ( $A$ ):



961 (c)

In  the double bonds are *trans* and *cis*. The first and third bonds are identical

963 (a)

*p*-nitrophenol is most acidic (among these) as it has electron withdrawing  $-\text{NO}_2$  gp.


965 (c)

Geometrical isomerism is found in compounds having coordination no. 4 (square planar and not tetrahedral shape) as well as coordination no. 6. Coordination no. 4 (square planar)




Showing geometrical isomerism

Coordination no. 6 (octahedral)



Showing geometrical isomerism.



966 (d)

Follow crystal field theory.

967 (d)

Due to resonance  $\text{C}_6\text{H}_6$  is stabilized and normal addition reactions (except addition of  $\text{H}_2$ ,  $\text{Cl}_2$  and ozonolysis) are not observed in  $\text{C}_6\text{H}_6$ .


969 (b)

EAN of Pt in  $[\text{PtCl}_6]^{2-} = 86$

971 (b)

$\text{CuCl} + \text{NH}_3 \rightarrow [\text{Cu}(\text{NH}_3)_2]\text{Cl}$ . The coordination no. = No. of ligands attached.

972 (a)



Longest system : 1,2,3,4,5,6

Next longest system : 6,7,8,1

Shortest system : 1,6

The IUPAC name of compound is 8-methyl bicyclo [4,3,0] nonane

973 (b)

Aniline on condensation with aromatic aldehyde gives Schiff's base.

974 (c)

Organometallic compounds are those compounds in which metal is directly attached to the carbon atom. In sodium ethoxide, sodium attached to oxygen atom, hence it is not an organometallic compound.

975 (d)


(i)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$

and

$\text{CH}_3\text{CH}_2\text{CH}_2 - \text{CH} - \text{CH}_3$

|  
OH

are Position isomers





Due the presence of asymmetry, optical isomerism is possible

(iii)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$  and  $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_2\text{CH}_3$  are functional isomerism

976 (c)

$\text{Co}^{3+}$  and  $\text{Pt}^{4+}$  have 6 coordination number.

$\text{CoCl}_3 \cdot 6\text{NH}_3$  and  $\text{PtCl}_4 \cdot 5\text{NH}_3$



Number of ionic species are same in the solution of both complexes, therefore their equimolar solutions will show same conductance.

977 (a)

Organometallic compounds are those compounds in which carbon atom is directly linked to metal.

But in sodium ethoxide as oxygen is attached to attached to sodium metal so, it is not a organometallic compound


978 (d)

$[\text{Cu}(\text{H}_2\text{O}_4)]\text{SO}_4 \cdot \text{H}_2\text{O}$  coordination number of Cu is 4.

979 (b)

Sodium nitroprusside is  $\text{Na}_2[\text{Fe}(\text{CN})_5\text{NO}]$ .


980 (d)



2-amino-3-hydroxy propanoic acid

981 (a)

$[\text{CoF}_6]^{3-}$  is an outer orbital complex ion. It involves outer orbital hybridisation. It has  $sp^3d^2$ -hybridisation because  $\text{F}^-$  is a weak ligand.



982 (b)

Effective atomic no. (EAN) = at. No. of central atom - oxidation state +  $2 \times$  (no. of ligands) =

$$28 - 0 + 2 \times 4 = 36$$

$$\text{EAN} = 28 - 2 + 2 \times 4 = 34.$$

983 (b)

Due to restricted rotation about the carbon-carbon single bond joining the two phenyl groups, the molecule as a whole is chiral and thus shows optical isomerism

984 (b)

Any side chain is oxidised to  $\text{COOH}$  gp.

985 (d)

These are facts about glycinate ligand.

986 (c)

According to Werner's theory, only those ions are precipitated which are attached to the metal atoms with ionic bonds and are present outside the coordination sphere.

987 (d)

$sp^2$ -hybridization leads to planar hexagonal shape.


988 (a)

Coordination isomerism is caused by interchange of ligands with the metal atoms.


989 (c)

Chlorophyll are green pigment in plant and contain magnesium instead of calcium


991 (a)



992 (a)






993 (d)



Tetraammine dichloro ferrum III chloride.

994 (b)



Nickel has two unpaired electrons and geometry is tetrahedral due to  $sp^3$  hybridisation.

995 (c)

It is a test for  $-\text{NH}_2$  gp attached on benzene nucleus following diazotisation and coupling reaction.



101 (d)



2 Ethane has an infinite number of conformation but staggered and eclipsed are preferred. Ethane molecule would exist in the staggered conformation due to its minimum energy and maximum stability

101 (a)



4 Follow IUPAC rules

101 (d)



5 The complex in which  $nd$  orbitals are used in hybridisation, are called outer orbital complex.





27.



28.




29.



101 (d)


6 In each case aromatisation leads to formation of  $\text{C}_6\text{H}_6$ .

101 (a)



101 (b)

8  $\beta$ -keto acids undergo decarboxylation most easily on heating.



101 (a)

9 Ni in  $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$  has two unpaired electrons in it.

102 (d)

4 An electron attracting group ( $-\text{NO}_2$ ) disperses the negative charge on phenoxide ion and thus, makes it more stable or increases the acidic character of phenol. The substitution is more effective at  $p$ -position than in the  $m$ -position as the former involves a resonating structure bearing negative charge on the carbon attached to the electron withdrawing group. Also presence of electron repelling gp. ( $-\text{CH}_3$ ) intensifies the negative charge on phenoxide ion and thus, makes phenol less acidic.

102 (d)



102 (c)

9 Let the ON of Pt in  $[\text{Pt}(\text{NH}_3)_5\text{Cl}]\text{Cl}_3$  is  $x$ .

$$x + 5 \times (0) - 1 - 3 = 0$$

$$x - 4 = 0$$


$$x = +4$$

103 (a)

0 Organometallic compounds have carbon-metal bond, hence  $\text{CH}_3\text{MgI}$ , tetraethyl tin and  $\text{KC}_4\text{H}_9$  are organometallic compounds while  $\text{C}_2\text{H}_5\text{ONa}$  is not an organometallic compound due to absence of carbon-metal bond.

103 (b)

1 Optical isomers of a compound which are non-superimposable but related to each other as an object and its mirror are called enantiomers





103 (d)

2 All involve  $dsp^2$ -hybridization.

103 (a)

3 *Cis*- isomer of  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$  is used as an anticancer drugs for treating several type of malignant tumours when it is inject into the blood stream the more reaction Cl groups are lost so, the Pt atom bonds to a N-atom in guanosine (a part of DNA). This molecule can bond to two different guanosine units and by bridging between them it upsets the normal reproduction of DNA

103 (a)



These are coupling reactions.

103 (a)

5 In  $\text{K}_4\text{Fe}(\text{CN})_6$ , the species retains its identity in solid as well as solution state

103 (c)

6 The formula of dichlorodioxalatochromium (III) is  $[\text{Cr}(\text{Cl}_2)(\text{ox})_2]^{3-}$

Primary valency of a metal (Cr) in the complex  
 $\equiv$  oxidation number of that metal

$$= +3$$

Secondary valency of chromium in complex  
 $=$  coordination number  
 $= +6$

( $\because$  Coordination number is the number of ligands attached to the central metal ion and oxalate ion is a bidentate ligand *i.e.*, can coordinate at two positions)


103 (a)

7 Friedel-Crafts reaction involves alkylation or acylation in benzene nucleus using alkylating or acylating reagents in presence of anhy.  $\text{AlCl}_3$ .

103 (d)

8 According to IUPAC system, the IUPAC name of a compound is written as single word as far as possible

103 (b)



104 (a)

0  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$  gives four mole of ions on complete ionisation.



104 (a)

2 It is a reason for the fact.

104 (d)

3 Halogens no doubt *o*- and *p*-directing gp. but they deactivate the ring.

104 (a)

4 In metal carbonyls CO has ox. no. equal to zero.

104 (d)


5 The dihedral angle is  $60^\circ$

104 (b)

6  $pK_a$  are 10.21, 10.14 and 7.15 respectively.

104 (d)

7 (i)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$  (ii)  $\text{CH}_3\text{CH}_2 - \underset{\text{OH}}{\text{CH}} - \text{CH}_3$



104 (c)

8  $\text{Na}_2[\text{Fe}(\text{CN})_5\text{NO}]$

Sodium pentacyanonitroso ferrate (II).

104 (d)


9  $\text{Na}^+$  does not possess the tendency to form complex ion because of non-availability of *d*-orbitals.

105 (b)

0 In the complex  $\text{K}_2\text{Fe}[\text{Fe}(\text{CN})_6]$  both the iron atoms are present in same oxidation state

105 (c)


1 Carbon bonded with four different groups is known as chiral carbon atom. In case of given compound



The number of chiral carbon atoms are two

105 (a)

2  $[\text{Co}(\text{NH}_3)_3\text{Cl}_3]$  does not give a precipitate with  $\text{AgNO}_3$  solution because all the chloride ions are non-ionizable.




105 (c)


3 The IUPAC name of the compound  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$  is pentaammine chloro cobalt (III) chloride.

105 (c)


4 The electronic configuration of Ni in  $[\text{Ni}(\text{CN})_4]^{2-}$ ,  $[\text{NiCl}_4]^{2-}$  and  $\text{Ni}(\text{CO})_4$  are as following  
 Ni in  $[\text{Ni}(\text{CN})_4]^{2-}$



$Ni^{2+}$  in  $[NiCl_4]^{2-}$



$Ni$  in  $[Ni(CO)_4]$



CO and  $CN^-$  are strong ligands so, they induces pairing of electrons so, their complexes are diamagnetic which,  $Cl^-$  is a weak ligand so, it does not induce the pairing of electrons so, its complex is paramagnetic

105 (b)


- 5 Presence of  $CH_3$  gp. (an electron repelling group) decreases acidic character in benzoic acid. Presence of electron withdrawing gps. ( $NO_2$ ,  $Cl$ ,  $SO_3H$ ) increases the acidic nature.

105 (a)

- 6  $Ni(CO)_4$  is a tetrahedral complex and is diamagnetic due to the absence of unpaired electron.

105 (d)

- 7 Such a carbon atom to which four different atoms or groups are attached is called asymmetric carbon



105 (d)

- 9 Coordination sphere is interchanged.

106 (a)

- 0 Racemic tartaric acid is optically inactive due to external compensation. Racemic tartaric acid is an equimolar mixture of optically active  $d$ - and  $l$ -forms. This form of tartaric acid is optically inactive due to external compensation

106 (d)

- 1  $-NH_2$  gp. is highly susceptible to oxidant ( $HNO_3$ ) and thus, first protected by acetylation.

106 (d)

- 2 All are the required facts for bridging ligands.

106 (a)

- 3  $C_6H_5COOH$  is monobasic acid;

$\therefore$  Mol. wt. = Eq. wt.

106 (d)

- 4 Cannizzaro's reaction.

106 (b)

- 5 NO is in  $NO^+$  form.

Let the oxidation state of Fe is  $x$

$$x + 0 \times 5 + (+1) = +2$$

$$x = 2 - 1$$

$$x = +1$$

Hence, the oxidation number of Fe in brown ring  $[Fe(H_2O)_5NO]^{2+}$  is +1.

106 (a)

- 6  $Cr^{3+}$ :  $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^3$ . The  $3d_{xy}^1, 3d_{xz}^1, 3d_{yz}^1$  has lower energy.

106 (a)

- 7 Number of moles of pentaqua chloro chromium III chloride in the solution.

$$n = \frac{0.01 \times 100}{1000} = 0.001$$

1 mole of  $[Cr(H_2O)_5Cl]Cl_2$  gives 2 moles of chloride ions.


Moles of chloride ions =  $0.001 \times 2 = 0.002$

Mass of AgCl produced =  $0.002 \times 143.5$

$$= 287.0 \times 10^{-3} g$$

106 (c)

- 8 The four isomers are:



106 (c)

- 9 Both optical isomerism and geometrical isomerism are examples of stereoisomerism.

107 (d)



$$x \quad 0 \quad -2$$

$$x + 0 + (-2) = 0 \Rightarrow x = +2 \text{ is valency and 4 is}$$

C.N. of Ni

107 (d)

- 1  $C_6H_5CHO$  does not reduce Fehling's solution.

107 (d)

- 3  $n$ -butane can exist in an infinite number of conformations

107 (a)

- 4 This is correct order of acidic nature of nitrobenzoic acids. Also follow ortho effect.

107 (d)

- 7 Positive charge on  $-\overset{+}{C}H_2$

is dispersed due to electron releasing nature of methoxy group.

108 (c)

0 An organometallic compound is considered as a substance contains a carbon-metal bond *e.g.*,  $\text{Al}_2(\text{CH}_3)_6$ .

108 (c)


1 Picramide is 2, 4, 6-trinitroaniline.

108 (d)

2 Except alkynes, chain isomerism is observed when the number of carbon atoms is four or more than four

108 (d)

3 The isomerism which arises due to restricted rotation about a bond in a molecule is known as geometrical isomerism



All of these form geometrical isomers

108 (a)

4 The choice of principal functional group is made on the basis of the following order

Carboxylic acid > sulphonic acid > anhydride > esters > acid halide > acid amide > nitrile > aldehyde > ketone > alcohol > amine

108 (d)

5  $[\text{Pt}(\text{NH}_3)\text{Cl}_2\text{Br}]\text{Cl} \rightleftharpoons [\text{Pt}(\text{NH}_3)\text{Cl}_2\text{Br}]^+ + \text{Cl}^-$   
 $\text{Cl}^- + \text{Ag}^+ \rightarrow \text{AgCl}$

108 (b)

7  $-\text{COOH}$  gp. reacts with  $\text{NaHCO}_3$  to give effervescence.

108 (d)




8  $\text{C}_6\text{H}_5\text{NO}_2 \xrightarrow{\text{Fe}/\text{H}_2\text{O}_2(v)} \text{C}_6\text{H}_5\text{NO}$

108 (c)

Complex ion	Hybridisation of central atom
$[\text{Fe}(\text{CN})_6]^{4-}$	$d^2sp^3$ (inner)
$[\text{Mn}(\text{CN})_6]^{4-}$	$d^2sp^3$ (inner)
$[\text{Co}(\text{NH}_3)_6]^{3+}$	$d^2sp^3$ (inner)
$[\text{Ni}(\text{NH}_3)_6]^{2+}$	$sp^3d^2$ (outer)


109 (a)

0 Electronic configuration of  $\text{Co}^{2+}$  ion



$\text{F}^-$  is a weak ligand. It cannot pair up electrons with *d*-subshell and forms outer orbital octahedral complex.


(b)  $[\text{Co}(\text{NH}_3)_6]^{3+}$  ion



4d




(c)  $[\text{Fe}(\text{CN})_6]^{3+}$  ion



4d



(d)  $[\text{Cr}(\text{NH}_3)_6]^{3+}$  ion



4d



$\text{NH}_3$  and  $\text{CN}^-$  are strong ligands. So, they form their inner orbital complex.

109 (c)

1 The name of complex is Carbonylchlorobis*trans*phosphineiridium (I).

109 (a)

3 Coordination isomerism is caused by the interchange of ligands between complex cation and complex anion

109 (c)


4 Due to bitter almond smell.

109 (b)

5 Compounds which do not show optical activity in spite of the presence of chiral carbon atoms are called *meso*-compounds

109 (d)

6 mmol of complex =  $30 \times 0.01 = 0.3$  Also, 1 mole of complex  $[\text{Cr}(\text{H}_2\text{O})_5\text{Cl}]\text{Cl}_2$  gives only two moles of chloride ion when dissolved in solution.



Hence, 0.6 mmol of  $\text{Ag}^+$  would be required for precipitation.

$$\Rightarrow 0.60 \text{ mmol of } \text{Ag}^+ = 0.1 \text{ M} \times V \text{ (in mL)}$$

$$\Rightarrow V = 6 \text{ mL}$$

109 (c)

7 This is Kekule's view for  $\text{C}_6\text{H}_6$  structure.

109 (c)

8 C = O double bond of a carbonyl group is a stronger bond ( $> \text{C} = \text{O}$ , 364 kJ/mol) than the C = C bond strength (250 kJ/mol) of the enol. Thus,  $\text{CH}_3\text{COCH}_3$  is more stable

109 (b)

9 Effective atomic number (EAN) = Atomic no. of metal

$$\text{-Oxidation no.} + \text{Coordination no.} \times 2$$

For  $[\text{CoF}_6]^{2-}$ , the oxidation state of cobalt is 4.

$$\begin{aligned} \text{EAN} &= (27-4) + 6 \times 2 \\ &= 23 + 12 = 35 \end{aligned}$$

110 (d)

3 Metal atom or cation acts as Lewis acid or electron pair acceptor.

110 (c)


4 The nitration of  $\text{C}_6\text{H}_6$  does not occur at room temperature. The solution becomes dark red due to absorption of  $\text{NO}_2$  given out by  $\text{HNO}_3$ .

110 (c)

5  $[\text{Co}(\text{en})_2\text{NO}_2\text{Cl}]\text{Br}$  exhibits linkage isomerism because the  $\text{NO}_2$  group can exist as nitrito ( $-\text{ONO}$ ) and nitro ( $-\text{NO}_2$ ) group. The linkage isomers of  $[\text{Co}(\text{en})_2\text{NO}_2\text{Cl}]\text{Br}$  are as  $[\text{Co}(\text{en})_2\text{NO}_2\text{Cl}]\text{Br}$  and  $[\text{Co}(\text{en})_2\text{ONOCl}]\text{Br}$ .

110 (c)


6 The compound will be numbered as and can be named as unbranched assemblies containing 3 or more identical cycles



1,1',2',1''-tercyclo propane

110 (c)


7




The gauche conformation is most stable due to presence of H-bonding between H atom of OH and Cl

110 (c)


8 The electronic configuration of Ni in  $[\text{Ni}(\text{CN})_4]^{2-}$ ,  $[\text{Ni}(\text{Cl})_4]^{2-}$  and  $\text{Ni}(\text{CO})_4$  are :  $\text{Ni}^{2+}$  in  $[\text{Ni}(\text{CN})_4]^{2-}$  :



$\text{Ni}^{2+}$  in  $[\text{Ni}(\text{Cl})_4]^{2-}$  :



$\text{Ni}^{2+}$  in  $[\text{Ni}(\text{CO})_4]$  :



110 (a)

9 Follow IUPAC rules.

111 (b)

0 Cd has no unpaired electron in  $[\text{CdCl}_4]^{2-}$  ion.

111 (d)

1 It is a fact.


111 (a)

2 On ionization it gives maximum number of (four) ions.

111 (a)

3 Chlorophyll contains Mg, hence (a) is incorrect statement.

111 (b)




111 (b)


5 Linkage isomerism is caused due to presence of ambidentate ligands.

$[\text{Pd}(\text{PPh}_3)_2(\text{NCS})_2]$  and  $[\text{Pd}(\text{PPh}_3)_2(\text{SCN})_2]$  are linkage isomers due to SCN, ambidentate ligand.

111 (a)



111 (c)




111 (d)

8 Halogen attached on side chain behaves as in aliphatic molecule.


111 (a)

9 MO theory reveals bond order in  $\text{C}_6\text{H}_6$  lies in between 1 and 2.

112 (a)



0.02mol (Y)



0.02mol (Z)

On using one liter solution we will get 0.01 mole Y and 0.01 mole Z





112 (c)

2 Metal-metal (Fe-Fe) bond pairs up the unpaired electrons.

112 (b)

3 Follow IUPAC rules.

112 (c)



☒ Indicates lone-pair of  $\text{NH}_3$  donated to Cr

30.  $d^2sp^3$  hybridisation, octahedral, thus, correct.

31. There are three unpaired electrons, hence paramagnetic, thus correct.

32.  $d^2sp^3$ -inner orbital complex, thus incorrect

33. Due to ionisable  $\text{Cl}^-$  ions, white precipitate with  $\text{AgNO}_3$ , thus correct.

Therefore, (c) is wrong.

112 (c)



5  $\text{NO}_2^+$  attacks at *ortho-para* for (P) w. r. t. OH  
 $\text{NO}_2^+$  attacks at *ortho-para* for (Q) w. r. t.  $\text{CH}_3$  and  $\text{OCH}_3$  both

$\text{NO}_2^+$  attacks at *ortho-para* for (S) with respect to  $-\text{OCOC}_6\text{H}_5$

112 (c)

6 Follow molecular orbital theory for  $\text{C}_6\text{H}_6$ .

112 (d)



112 (d)

9 The number of atoms of the ligand that are directly bounded to the central metal atom or ion by coordinated bond is known as the coordination number of the metal atom or ion. Coordination number of metal = number of  $\sigma$  bonds formed by metal with ligand

113 (c)

0 Follow MO diagram for  $\text{C}_6\text{H}_6$ .


113 (b)

2  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$  has four unpaired electrons;  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$ ,  $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$  and  $[\text{Zn}(\text{H}_2\text{O})_6]^{2+}$  have 3, 1, 0 unpaired electrons, respectively.

113 (a)


3 Larger is the ligand, less stable is metal-ligand bond.

113 (c)



113 (a)

5 The name of reaction is Baeyer-Villiger oxidation.





113 (d)

6 Pyridine undergoes  $S_E$  reactions at 3-position but under vigorous conditions, nitration, sulphonation and halogenation occurs only at  $300^\circ\text{C}$ . Friedel-Crafts reaction is not observed in pyridine because electron pair on N-atom (Lewis base) form complex with  $\text{AlCl}_3$  (Lewis acid) and

a+ve charge on N-atom so produced decreases the activity of pyridine for  $S_E$  reaction.

113 (d)

7  $2Cl^-$  ions are inisable



113 (d)

8 Benzene is very good solvent.

113 (a)

9 Salol is phenyl salicylate and is used as antiseptic.


114 (a)

0  $-NO_2$  gp. is deactivating gp. for  $S_E$  reaction.

114 (d)


1 Geometrical isomerism is found in compounds having coordination no. 4 (square planar and not tetrahedral shape) as well as coordination no.6.

Coordination no. 4 (square planar)




Showing geometrical isomerism  $M_{ABCD}$ .

Coordination no. 6 (octahedral)




Showing geometrical isomerism.




114 (a)

3 The presence of  $m$ -directing groups in benzene nucleus simply decreases electron density at  $o$ - and  $p$ - whereas no change in electron density at  $m$ -position is noticed.



On the contrary  $o$ - and  $p$ -directing groups in nucleus increase the electron density at  $o$ - and  $p$ - position.



Thus, presence of  $o$ - and  $p$ -directing groups provide seats for  $S_E$  reactions or activates the ring, whereas presence of  $m$ -directing groups does not activate the ring and thus, deactivate the ring for  $S_E$  reactions.

114 (a)

4 Aniline is basic.

114 (d)

5 Ferrocene is diphenyl iron complex.

114 (c)

6 Effective atomic number (EAN)

$$= \text{Atomic no.} - O.S. + 2 \times C.N.$$

$$= 28 - 0 + 2 \times 4$$

$$= 28 + 8$$

$$= 36$$

114 (d)

8  $[Co(en)_2Cl_2]$  forms optical and geometrical isomers.

114 (c)



9 Only *cis*-octahedral compounds show optical activity.

115 (b)

0 A square planar complex results from  $dsp^2$ -hybridisation involving  $(n-1) d_{x^2-y^2}, ns, np_x$  and  $np_y$  atomic orbitals.

115 (a)

2  $HNO_3$  accepts a proton from  $H_2SO_4$ .



115 (c)

3 Lithium tetrahydroaluminate is  $Li[Al(H)_4]$