

Differential Calculus

Single Correct Answer Type

1. Let $f(x) = 4x + 8\cos x - \ln\{\cos x(1+\sin x)\} + \tan x - 2\sec x - 6$. If $f(x) > 0 \forall x \in (0, a)$ then

a) $a = \frac{\pi}{6}$ b) $a = \frac{\pi}{3}$ c) $a = \frac{\pi}{2}$ d) none of these

Ans. a

Sol.
$$\begin{aligned} f'(x) &= 4 - 8\sin x - \frac{(-\sin x + \cos^2 x - \sin^2 x)}{\cos x(1+\sin x)} + \sec^2 x - \sec x \tan x \\ &= 4(1-2\sin x) + \sec^2 x(1-2\sin x) - 4\sec(1-2\sin x) \\ &= f(x) = (\sec x - 2)^2(1-2\sin x) \end{aligned}$$

If $f(x) > 0 \forall x \in (0, a)$, then $f(x)$ is increasing in $(0, a) \Rightarrow a = \frac{\pi}{6}$

2. If $f(x)$ is continuous for all real values of x , then $\sum_{r=1}^n \int_0^1 f(r-1+x) dx =$

a) $\int_0^n f(x) dx$ b) $\int_0^1 f(x) dx$ c) $n \int_0^1 f(x) dx$ d) $(n-1) \int_0^1 f(x) dx$

Ans. a

Sol.
$$\begin{aligned} \sum_{r=1}^n \int_0^1 f(r-1+x) dx &= \int_0^1 f(x) dx + \int_0^1 f(1+x) dx + \int_0^1 f(2+x) dx + \dots + \int_0^1 f(n-1+x) dx \\ &= \int_0^1 f(x) dx + \int_1^2 f(x) dx + \int_2^3 f(x) dx + \dots + \int_{n-1}^n f(x) dx = \int_0^n f(x) dx \end{aligned}$$

3. The coordinates of the point on the curve $x^3 = y(x-a)^2$, $a > 0$ where the ordinate is minimum

a) $(2a, 8a)$ b) $\left(-2a, \frac{-8a}{9}\right)$ c) $\left(3a, \frac{27a}{4}\right)$ d) $\left(-3a, \frac{-27a}{16}\right)$

Ans. c

The ordinates of any point on the curve is given by $y = \frac{x^3}{(x-a)^2}$

Sol.
$$\frac{dy}{dx} = \frac{x^2(x-3a)}{(x-a)^3}$$

Now, $\frac{dy}{dx} = 0 \Rightarrow x = 0$ or $x = 3a$

$$\frac{d^2y}{dx^2} \Big|_{x=0} = 0 \text{ and } \frac{d^2y}{dx^2} \Big|_{x=3a} = \frac{72a^5}{(2a)^6} > 0$$

Hence y is minimum at $x = 3a$ and is equal to $\frac{27a}{4}$

4. Let $I_n = \int_0^{\frac{\pi}{2}} \frac{\sin(2n-1)x}{\sin x} dx$, $J_n = \int_0^{\frac{\pi}{2}} \frac{\sin^2 nx}{\sin^2 x} dx$, $n \in N$, then

a) $J_{(n+1)} - J_n = I_n$ b) $J_{(n+1)} - J_n = I_{(n+1)}$ c) $J_{n+1} + J_n = J_n$ d) $J_{n+1} + J_{n+1} = J_n$

Ans. b

Sol.
$$J_n - J_{n-1} = \int_0^{\frac{\pi}{2}} \frac{\sin^2 nx - \sin^2(n-1)x}{\sin^2 x} dx - \int_0^{\frac{\pi}{2}} \frac{\sin(2n-1)x - \sin x}{\sin^2 x} dx = I_n$$

i.e. $J_n - J_{n-1} = I_n \Rightarrow J_{n+1} - J_n = I_{n+1}$

5. A curve whose concavity is directly proportional to the logarithm of its x-coordinates at any of the curve, is given by

a) $c_1 x^2 (2 \log x - 3) + c_2 x + c_3$ b) $c_1 x^2 (2 \log x + 3) + c_2 x + c_3$
 c) $c_1 x^2 (2 \log x) + c_2$ d) none of these

Ans. a

Sol.
$$\frac{d^2y}{dx^2} = k \log x \Rightarrow \frac{dy}{dx} = k(x \log x - x) + A$$

$$\Rightarrow y = k \left[\frac{1}{2} x^2 \log x - \int \frac{1}{2} x^2 \cdot \frac{1}{x} - \frac{x^2}{2} dx \right] + Ax + B$$

$$\Rightarrow y = \frac{k}{4} \{2x^2 \log x - x^2 - 2x^2\} + Ax + B$$

$$\Rightarrow y = c_1 (2 \log x - 3)x^2 + c_2 x + c_3$$

6. The domain of the function $f(x) = \sqrt{3 - 2^x - 2^{1-x}} + \sqrt{\sin^{-1} x}$ is

a) $[-1, 0]$ b) $[0, 1]$ c) $\left[\frac{1}{2}, 1\right]$ d) $[1, 2]$

Ans. b

Sol. $\sin^{-1} x \geq 0 \Rightarrow 0 \leq x \leq 1$
 and $2^x + 2^{1-x} \leq 3 \Rightarrow 2^x + 2 \cdot 2^{-x} - 3 \leq 0$
 Put $2^x = t$, then $t^2 - 3t + 2 \leq 0 \Rightarrow (t-2)(t-1) \leq 0$
 $\Rightarrow 1 \leq t \leq 2$ i.e. $1 \leq 2^x \leq 2$
 $\Rightarrow 0 \leq x \leq 1$

7. Area bounded by the curve $y = \sin x$, $y = \cos x$, $x = -\frac{\pi}{3}$, $x = 2\pi$

a) $4\sqrt{2} - \left(\frac{\sqrt{3}+1}{2}\right)$ b) $\sqrt{2} + \left(\frac{\sqrt{3}+1}{2}\right)$ c) $\sqrt{2} - \left(\frac{\sqrt{3}+2}{2}\right)$ d) $4\sqrt{2} + \left(\frac{\sqrt{3}+1}{2}\right)$

Ans. d

Sol. $A = \int_{-\pi/3}^{2\pi} |\sin x - \cos x| dx \Rightarrow 4\sqrt{2} + \frac{\sqrt{3}+1}{2}$

8. Let $f(1) = 1$ and $f(n) = 2 \sum_{r=1}^{n-1} f(r)$, then $\sum_{n=1}^m f(n)$ is equal to

a) $3^{m-1} - 1$ b) 3^{m-1} c) $3^m - 1$ d) none of these

Ans. b

Sol. $f(n) = 2(f(1) + f(2) + \dots + f(n-1))$

$$\therefore f(n+1) = 2(f(1) + f(2) + \dots + f(n))$$

$$\Rightarrow f(n+1) = 3f(n) \text{ for } n \geq 2$$

$$\text{Also } f(2) = 2f(1) = 2$$

$$f(3) = 3f(2) = 2 \cdot 3$$

$$\sum_{n=1}^m f(n) = f(1) + f(2) + \dots + f(m)$$

$$= 1 + 2 + 2 \cdot 3 + 2 \cdot 3^2 + \dots + 2 \cdot 3^{m-2} = 1 + 2(1 + 3 + 3^2 + \dots + 3^{m-2})$$

9. $I = \int \frac{2+3\cos\theta}{\sin\theta+2\cos\theta+3} d\theta$, then

a) $I = \frac{6\theta}{5} + \frac{3}{5} \log|\sin\theta+2\cos\theta+3| - \frac{8}{5} \tan^{-1}\left(\frac{\tan\left(\frac{\theta}{2}\right)+1}{2}\right) + c$

b) $I = \frac{6\theta}{5} - \frac{3}{5} \log|\sin\theta+2\cos\theta+3| - \frac{8}{5} \tan^{-1}\left(\frac{\tan\left(\frac{\theta}{2}\right)+1}{2}\right) + c$

c) $I = \frac{6\theta}{5} - \frac{3}{5} \log|\sin\theta+2\cos\theta+3|$

d) none of these

Ans. a

Sol. $2+3\cos\theta = I(\sin\theta+2\cos\theta+3) + m(\cos\theta-2\sin\theta) + n$, then integrate

10. The value of $\lim_{n \rightarrow \infty} \left(\frac{\sqrt{n}}{(3+4\sqrt{n})^2} + \frac{\sqrt{n}}{\sqrt{2}(3\sqrt{2}+4\sqrt{n})^2} + \dots + \frac{1}{49n} \right)$ is equal to

a) $\frac{1}{14}$

b) $\frac{2}{7}$

c) $\frac{3}{7}$

d) none of these

Ans. a

Sol. $\lim_{n \rightarrow \infty} \sum_{r=1}^n \frac{\sqrt{n}}{\sqrt{r}(3\sqrt{r}+4\sqrt{n})^2}$

$$\text{Put } \frac{r}{n} = x \Rightarrow \frac{1}{n} = dx$$

$$= \int_0^1 \frac{dx}{\sqrt{x}(3\sqrt{x}+4)^2} = \frac{1}{14}$$

11. Solution of differential equation $xdy - (y + xy^3(1 + \log x))dx = 0$

a) $\frac{-x^2}{y^2} = \frac{2x}{3} \left(\frac{2}{3} + \log x \right) + c$

b) $\frac{x^2}{y^2} = \frac{2x^2}{3} \left(\frac{2}{3} + \log x \right) + c$

c) $\frac{-x^2}{y^2} = \frac{2x^3}{3} \left(\frac{2}{3} + \log x \right) + c$ d) none of these

Ans. c

Sol. $-d\left(\frac{x}{y}\right) = xy(1 + \log x)dx$

$\int -\frac{x}{y} d\left(\frac{x}{y}\right) = \int x^2(1 + \log x)dx$ gives solution

12. Let $f: R \rightarrow R$ and $g: R \rightarrow R$ be twice differentiable function satisfying $f''(x) = g''(x)$, $2f'(1) = g'(1) = 4$ and $3f(2) = g(2) = 9$. The value of $f(4) - g(4)$ is equal to

- a) -6 b) -16 c) -10 d) -8

Ans. c

Sol. $f'(x) = g(x) - 2$

$f(x) = g(x) - 2x - 2$

$f(u) - g(u) = -10$

13. Let a, b, c be three real numbers such that $a < b < c$. Let $f(x)$ be continuous $\forall x \in [a, c]$ and differentiable $\forall x \in (a, c)$. If $f''(x) > 0 \forall x \in (a, c)$ then

- a) $(c-b)f(a) + (b-a)f(c) > (c-a)f(b)$ b) $(c-b)f(a) + (a-c)f(b) < (a-b)f(c)$
 c) $f(a) < f(b) < f(c)$ d) none of these

Ans. a

Sol. By LMVT

$$\frac{f(b)-f(a)}{b-a} > \frac{f(c)-f(b)}{c-b}$$

14. The solution of $y^5 x + y - x \frac{dy}{dx} = 0$ is

- a) $\frac{x^4}{4} + \frac{1}{5} \left(\frac{x}{y} \right)^5 = c$ b) $\frac{x^5}{5} + \frac{1}{4} \left(\frac{x}{y} \right)^4 = c$ c) $\left(\frac{x}{y} \right)^5 + \frac{x^4}{4} = c$ d) $(xy)^4 + \frac{x^5}{5} = c$

Ans. b

Sol. $y^5 x dx + y dy - x dy = 0$, multiply by x^3 / y^5

$$\Rightarrow x^4 dx + \frac{x^3}{y^3} (d(x/y)) = 0$$

$$\Rightarrow \frac{x^5}{5} + \frac{1}{4} \left(\frac{x}{y} \right)^4 = c$$

15. A point P lying inside the curve $y = \sqrt{2ax - x^2}$ is moving such that its shortest distance from the curve at any position is greater than its distance from x-axis. The point P encloses a region whose area is equal to

- a) $\frac{\pi a^2}{2}$ b) $\frac{a^2}{3}$ c) $\frac{2a^2}{3}$ d) $\left(\frac{3\pi - 4}{6} \right) a^2$

Ans. c

Sol.

$$y = \sqrt{2ax - x^2} \Rightarrow (x-a)^2 + y^2 = a^2$$


Let P(h, k) be a point then BP > PN

For the boundary condition BP = PN = k

$$\text{Now } AP = a - h = \sqrt{(h-a)^2 + k^2} \Rightarrow k = h - \frac{h^2}{2a}$$

$$\therefore \text{boundary of the region is } y = x - \frac{x^2}{2a}$$

$$\text{Required area} = 2 \int_0^a \left(x - \frac{x^2}{2a} \right) dx = \frac{2a^2}{3}$$



16. If $\log_x (\log_y k) > 0$ where $x, k \in (0, 1)$ then $y \in$
 a) (0, x) b) (0, k) c) (k, 1) d) \mathbb{R}^+

Ans. c

Sol. $\log_y k < 1$

case 1 : if $y > 1 \Rightarrow k < y$

for $\log_y k > 0 \Rightarrow k > 1$ which is not possible

case 2 : if $y < 1 \Rightarrow k > y$

and for $\log_y k > 0 \Rightarrow k < 1$ which is true

17. Period of $f(x) = x - [x + \lambda] - \mu$ where $\lambda, \mu \in \mathbb{R}$ and $[\cdot]$ denotes the g.i.f is
 a) λ b) μ c) $|\lambda - \mu|$ d) 1

Ans. d

$$\begin{aligned} f(x) &= x - [x + \lambda] - \mu = x + \lambda - [x + \lambda] - (\lambda + \mu) \\ &= \{x + \lambda\} - (\lambda + \mu) \\ \therefore \text{Period of } f(x) &= 1 \end{aligned}$$

18. If $f(x) = 2\sin^3 x - 3\sin^2 x + 12\sin x + 5 \forall x \in \left(0, \frac{\pi}{2}\right)$, then

- a) f is increasing in $\left(0, \frac{\pi}{2}\right)$ b) f is decreasing in $\left(0, \frac{\pi}{2}\right)$
 c) f is increasing $\left(0, \frac{\pi}{4}\right)$ and decreasing in $\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$
 d) f is decreasing in $\left(0, \frac{\pi}{4}\right)$ and increasing in $\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$

Ans. a

$$\text{Sol. } f'(x) = 6\cos x (\sin^2 x - \sin x + 2) > 0 \forall x \in \left(0, \frac{\pi}{2}\right)$$

Thus f(x) is increasing in $\left(0, \frac{\pi}{2}\right)$

19. Total number of points of non-differentiability of $f(x) = [3 + 4\sin x]$ in $[\pi, 2\pi]$ where $[\cdot]$ denote the g.i.f are

- a) 5 b) 6 c) 8 d) 9


Ans. c

Sol.

$$f(x) = 3 + [4 \sin x]$$

$f(x)$ is non-differentiable where $g(x) = [4 \sin x]$ is non differentiable

In $[\pi, 2\pi]$, $g(x)$ is clearly non-differentiable at 8 points.



20. If $f(x) + 2f(1-x) = x^2 + 1 \forall x \in R$ and $\int_0^k f(x) dx = 0$, then k equals to
 a) 3 b) 2 c) 4 d) none of these

Ans. a

Sol. Putting $(1-x)$ for x and subtracting we get $f(x) = \frac{x^2 - 4x + 3}{3}$

$$\text{Now } \int_0^k \frac{x^2 - 4x + 3}{3} dx = 0 \Rightarrow \frac{k^3}{3} - 2k^2 + 3k = 0$$

$$\Rightarrow k = 3$$

21. A point $P(x, y)$ moves in such a way that $[x+y+1] = [x]$ (where $[]$ denotes g.i.f) and $x \in (0, 2)$. Then the area representing all the possible positions of P equals

- a) $\sqrt{2}$ sq. units b) $2\sqrt{2}$ sq. units c) $4\sqrt{2}$ sq. units d) none of these

Ans. d

Sol.

If $x \in (0, 1)$


Then $-1 \leq x+y < 0$

and if $x \in (1, 2)$

$0 \leq x+y < 1$

Required area =

$$4 \left(\frac{1}{2} \cdot 1 \cdot \sqrt{2} \sin \frac{\pi}{4} \right) = 2 \text{ sq units}$$



22. Let $f(x)$ be a polynomial with real coefficients satisfies $f(x) = f'(x) \times f'''(x)$. If $f(x) = 0$ satisfies $x = 1, 2, 3$ only then the value of $f'(1) \times f'(2) \times f'(3) =$
 a) positive b) negative c) 0 d) inadequate data

Ans. c

Sol. $f(x) = f'(x) \times f'''(x)$ is satisfied by only the polynomial of degree 4.

Since $f(x) = 0$ satisfies $x = 1, 2, 3$ only. It is clear one of the root is twice repeated.

$$\Rightarrow f'(1) f'(2) f'(3) = 0$$

23. The value of $\lim_{n \rightarrow \infty} \left(\frac{n!}{(mn)^n} \right)^{1/n}$ is

a) em

b) $\frac{e}{m}$ c) $\frac{1}{em}$

d) none of these

Ans. c

$$\text{Sol. } L = \lim_{n \rightarrow \infty} \frac{1}{m} \left(\frac{1}{n} \cdot \frac{2}{n} \cdot \frac{3}{n} \cdots \frac{n}{n} \right)^{1/n}$$

$$\ln L = \lim_{n \rightarrow \infty} \left[\ln \left(\frac{1}{m} \right) + \frac{1}{n} \left(\ln \frac{1}{n} + \ln \frac{2}{n} + \dots + \ln \frac{n}{n} \right) \right]$$

$$= \ln m + \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{r=1}^n \ln \left(\frac{r}{n} \right) = -\ln m + \int_0^1 \ln x dx = -\ln m - 1 = \ln \left(\frac{1}{em} \right)$$

$$\therefore L = \frac{1}{em}$$

24. Let $A = \{1, 2, 3, 4, 5\}$ and $f : A \rightarrow A$ be an into function such that $f(i) \neq i \forall i \in A$, then number of such functions f are

a) 1024

b) 904

c) 984

d) none of these

Ans. d

Sol. Total number of functions for which $f(i) \neq i = 4^5$

and number of onto functions in which $f(i) \neq i = 44$

\Rightarrow required numbers of functions = 980

25. The area of the region bounded between the curves $y = e|x|\ln|x|$, $x^2 + y^2 - 2(|x| + |y|) + 1 \geq 0$ and x-axis where $|x| \leq 1$, if α is the x-coordinate of the point of intersection of curves in 1st quadrant, is

$$\text{a) } 4 \left[\int_0^\alpha ex \ln x dx + \int_\alpha^1 \left(1 - \sqrt{1 - (x-1)^2} \right) dx \right]$$

$$\text{b) } \left[\int_0^\alpha ex \ln x dx - \int_1^\alpha \left(1 - \sqrt{1 - (x-1)^2} \right) dx \right]$$

$$\text{c) } 2 \left[- \int_0^\alpha ex \ln x dx + \int_\alpha^1 \left(1 - \sqrt{1 - (x-1)^2} \right) dx \right]$$


$$\text{d) } 2 \left[\int_0^\alpha ex \ln x dx + \int_\alpha^1 \left(1 - \sqrt{1 - (x-1)^2} \right) dx \right]$$

Ans. c

Sol.

Required area is

$$2 \left[\int_0^\alpha ex \ln x dx + \int_1^\alpha \left(1 - \sqrt{1 - (x-1)^2} \right) dx \right]$$



26. The value of $\lim_{n \rightarrow \infty} n \left[\frac{1}{3n^2 + 8n + 4} + \frac{1}{3n^2 + 16n + 16} + \dots n \text{ terms} \right]$ is

$$\text{a) } \frac{1}{4} \ln \left(\frac{9}{5} \right)$$

$$\text{b) } \frac{1}{5} \ln \left(\frac{9}{5} \right)$$

$$\text{c) } \frac{1}{4} \ln \left(\frac{8}{5} \right)$$

$$\text{d) } \frac{1}{4} \ln \left(\frac{9}{7} \right)$$

Ans. a


Sol. Use definite integral of first principal as a limit of sum

$$\lim_{n \rightarrow \infty} \sum_{r=1}^n \frac{1}{4\left(1 + \frac{r}{n}\right)^2 - 1} \cdot \frac{1}{n}$$

27. The area of the region containing the points satisfying $|y| + \frac{1}{2} \leq e^{-|x|}$, $\max(|x|, |y|) \leq 2$ is

a) $2\log\left(\frac{e}{2}\right)$ b) $2\log\left(\frac{2e}{3}\right)$ c) $3\log\left(\frac{e}{2}\right)$ d) $3\log\left(\frac{2e}{3}\right)$

Ans. a



28. If $y = 2^{\frac{1}{2^{1-x}}}$; then $\lim_{x \rightarrow 1^+} y$ is

a) -1 b) 1 c) 0 d) $\frac{1}{2}$

Ans. b

Sol. $\lim_{h \rightarrow 0} 2^{\frac{1}{2^{1-(1+h)}}} = 2^{-0} = 1$

29. If $y = \frac{2x+5}{3x+10}$, then $2\left(\frac{dy}{dx}\right)\left(\frac{d^3y}{dx^3}\right)$ is equal to

a) $\left(\frac{d^2y}{dx^2}\right)^2$ b) $3\frac{d^2y}{dx^2}$ c) $3\left(\frac{d^2y}{dx^2}\right)^2$ d) $3\frac{d^2x}{dy^2}$

Ans. c

Sol. $3xy + 10y = 2x + 5$, now differentiate 3 times.

30. If the number of solutions of $\ln|\sin x| = -x^2 + 2x$ when $x \in (0, \pi)$ is m and when

$x \in \left[-\frac{\pi}{2}, \frac{3\pi}{2}\right]$ is n, then $(m+n)$ is equal to

a) 2 b) 4 c) 6 d) 1

Ans. a

Sol. $m = 0, n = 2$

31. If x, {x} and 2[x] represent the segments of a focal chord and length of latus rectum of an ellipse respectively, then length of major axis of ellipse is always greater than (where $x \in \mathbb{Z}$)

a) 7 b) 5 c) 8 d) 2

Ans. d

Sol. Clearly, x , $[x]$ and $\{x\}$ are in H.P $\Rightarrow [x] = \frac{2x\{x\}}{x+\{x\}} \Rightarrow [x] = 1$

$$\Rightarrow \frac{b^2}{a} = 1 \Rightarrow a(1-e^2) = 1 \Rightarrow 2a > 2 \quad [\text{since } 0 < e < 1]$$

32. The value of $\int_3^6 (\sqrt{x+\sqrt{12x-36}} + \sqrt{x-\sqrt{12x-36}}) dx$ is equal to
 a) $6\sqrt{3}$ b) $4\sqrt{3}$ c) $12\sqrt{3}$ d) none of these

Ans. a

Sol. $I = \int_3^6 ((\sqrt{x-3} + \sqrt{3}) + (\sqrt{3} - \sqrt{x-3})) dx = 6\sqrt{3}$

33. If integral $\int \frac{dx}{(\sec x + \csc x + \tan x + \cot x)^2} = \frac{x}{a} + \frac{\sqrt{2} \cos x + \frac{p \sin x}{4}}{b} + \frac{\cos 2x}{c} + d$, then
 a + b + c is equal to
 a) -2 b) -4 c) 2 d) none of these

Ans. b

Sol. Clearly,

$$I = \int \frac{\sin^2 x \cos^2 x}{(\sin x + \cos x + 1)^2} dx = \frac{1}{4} \int \frac{((\sin x + \cos x)^2 - 1)^2}{(\sin x + \cos x + 1)} dx = \frac{1}{4} \int (\sin x + \cos x - 1)^2 dx$$

On simplifying a + b + c = -4

34. If $I_n = \int_{-n}^n (\{x+1\}\{x^2+2\} + \{x^2+3\}\{x^2+4\}) dx$, (where $\{.\}$ denotes the fractional part)
 then I_1 is equal to
 a) $-\frac{1}{3}$ b) $-\frac{2}{3}$ c) $\frac{1}{3}$ d) none of these

Ans. b

Sol. $I_1 = \int_{-1}^1 (\{x\} + \{x^3\}) \{x^2\} dx = -2 \int_0^1 \{x^2\} dx = -2 \times \frac{x^3}{3} \Big|_0^1 = -\frac{2}{3}$

35. Area bounded by $y = f^{-1}(x)$ and tangent and normal drawn to it at the points with abscissae π and 2π , where $f(x) = \sin x - x$ is

$$\text{a) } \frac{p^2}{2} - 1 \quad \text{b) } \frac{p^2}{2} - 2 \quad \text{c) } \frac{p^2}{2} - 4 \quad \text{d) } \frac{p^2}{2}$$

Ans. b

Sol. Required area $A = \int_{\pi}^{2\pi} ((\sin x - x) + 2\pi) dx = \frac{\pi^2}{2} - 2 \text{ sq.units}$

36. Let a curve $y = f(x)$, $f(x) \neq 0$ for $x \in R$ has property that for every point P on the curve length of subnormal is equal to abscissa of P. If $f(1) = 3$, then $f(4)$ is equal to
 a) $-2\sqrt{6}$ b) $2\sqrt{6}$ c) $3\sqrt{5}$ d) none of these

Ans. b

Sol. Given $y \frac{dy}{dx} = x$

$$y dy = x dx$$

$$y^2 = x^2 + c$$

$$f(1) = 3 \Rightarrow 9 - 1 + c = 8 \Rightarrow c = 8$$

$$\Rightarrow y^2 = x^2 + 8$$

$$f(x) = \sqrt{x^2 + 8}$$

$$f(4) = \sqrt{16 + 8} = 2\sqrt{6}$$

37. Range of $f(x) = \cos^{-1}\left(\frac{x^2 + x + 1}{x^4 + 1}\right)$ is

a) $\left[0, \frac{\pi}{2}\right]$

b) $\left[0, \frac{\pi}{2}\right)$

c) $\left(0, \frac{\pi}{2}\right]$

d) $[0, \pi]$

Ans. b

Sol. Let $g(x) = \frac{x^2 + x + 1}{x^4 + 1}$

$$\Rightarrow 0 < g(x) \leq 1$$

So range of $f(x)$ is $\left[0, \frac{\pi}{2}\right)$

38. If $f(x) = 0$ is a cubic equation with positive and distinct roots α, β, γ such that β is H.M of the roots of $f'(x) = 0$, then α, β and γ are in

a) A.P

b) G.P

c) H.P

d) none of these

Ans. b

Sol. $f(x) = (x - \alpha)(x - \beta)(x - \gamma)$

$$\Rightarrow f'(x) = 3x^2 - 2x(\alpha + \beta + \gamma) + \alpha\beta + \beta\gamma + \gamma\alpha$$

$$\Rightarrow \beta = \frac{2\alpha_1\beta_1}{\alpha_1 + \beta_1} \text{ (where } \alpha_1, \beta_1 \text{ are the roots of } f'(x) = 0)$$

$$\Rightarrow \beta^2 = \gamma\alpha$$

39. Let a curve $y = f(x)$, $f(x) \geq 0 \forall x \in R$ has property that for every point P on the curve, the length of subnormal is equal to abscissa of P. If $f(1) = 3$, then $f(4)$ is equal to

a) $-2\sqrt{6}$

b) $2\sqrt{6}$

c) $3\sqrt{5}$

d) none of these

Ans. b

Sol. $y \frac{dy}{dx} = x \Rightarrow y^2 = x^2 + c$

$$f(x) = \sqrt{x^2 + 8} \Rightarrow f(4) = 2\sqrt{6}$$

40. If $\int_0^{\pi/2} \frac{dx}{a^2 \cos^2 x + b^2 \sin^2 x} = \frac{\pi}{2ab}$, then the value of $\int_0^{\pi/2} \frac{dx}{(4 \cos^2 x + 9 \sin^2 x)^2}$ is equal to

a) $\frac{11\pi}{864}$

b) $\frac{13\pi}{864}$

c) $\frac{17\pi}{864}$

d) none of these

Ans. b

Sol. $I = \int_0^{\pi/2} \frac{dx}{a^2 \cos^2 x + b^2 \sin^2 x} = \frac{\pi}{2ab}$

$$\frac{dI}{da} = \frac{-\pi}{2a^2 b}$$

$$\Rightarrow \int_0^{\pi/2} \frac{\cos^2 x dx}{(a^2 \cos^2 x + b^2 \sin^2 x)} = \frac{\pi}{4a^3 b}$$

differentiating with respect to b

$$\int_0^{\pi/2} \frac{\sin^2 x dx}{(a^2 \cos^2 x + b^2 \sin^2 x)} = \frac{\pi}{4a^3 b}$$

$$\Rightarrow \int_0^{\pi/2} \frac{dx}{(a^2 \cos^2 x + b^2 \sin^2 x)} = \frac{\pi}{2ab} \left[\frac{1}{a^2} + \frac{1}{b^2} \right] = \frac{\pi}{24} \left[\frac{1}{4} + \frac{1}{9} \right] = \frac{13\pi}{864}$$

41. If $\int \frac{dx}{\cos^3 x - \sin^3 x} = A \tan^{-1}(\sin x + \cos x) + B \ln f(x) + C$, then A is equal to

a) $\frac{2}{3}$ b) $\frac{2}{5}$ c) $-\frac{2}{3}$ d) none of these

Ans. a

Sol. $I = \int \frac{dx}{(\cos x - \sin x) \left(1 + \frac{\sin 2x}{2} \right)} = \int \frac{\cos x - \sin x}{(\cos x - \sin x)^2 \left(1 + \frac{\sin 2x}{2} \right)} dx$

Put $\cos x + \sin x = t$

$$I = \frac{2}{3} \tan^{-1}(\sin x + \cos x) - \frac{2}{3\sqrt{2}} \ln f(x) + C$$

42. Solution of the differential equation $y(2x^4 + y) \frac{dy}{dx} = (1 - 4xy^2)x^2$ is given by

a) $3(x^2y)^2 + y^3 - x^3 = c$ b) $xy^2 + \frac{y^3}{3} - \frac{x^3}{3} + c = 0$
 c) $\frac{2}{5}yx^5 + \frac{y^3}{3} = \frac{x^3}{3} - \frac{4xy^3}{3} + c$ d) none of these

Ans. a

Sol. Given equation can be written as

$$2x^2y(x^2 dy + 2xy dx) + y^2 dy - x^2 dx = 0$$

$$\text{or } 2x^2y d(x^2y) + y^2 dy - x^2 dx = 0$$

Integrating, we get

$$3(x^2y)^2 + y^3 - x^3 = c$$

43. If $I = \int_0^{\pi} \frac{\cos x}{(x+2)^2} dx$, then $\int_0^{\pi} \frac{\sin x}{x+1} dx$ is equal to

a) 21 b) $\frac{1}{\pi+2} - \frac{1}{2} - 1$ c) 0 d) $\frac{1}{\pi+2} + \frac{1}{2} - 1$

Ans. d


Sol. $I = \int_0^{\pi} \cos x d \left(-\frac{1}{x+2} \right) = \left[\frac{-\cos x}{x+2} \right]_0^{\pi} - \int_0^{\pi} \frac{\sin x}{x+2} dx$

$$= \frac{1}{\pi+2} + \frac{1}{2} - \int_0^{\pi/2} \frac{\sin 2x}{x+1} dx$$

44. The number of solutions of $\sin \pi x = |\log|x||$ is

a) infinite b) 8 c) 6 d) 0

Ans. c



45. If $f(x) = |x^2 + (k-1)|x| - k|$ is non differentiable at five real points, then k will lie in

a) $(-\infty, 0)$ b) $(0, \infty)$ c) $(-\infty, 0) - \{-1\}$ d) $(0, \infty) - \{1\}$

Ans. c

Sol. $f(x) = |x^2 + (k-1)|x| - k| = |(|x|-1)(|x|+k)|$

Both roots of $(x-1)(x+k) = 0$ should be positive and distinct

$$\Rightarrow k \in (-\infty, 0) - \{-1\}$$

46. Let $g(x) = \int_a^x f(t) dt$ and $f(x)$ satisfies the following condition

$f(x+y) = f(x) + f(y) + 2xy - 1, \forall x, y \in R$ and $f'(0) = \sqrt{3+a-a^2}$, then the exhaustive set of values of x where $g(x)$ increases is

a) $(-\infty, -\frac{3}{2})$ b) $(-\frac{3}{2}, 0)$ c) $(0, \infty)$ d) $(-\infty, \infty)$

Ans. d

Sol. $f(x) = x^2 + (\sqrt{3+a-a^2})x + 1$

$$g'(x) = f(x) > 0, \forall x \in R$$

47. Number of positive continuous function $f(x)$ defined in $[0,1]$ for which

$$\int_0^1 f(x) dx = 1, \int_0^1 xf(x) dx = 2, \int_0^1 x^2 f(x) dx = 4, \text{ is}$$

a) 1 b) 4 c) infinite d) none of these

Ans. d

Sol. Multiplying these three integral by 4, -4, 1 and adding we get $\int_0^1 f(x)(x-2)^2 dx = 0$.

Hence there does not exist any function satisfying these conditions.

48. Tangents are drawn at the point of intersection P of ellipse $x^2 + 2y^2 = 50$ and hyperbola $\frac{x^2}{16} - \frac{y^2}{9} = 1$, in the first quadrant. The area of the circle passing through the point P which cuts the intercept of 2 unit length each from these tangents, is

a) 2π b) $\sqrt{2}\pi$ c) 4π d) 6π

Ans. a

Sol. Given conic are confocal so they cut orthogonally.

49. Let $f(x) = x^3 + \frac{1}{x^3}$, $x \neq 0$. If the intervals in which $f(x)$ increases are $(-\infty, a]$ and $[b, \infty)$ then $\min(b-a)$ is equal to

a) 0 b) 2 c) 3 d) 4

Ans. b

Sol. Here $f'(x) = 3x^2 - \frac{3}{x^4} \geq 0 \Rightarrow x^6 - 1 \geq 0 \Rightarrow x \in (-\infty, 1] \cup [1, \infty)$
 $\therefore \min(b-a) = \min(b) - \max(a) = 1 - (-1) = 2$

50. Let $y = f(x)$, $f : R \rightarrow R$ be an odd differentiable function such that $f'''(x) > 0$ and $g(\alpha, \beta) = \sin^8\alpha + \cos^8\beta + 2 - 4 \sin^2\alpha \cos^2\beta$. If $f'''(g(a, b)) = 0$, then $\sin^2 a + \sin^2 b$ is equal to

a) 0 b) 1 c) 2 d) 3

Ans. b

Sol. $f''(x)$ is odd function $\Rightarrow g(\alpha, \beta) = 0$
 $\Rightarrow (\sin^4 \alpha - 1)^2 + (\cos^4 \beta - 1)^2 + 2(\sin^2 \alpha - \cos^2 \beta)^2 = 0$
 $\Rightarrow \sin^2 \alpha + \sin^2 \beta = 1$

51. If $f(x) = \int_0^x (f(t))^2 dt$, $f : R \rightarrow R^+$ be differentiable function and $f(g(x))$ is differentiable function at $x = a$, then

a) $g(x)$ must be differentiable at $x = a$ b) $g(x)$ may be non-differentiable at $x = a$
c) $g(x)$ may be discontinuous at $x = a$ d) none of these

Ans. a

Sol. Here, $f'(x) = (f(x))^2 > 0$; $\frac{d}{dx} f(g(x))|_{x=a} = f'(g(x)) \lim_{x \rightarrow a} \frac{g(x) - g(a)}{x - a}$
As $f'(g(x)) \neq 0$

$g(x)$ must be differentiable at $x = a$.

52. A polynomial of 6th degree $f(x)$ satisfies $f(x) = f(2-x)$ " $x \in R$, if $f(x) = 0$ has 4 distinct and two equal roots, then sum of roots of $f(x) = 0$ is

a) 4 b) 5 c) 6 d) 7

Ans. c

Sol. Let α be the root of $f(x) = 0 \Rightarrow f(\alpha) = f(2-\alpha) = 0$

$f(x)$ has 4 distinct and two equal roots.

\therefore sum of roots = 6

53. The number of integral solutions of equation $\int_0^x \frac{\ln t dt}{x^2 + t^2} - p \ln 2 = 0 ; x > 0$ is
- a) 0 b) 1 c) 2 d) 3

Ans. c

Sol. We have on putting $t = \frac{x^2}{2}$ and solving

$$\int_0^\infty \frac{\ln x}{x^2 + t^2} dt = \frac{2\pi \ln x}{x} \Rightarrow \frac{\ln x}{x} = \frac{\ln 2}{2}$$

$\Rightarrow x = 2$ and 4; two solutions.

54. If $f(x) = \begin{cases} e^{x-1}, & 0 \leq x \leq 1 \\ x+1 - \{x\}, & 1 < x < 3 \end{cases}$ and $g(x) = x^2 - ax + b$, such that $f(x), g(x)$ is continuous in $[0, 3]$ then the values of a and b is

- a) 2, 3 b) 3, 2 c) $\frac{3}{2}, 1$ d) none of these

Ans. b

Sol. Clearly $f(x)$ is discontinuous at $x = 1$ and 2, for $f(x), g(x)$ to be continuous at $x = 1$ and 2; $g(1)$ and $g(2) = 0 \Rightarrow a = 3$ and $b = 2$

55. $\int_0^{16n^2/p} \cos \frac{p}{2} \sin \frac{u}{n} du$ is
- a) 0 b) 1 c) 2 d) 3

Ans. a

Sol. Let $\frac{x\pi}{n} = t \Rightarrow \int_0^{16n^2} \cos \frac{\pi}{2} \left[\frac{\pi x}{n} \right] dx = \frac{n}{\pi} \int_0^{16n} \cos \frac{\pi}{2} [t] dt = \frac{4n^2}{\pi} \int_0^4 \cos \frac{\pi}{2} [t] dt = 0$

56. If $\int_{-2}^1 (ax^2 - 5) dx = 0$ and $\int_1^2 (bx + c) dx = 0$ then
- a) $ax^2 - bx + c = 0$ has atleast one root in $(1, 2)$
 b) $ax^2 - bx + c = 0$ has atleast one root in $(-2, -1)$
 c) $ax^2 + bx + c = 0$ has atleast one root in $(-2, -1)$
 d) none of these

Ans. b

Sol. We have $\int_{-2}^{-1} (ax^2 - 5) dx + \int_1^2 (bx + c) dx + 5 = \int_{-2}^{-1} (ax^2 - 5 - bx + c + 5) dx = 0$
 $\Rightarrow ax^2 - bx + c = 0$ has atleast one root in $(-2, -1)$