

- c) $2\text{Cr}^{3+} + 7\text{H}_2\text{O} \rightarrow \text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^-$ d) $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + \text{e}^-$
16. The anion nitrate is converted into ammonium ion. The equivalent mass of nitrate ion in the reaction would be:
 a) 6.20 b) 7.75 c) 10.5 d) 21.0
17. Which acts as a reducing agent?
 a) HNO_3 b) KMnO_4 c) H_2SO_4 d) $(\text{COOH})_2$
18. What weight of HNO_3 is needed to convert 5 g I_2 into HIO_3 , $\text{HNO}_3 \rightarrow \text{NO}$?
 a) 4.13 g b) 24.8 g c) 6.2 g d) 10.2 g
19. When SO_2 is passed in acidified potassium dichromate solution, the oxidation state of S is changed from :
 a) + 4 to 0 b) +4 to +2 c) +4 to +6 d) +6 to +4
20. Among the properties given below, the set of properties shown by CN^- ion towards metal species is :
 1. Reducing; 2. Oxidising; 3. Complexation
 a) 1, 3 b) 1, 2, 3 c) 1, 2 d) 2, 3
21. Solution of sodium metal in liquid NH_3 is strongly reducing due to the presence of :
 a) Sodium atoms b) Solvated electrons c) NaOH d) Sodium amide
22. Oxidation numbers of Fe in Fe_3O_4 are :
 a) +2 and +3 b) +1 and +2 c) +1 and +3 d) None of these
23. It is found that V forms a double salt isomorphous with Mohr's salt. The oxidation number of V in this compound is :
 a) + 3 b) + 2 c) + 4 d) -4
24. MnO_4^- is a good oxidising agent in different medium changing to
 $\text{MnO}_4^- \rightarrow \text{Mn}^{2+}$
 $\rightarrow \text{MnO}_4^{2-}$
 $\rightarrow \text{MnO}_2$
 $\rightarrow \text{Mn}_2\text{O}_3$
 Changes in oxidation number respectively are
 a) 1,3,4,5 b) 5,4,3,2 c) 5,1,3,4 d) 2,6,4,3
25. The oxidation number of Ba in barium peroxide is :
 a) +2 b) -1 c) +4 d) +6
26. Strongest reducing agent among the following is :
 a) K b) Mg c) Al d) Ba
27. The eq. wt. of $\text{Na}_2\text{S}_2\text{O}_3$ as reductant, in the reaction, $\text{Na}_2\text{S}_2\text{O}_3 + 5\text{H}_2\text{O} + 4\text{Cl}_2 \rightarrow 2\text{NaHSO}_4 + 8\text{HCl}$:
 a) (Mol. wt.)/1 b) (Mol. wt.)/2 c) (Mol. wt.)/6 d) (Mol. wt.)/8
28. When Fe metal is rusted then Fe is :
 a) Oxidised b) Reduced c) Hydrolysed d) Precipitated
29. The value of n in $\text{MnO}_4^- + 8\text{H}^+ + n\text{e}^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$ is
 a) 5 b) 4 c) 2 d) 3
30. In nitric oxide (NO), the oxidation state of nitrogen is :
 a) -2 b) +1 c) -1 d) +2
31. Reaction of acidified KMnO_4 with ferrous oxalate gives oxidation products containing :
 a) Fe^{3+} b) CO_2 c) Both (a) and (b) d) None of these
32. How many litre a 0.5 N solution of an oxidising agent are reduced by 2 litre of 2.0 N solution of a reducing agent?
 a) 8 litre b) 4 litre c) 6 litre d) 7 litre
33. In which of the following oxygen shows -1 oxidation state?
 a) H_2O_2 b) CO_2 c) H_2O d) OF_2
34. The coefficients of I^- , IO_3^- and H^+ in the redox reaction, $\text{I}^- + \text{IO}_3^- + \text{H}^+ \rightarrow \text{I}_2 + \text{H}_2\text{O}$ in the balanced form respectively are
 a) 5, 1, 6 b) 1, 5, 6 c) 6, 1, 5 d) 5, 6, 1
35. Which compound shows highest oxidation number for chlorine?

- a) Show a decrease in their oxidation number during a change
 b) Gain electrons during a change
 c) Oxidise others and reduce themselves
 d) All of the above
56. One gas bleaches the colour of the flowers by reduction while the other by oxidation. The gases are :
 a) CO, Cl₂ b) H₂S, Br₂ c) SO₂, Cl₂ d) NH₃, SO₃
57. 5 g of a sample of bleaching powder is treated with excess acetic acid and KI solution. The liberated I₂ required 50 mL of N/10 hypo. The percentage of available chlorine in the sample is :
 a) 3.55 b) 7.0 c) 35.5 d) 28.2% Cl₂
58. The oxidation number of iodine in IF₅ is :
 a) +5 b) -5 c) -1 d) +1
59. The eq. wt. of FeC₂O₄ in , FeC₂O₄ → Fe³⁺ + 2CO₂ is :
 a) its mol. wt. b) mol. wt./3 c) mol. wt./4 d) None of these
60. Moles of H₂O₂ required for decolorizing 1 mole of acidified KMnO₄ are :
 a) 1/2 b) 3/2 c) 5/2 d) 7/2
61. Oxidation number of sulphur in Caro's acid is
 a) +6 b) +4 c) +8 d) +7
62. The equivalent weight of a reductant or an oxidant is given by :
 a) Eq. wt. = $\frac{\text{mol. weight of reductant or oxidant}}{\text{no. of electrons lost or gained by 1 molecule of reductant or oxidant}}$
 b) Eq. wt. = $\frac{\text{mol. wt.}}{\text{valence}}$
 c) Eq. wt. = $\frac{\text{mol. wt.}}{\text{total charge on cation or anion}}$
 d) All of the above
63. In presence of dil. H₂SO₄. The equivalent weight of KMnO₄ is :
 a) 1/5 of its molecular weight
 b) 1/6 of its molecular weight
 c) 1/10 of its molecular weight
 d) 1/2 of its molecular weight
64. Respiration is :
 a) Oxidation b) Reduction c) Both (a) and (b) d) None of these
65. $a\text{K}_2\text{Cr}_2\text{O}_7 + b\text{KCl} + c\text{H}_2\text{SO}_4 \rightarrow x\text{CrO}_2\text{Cl}_2 + y\text{KHSO}_4 + z\text{H}_2\text{O}$.
 The above equation balances when
 a) $a = 2, b = 4, c = 6$ and $x = 2, y = 6, z = 3$
 b) $a = 4, b = 2, c = 6$ and $x = 6, y = 2, z = 3$
 c) $a = 6, b = 4, c = 2$ and $x = 6, y = 3, z = 2$
 d) $a = 1, b = 4, c = 6$ and $x = 2, y = 6, z = 3$
66. Which of the following shows highest ox. no. in combined state?
 a) Os b) Ru c) Both (a) and (b) d) None of these
67. The oxidation number of sulphur in H₂S₂O₈ is :
 a) +2 b) +6 c) +7 d) +14
68. In the following reaction
 $M^{x+} + \text{MnO}_4 \rightarrow \text{MO}_3 + \text{Mn}^{2+} + \frac{1}{2}\text{O}_2$,
 If one mole of MnO₄ oxidises 2.5 moles of M^{x+} then the value of x is
 a) 5 b) 3 c) 4 d) 2
69. What volume of N K₂Cr₂O₇ solution is required to oxidise (in acid solution) a solution containing 10 g of FeSO₄? (mol.wt.of FeSO₄ = 152)
 a) 65.78 mL b) 134 mL c) 35 mL d) 33.5 mL

70. Bleaching action of chlorine in presence of moisture is :
 a) Reduction b) Oxidation c) Hydrolysis d) substitution
71. A mixture of potassium chlorate, oxalic acid and sulphuric acid is heated. During the reaction which element undergoes maximum change in the oxidation number?
 a) Cl
 b) C
 c) S
 d) H
72. Stannous chloride gives a white precipitate with a solution of mercuric chloride. In this process mercuric chloride is :
 a) Oxidized
 b) Reduced
 c) Converted into a complex compound containing Sn and Hg
 d) Converted into a chloro complex of Hg
73. In the titration of CuSO_4 vs. Hypo in presence of KI, which statement is wrong?
 a) It is iodometric titration
 b) I_2 with starch gives blue colour
 c) CuSO_4 is reduced to white Cu_2I_2 during redox change
 d) The solution before titration, on addition of KI appears blue
74. Manganese acts as strongest oxidising agent in the oxidation state
 a) +7 b) +2 c) +4 d) +5
75. The value of 'n' in the reaction
 $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + n\text{Fe}^{2+} \rightarrow 2\text{Cr}^{3+} + n\text{Fe}^{3+} + 7\text{H}_2\text{O}$
 will be
 a) 2 b) 3 c) 6 d) 7
76. In a reaction 4 mole of electrons are transferred to one mole of HNO_3 when it acts as an oxidant. The possible reduction product is :
 a) (1/2) mole N_2 b) (1/2) mole N_2O c) 1 mole of NO_2 d) 1 mole NH_3
77. The oxidation number of phosphorus in PO_4^{3-} , P_4O_{10} and $\text{P}_2\text{O}_7^{4-}$ is :
 a) +3 b) +2 c) -3 d) +5
78. In the equation ,
 $\text{CrO}_4^{2-} + \text{SO}_3^{2-} \rightarrow \text{Cr(OH)}_4 + \text{SO}_4^{2-}$
 the oxidation number of Cr changes from
 a) 6 to 4 b) 6 to 3 c) 8 to 4 d) 4 to 3
79. Oxidation numbers of P in PO_4^{3-} of S in SO_4^{2-} and that of Cr in $\text{Cr}_2\text{O}_7^{2-}$ are respectively :
 a) -3, +6 and +6 b) +5, +6 and +6 c) +3, +6 and +5 d) +5, +3 and +6
80. In alkaline condition KMnO_4 reacts as follows,
 $2\text{KMnO}_4 + 2\text{KOH} \rightarrow 2\text{K}_2\text{MnO}_4 + \text{H}_2\text{O} + \text{O}$
 Therefore, its equivalent weight will be :
 a) 31.6 b) 52.7 c) 79.0 d) 158.0
81. Oxidation number of S in SO_4^{2-}
 a) +6 b) +3 c) +2 d) -2
82. Which of the following is redox reaction?
 a) $\text{N}_2\text{O}_5 + \text{H}_2\text{O} \rightarrow 2\text{HNO}_3$
 b) $\text{AgNO}_3 + \text{KI} \rightarrow \text{AgI} + \text{KNO}_3$
 c) $\text{BaO}_2 + \text{H}_2\text{SO}_4 \rightarrow \text{BaSO}_4 + \text{H}_2\text{O}_2$
 d) $\text{SnCl}_2 + \text{HgCl}_2 \rightarrow \text{SnCl}_4 + \text{Hg}$
83. In which of the following compounds, the oxidation number of iodine is fractional?
 a) IF_3 b) IF_5 c) I_3^- d) IF_7
84. The oxidation number of Cl in KClO_3 is :

- a) +5 b) -5 c) +3 d) -3
85. The oxidation number of oxygen in $\text{KO}_3, \text{Na}_2\text{O}_2$ is
a) 3,2 b) 1,0 c) 0,1 d) -0.33,-1
86. In the reaction, $\text{I}_2 + 2\text{S}_2\text{O}_3^{2-} \rightarrow 2\text{I}^- + \text{S}_4\text{O}_6^{2-}$, Equivalent weight of iodine will be equal to:
a) Its molecular weight
b) 1/2 of its molecular weight
c) 1/4 of its molecular weight
d) Twice the molecular weight
87. The maximum oxidation number of transition metals may be:
a) +4 b) +6 c) +8 d) +10
88. The ratio of amounts of H_2S needed to precipitate all the metal ions from 100 mL 1M AgNO_3 and 100 mL of 1M CuSO_4 is :
a) 1 : 2 b) 2 : 1 c) Zero d) infinite
89. Oxidation state of sulphur in $\text{Na}_2\text{S}_2\text{O}_3$ and $\text{Na}_2\text{S}_4\text{O}_6$
a) 4 and 6 b) 3 and 5 c) 2 and 2.5 d) 6 and 6
90. Number of K^+ ions and mole of K^+ ions present in 1 litre of $\frac{N}{5}$ KMnO_4 acidified solution respectively are :
a) 0.04 and 2.4×10^{22}
b) 2.4×10^{22} and 0.04
c) 200 and 6.023×10^{23}
d) 6.023×10^{23} and 200
91. Conversion of PbSO_4 to PbS is :
a) Reduction of S b) Oxidation of S c) Dissociation d) None of these
92. Which change requires a reducing agent?
a) $\text{CrO}_4^{2-} \rightarrow \text{CrO}_7^{2-}$ b) $\text{BrO}_3^- \rightarrow \text{BrO}^-$ c) $\text{H}_2\text{O}_2 \rightarrow \text{O}_2$ d) $\text{Al(OH)}_3 \rightarrow \text{Al(OH)}_4^-$
93. In the reaction, $\text{N}_2 \rightarrow \text{NH}_3$. The eq.wt. of N_2 and NH_3 are respectively equal to :
a) $\frac{28}{3}, \frac{17}{3}$ b) $\frac{28}{6}, \frac{17}{3}$ c) $\frac{28}{2}, \frac{17}{2}$ d) $\frac{28}{5}, \frac{17}{5}$
94. Which acts as reducing agent as well as oxidising agent?
a) O_3 b) ClO_4^- c) F_2 d) MnO_4^-
95. When Cl_2 gas reacts with hot and concentrated sodium hydroxide solution, the oxidation number of chlorine changes from :
a) Zero to -1 and zero to +3
b) Zero to +1 and zero to -3
c) Zero to +1 and zero to -5
d) Zero to -1 and zero to +5
96. Which of the following is not a redox reaction?
a) $2\text{Na} + \text{Cl}_2 \rightarrow 2\text{NaCl}$ b) $\text{C} + \text{O}_2 \rightarrow \text{CO}_2$
c) $\text{AgNO}_3 + \text{NaCl} \rightarrow \text{AgCl} + \text{NaNO}_3$ d) $\text{Zn} + \text{H}_2\text{SO}_4 \rightarrow \text{ZnSO}_4 + \text{H}_2$
97. The difference in the oxidation numbers of the two types of sulphur atoms in $\text{Na}_2\text{S}_4\text{O}_6$ is
a) 4 b) 5 c) 6 d) 7
98. A compound contains atoms X, Y, Z. The oxidation number of X is +2, Y is +5 and Z is -2. The possible formula of the compound is :
a) XY_1Z_2 b) $\text{Y}_2(\text{XZ}_3)_2$ c) $\text{X}_3(\text{YZ}_4)_2$ d) $\text{X}_3(\text{Y}_4\text{Z})_2$
99. The equivalent weight of SnCl_2 in the reaction, $\text{SnCl}_2 + \text{Cl}_2 \rightarrow \text{SnCl}_4$ is :
a) 49 b) 95 c) 45 d) 59
100. What is the ox. no. of Mn in K_2MnO_4 ?
a) +4 b) +6 c) +2 d) +8
101. The stable oxidation states of Mn are :
a) +2, +3 b) +3, +7 c) +2, +7 d) +3, +5
102. 25 mL of 0.50 M H_2O_2 solution is added to 50 mL of 0.20 M KMnO_4 in acidic solution. Which of the

- following statements is true?
- a) 0.010 mole of oxygen is liberated
 b) 0.005 mole of KMnO_4 are left
 c) 0.030 g atom of oxygen gas is evolved
 d) 0.0025 mole H_2O_2 does not react with KMnO_4
103. Oxidation number of carbon in KCN is :
 a) +2 b) -2 c) +1 d) +3
104. The oxidation state of Ni in $\text{Ni}(\text{CO})_4$ is :
 a) Zero b) +4 c) +8 d) +2
105. M is the molecular weight of KMnO_4 . The equivalent weight of KMnO_4 when it is converted into K_2MnO_4 is :
 a) M b) $M/3$ c) $M/5$ d) $M/7$
106. Oxidation number of Mn in K_2MnO_4 and MnSO_4 are respectively:
 a) +7 and +2 b) +6 and +2 c) +5 and +2 d) +2 and +6
107. Which is the best description of behaviour of bromine in the reaction given below?
 $\text{H}_2\text{O} + \text{Br}_2 \rightarrow \text{HBr} + \text{HOBr}$
 a) Proton accepted only b) Both oxidised and reduced
 c) Oxidised only d) Reduced only
108. The oxidation number of P in KH_2PO_2 is :
 a) +1 b) +3 c) -3 d) +5
109. LiAlH_4 is used as :
 a) Oxidising agent b) Reducing agent c) A mordant d) Water softener
110. The brown ring complex $[\text{Fe}(\text{H}_2\text{O})_5\text{NO}^+]\text{SO}_4$ has ox.no. of Fe :
 a) +1 b) +2 c) +3 d) +4
111. The oxidation state of Fe in Fe_3O_4 is
 a) +3 b) $8/3$ c) +6 d) +2
112. In the reactions; $\text{As}_2\text{S}_3 + \text{HNO}_3 \rightarrow \text{H}_3\text{AsO}_4 + \text{H}_2\text{SO}_4 + \text{NO}$, the element oxidized is/ are :
 a) As only b) S only c) N only d) As and S both
113. The eq. wt. of KMnO_4 in the reaction, $\text{MnO}_4^- + \text{Mn}^{2+} + \text{H}_2\text{O} \rightarrow \text{MnO}_2 + \text{H}^+$ (unbalanced) is :
 a) 52.7 b) 158 c) 31.6 d) None of these
114. NO_3^- ions are converted to NH_4^+ ions by a suitable reactant. The equivalent mass of NO_3^- and NH_4^+ are :
 a) 7.75, 2.25 b) 7.75, 7.75 c) 2.25, 7.75 d) 2.25, 2.25
115. Oxidation number of chlorine in HClO_4 is :
 a) +1 b) -1 c) -7 d) +7
116. Iodine has +7 oxidation state in :
 a) HIO_4 b) H_3IO_5 c) H_5IO_6 d) all of these
117. The violent reaction between sodium and water is an example of :
 a) Reduction
 b) Oxidation
 c) Redox reaction
 d) neutralisation reaction
118. Oxidation number of Fe in $\text{K}_3[\text{Fe}(\text{CN})_6]$ is :
 a) +2 b) +3 c) +4 d) +1
119. One mole of acidified $\text{K}_2\text{Cr}_2\text{O}_7$ on reaction with excess KI will liberate.....mole(s) of I_2 .
 a) 6 b) 1 c) 7 d) 3
120. In the preparation of chlorine from HCl, MnO_2 acts as :
 a) Reducing agent b) oxidising agent c) Catalytic agent d) Dehydrating agent
121. What volume of O_2 measured at standard conditions will be formed by the action of 100 mL of 0.5 N KMnO_4 on hydrogen peroxide in an acidic solution? The skeleton equation for the reaction is,
 $\text{KMnO}_4 + \text{H}_2\text{SO}_4 + \text{H}_2\text{O}_2 \rightarrow \text{KHSO}_4 + \text{MnSO}_4 + \text{H}_2\text{O} + \text{O}_2$:

- a) 0.12 litre b) 0.28 litre c) 0.56 litre d) 1.12 litre
122. Which quantities are conserved in all oxidation-reduction reactions?
 a) Charge only b) Mass only
 c) Both charge and mass d) Neither charge nor mass
123. Which substance serves as a reducing agent in the following reaction,
 $14\text{H}^+ + \text{Cr}_2\text{O}_7^{2-} + 3\text{Ni} \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O} + 3\text{Ni}^{2+}$?
 a) H_2O b) Ni c) H^+ d) $\text{Cr}_2\text{O}_7^{2-}$
124. Which of the following chemical reactions depicts the oxidising behaviour of H_2SO_4 ?
 a) $2\text{HI} + \text{H}_2\text{SO}_4 \rightarrow \text{I}_2 + \text{SO}_2 + 2\text{H}_2\text{O}$ b) $\text{Ca}(\text{OH})_2 + \text{H}_2\text{SO}_4 \rightarrow \text{CaSO}_4 + 2\text{H}_2\text{O}$
 c) $\text{NaCl} + \text{H}_2\text{SO}_4 \rightarrow \text{NaHSO}_4 + \text{HCl}$ d) $2\text{PCl}_5 + \text{H}_2\text{SO}_4 \rightarrow 2\text{POCl}_3 + 2\text{HCl} + \text{SO}_2\text{Cl}_2$
125. In the aluminothermic process, aluminium acts as :
 a) An oxidising agent b) A flux c) A reducing agent d) A solder
126. In the reaction, $\text{SO}_2 + 2\text{H}_2\text{S} \rightarrow 3\text{S} + 2\text{H}_2\text{O}$ the substance that oxidizes is,
 a) H_2S b) SO_2 c) S d) H_2O
127. The oxidation number of sulphur in S_8 , S_2F_2 , H_2S respectively are :
 a) 0, +1 and -2 b) +2, +1 and -2 c) 0, +1 and +2 d) -2, +1 and -2
128. Maximum oxidation state is present in :
 a) CrO_2Cl_2 and MnO_4^-
 b) MnO_2
 c) $[\text{Fe}(\text{CN})_6]^{3-}$ and $[\text{Co}(\text{CN})_6]^{3-}$
 d) MnO
129. With which element oxygen shows positive oxidation state in its compounds?
 a) Na b) Cl c) N d) F
130. What is the oxidation number of chlorine in ClO_3^- ?
 a) +5 b) +3 c) +4 d) +2
131. NaClO solution reacts with H_2SO_3 as, $\text{NaClO} + \text{H}_2\text{SO}_3 \rightarrow \text{NaCl} + \text{H}_2\text{SO}_4$
 A solution of NaClO used in the above reaction contained 15 g of NaClO per litre. The normality of the solution would be :
 a) 0.40 b) 0.20 c) 0.60 d) 0.80
132. In sodium hydride, oxidation state of sodium is :
 a) Zero b) +1 c) -1 d) +2
133. The oxidation number of xenon in XeOF_2 is
 a) Zero b) 2 c) 4 d) 3
134. Which is not a redox reaction?
 a) $\text{H}_2 + \text{Br}_2 \rightarrow 2\text{HBr}$
 b) $\text{NH}_4\text{Cl} \rightarrow \text{NH}_3 + \text{HCl}$
 c) $\text{NH}_4\text{NO}_3 \rightarrow \text{N}_2\text{O} + 2\text{H}_2\text{O}$
 d) $\text{Fe} + \text{S} \rightarrow \text{FeS}$
135. In $\text{C} + \text{H}_2\text{O} \rightarrow \text{CO} + \text{H}_2$; H_2O acts as :
 a) Oxidant b) Reductant c) Both (a) and (b) d) None of these
136. Millimole of a solute in a solution can be given by :
 a) $M \times V_{\text{in litre}}$ b) $M \times V_{\text{in mL}}$ c) $\frac{\text{wt.}}{\text{mol. wt.}} \times 1000$ d) Both (b) and (c)
137. The oxidation number of carbon in $\text{H}_2\text{C}_2\text{O}_4$ is :
 a) +2 b) +3 c) +4 d) +1
138. What is the oxidation state of P in $\text{Ba}(\text{H}_2\text{PO}_2)_2$?
 a) +1 b) +2 c) +3 d) -1
139. Oxidation state of +1 for phosphorus is found in :
 a) H_3PO_3 b) H_3PO_4 c) H_3PO_2 d) $\text{H}_4\text{P}_2\text{O}_7$
140. Oxidation number of S in $(\text{CH}_3)_2\text{SO}$ is :
 a) Zero b) +1 c) +2 d) +3

141. In which reaction the underlined substance has been reduced?
- Carbon monoxide + copper oxide \rightarrow carbon dioxide + copper
 - Copper oxide + hydrochloric acid \rightarrow water + copper chloride
 - Steam + iron \rightarrow hydrogen + iron oxide
 - Hydrogen + iron oxide \rightarrow water + iron
142. The decomposition of KClO_3 to KCl and O_2 on heating is an example of :
- Intermolecular redox change
 - Intramolecular redox change
 - Disproportionation or auto redox change
 - None of the above
143. Mohr's salt is oxidised to in presence of acidized KMnO_4 .
- Fe^{2+}
 - Fe^{3+}
 - Fe
 - None of these
144. Fluorine is a strong oxidising agent because :
- It has several isotopes
 - It is very small and has 7 electrons in valency shell
 - Its valency is one
 - It is the first member of the halogen series
145. In the conversion of Br_2 to BrO_3^- , the oxidation number of Br changes from
- Zero to +5
 - +1 to +5
 - Zero to -3
 - +2 to +5
146. The oxidation number of Cr in CrO_5 is
- +3
 - +5
 - +6
 - 0
147. An indicator used for redox reaction is itself :
- Either an oxidant or a reductant
 - Neither an oxidant nor a reductant
 - Acid or base
 - None of the above
148. CrO_5 reacts with H_2SO_4 to give $\text{Cr}_2(\text{SO}_4)_3$, H_2O and O_2 . Moles of O_2 liberated by 1 mole of CrO_5 in this reaction are :
- 2.5
 - 1.25
 - 4.5
 - 1.75
149. In the following reaction, $4\text{P} + 3\text{KOH} + 3\text{H}_2\text{O} \rightarrow 3\text{KH}_2\text{PO}_2 + \text{PH}_3$
- P is only oxidized
 - P is only reduced
 - P is both oxidized as well as reduced
 - None of the above
150. Oxidation number of P in $\text{P}_2\text{O}_7^{4-}$ is :
- +3
 - +4
 - +5
 - +6
151. In the conversion of $\text{K}_2\text{Cr}_2\text{O}_7$ to K_2CrO_4 the oxidation number of chromium :
- Increases
 - Remains the same
 - Decreases
 - None of these
152. In which of the following, the oxidation number of oxygen has been arranged in increasing order?
- $\text{OF}_2 < \text{KO}_2 < \text{BaO}_2 < \text{O}_3$
 - $\text{BaO}_2 < \text{KO}_2 < \text{O}_3 < \text{OF}_2$
 - $\text{BaO}_2 < \text{O}_3 < \text{OF}_2 < \text{KO}_2$
 - None of these
153. Oxidation number of sodium in sodium amalgam is :
- +2
 - +1
 - 2
 - zero
154. The apparatus in which standard solution is prepared is known as :
- Measuring flask
 - Round bottom flask
 - Burette
 - None of these
155. $\text{K}_3\text{Fe}(\text{CN})_6$ is used as Indicator for FeSO_4 vs. $\text{K}_2\text{Cr}_2\text{O}_7$ titrations.
- Self
 - External
 - Internal
 - Not an
156. The oxidation number of N in N_2H_5^+ is :
- 2
 - +3
 - +2
 - 3
157. Which can act as oxidant?
- H_2O_2
 - H_2S
 - NH_3
 - None of these
158. What weight of HNO_3 is needed to convert 5 g of iodine into iodic acid according to the reaction, $\text{I}_2 +$

- $\text{HNO}_3 \rightarrow \text{HIO}_3 + \text{NO}_2 + \text{H}_2\text{O} ?$
- a) 12.4 g b) 24.8 g c) 0.248 g d) 49.6 g
159. In which SO_2 acts as oxidant, while reacting with :
- a) Acidified KMnO_4 b) Acidified $\text{K}_2\text{Cr}_2\text{O}_7$ c) H_2S d) Acidified $\text{C}_2\text{H}_5\text{OH}$
160. HBr and HI reduce H_2SO_4 , HCl can reduce KMnO_4 and HF can reduce:
- a) H_2SO_4 b) $\text{K}_2\text{Cr}_2\text{O}_7$ c) KMnO_4 d) None of these
161. Equivalent mass of $\text{Na}_2\text{S}_2\text{O}_3$ in its reaction with I_2 is equal to :
- a) Molar mass b) Molar mass / 2 c) Molar mass / 3 d) Molar mass / 4
162. Which of the following change represents a disproportionation reaction(s)?
- a) $\text{Cl}_2 + 2\text{OH}^- \rightarrow \text{ClO}^- + \text{Cl}^- + \text{H}_2\text{O}$
 b) $\text{Cu}_2\text{O} + 2\text{H}^+ \rightarrow \text{Cu} + \text{Cu}^{2+} + \text{H}_2\text{O}$
 c) $2\text{HCuCl}_2 \xrightarrow[\text{water}]{\text{Dilution with}} \text{Cu} + \text{Cu}^{2+} + 4\text{Cl}^- + 2\text{H}^+$
 d) All of the above
163. Oxidation number of 'N' in N_3H (hydrazoic acid) is
- a) $-\frac{1}{3}$ b) +3 c) 0 d) -3
164. Ceric ammonium sulphate and potassium permanganate are used as oxidising agents in acidic medium for oxidation of ferrous ammonium sulphate to ferric sulphate. The ratio of number of moles of ceric ammonium sulphate required per mole of ferrous ammonium sulphate to the number of moles of KMnO_4 required per mole of ferrous ammonium sulphate, is
- a) 5.0 b) 0.2 c) 0.6 d) 2.0
165. Eq.wt. of NH_3 in, $\text{NH}_3 + \text{O}_2 \rightarrow \text{NO} + \text{H}_2\text{O}$ is :
- a) 3.4 b) 17 c) 8.5 d) None of these
166. Carbon is in the lowest oxidation state in :
- a) CH_4 b) CCl_4 c) CO_2 d) CF_4
167. When the ion $\text{Cr}_2\text{O}_7^{2-}$ acts as an oxidant in acidic aqueous solution the ion Cr^{3+} is formed. How many mole of Sn^{2+} would be oxidised to Sn^{4+} by one of $\text{Cr}_2\text{O}_7^{2-}$ ions?
- a) 2/3 b) 3/2 c) 2 d) 3
168. 100 mL of 0.1 M solution of a reductant is diluted to 1 litre, which of the following changes?
- a) Molarity b) Millimole c) Milliequivalent d) None of these
169. If H_2S is passed through an acidified $\text{K}_2\text{Cr}_2\text{O}_7$ solution, the colour of the solution :
- a) Will remain unchanged
 b) Will change to deep red
 c) Will change to dark green
 d) Will change to dark brown
170. Ozone tails mercury. The reaction isof Hg.
- a) Reduction b) Oxidation c) Substitution d) None of these
171. The oxidation number of Cr in $[\text{Cr}(\text{NH}_3)_4\text{Cl}_2]^+$ is :
- a) +3 b) +2 c) +1 d) zero
172. In the reaction, $\text{VO} + \text{Fe}_2\text{O}_3 \rightarrow \text{FeO} + \text{V}_2\text{O}_5$. The eq.wt. of V_2O_5 is equal to its :
- a) mol. wt. b) mol. wt./8 c) mol. wt./6 d) None of these
173. The eq. wt. of K_2CrO_4 as an oxidising agent in acid medium is :
- a) (mol. wt.)/2 b) $(2 \times \text{mol. wt.})/3$ c) (mol. wt.)/3 d) (mol. wt.)/6
174. Which reaction involves neither oxidation nor reduction?
- a) $\text{CrO}_4^{2-} \rightarrow \text{Cr}_2\text{O}_7^{2-}$ b) $\text{Cr} \rightarrow \text{CrCl}_3$ c) $\text{Na} \rightarrow \text{Na}^+$ d) $2\text{S}_2\text{O}_3^{2-} \rightarrow \text{S}_4\text{O}_6^{2-}$
175. The number of equivalent per mole of H_2S used in its oxidation to SO_2 is :
- a) 3 b) 6 c) 4 d) 2
176. Oxidation number of sulphur in Na_2SO_4 is :
- a) +2 b) +4 c) +6 d) -2
177. Which can have both +ve and -ve oxidation states?

193. In which of the following oxidation number of chlorine is +5?
 a) HClO b) HClO₂ c) HClO₃ d) HClO₄
194. In the reaction, $\text{Zn} + 2\text{H}^+ + 2\text{Cl}^- \rightarrow \text{Zn}^{2+} + 2\text{Cl}^- + \text{H}_2$, the spectator ion is :
 a) Cl⁻ b) Zn²⁺ c) H⁺ d) All of these
195. Turn bull's blue is :
 a) Fe₃[Fe(CN)₆]₂ b) K₄Fe(CN)₆ c) K₃Fe(CN)₆ d) Na₄Fe(CN)₆
196. The oxidation state shown by silicon when it combines with strongly electropositive metals is
 a) -2 b) -4 c) +4 d) +2
197. The compound that can work both as an oxidising and reducing agent is:
 a) KMnO₄ b) H₂O₂ c) Fe₂(SO₄)₃ d) K₂Cr₂O₇
198. An element A in a compound ABD has oxidation number Aⁿ⁻. It is oxidized by Cr₂O₇²⁻ in acidic medium. In the experiment 1.68 × 10⁻³ mole of K₂Cr₂O₇ were used for 3.26 × 10⁻³ mole of ABD. The new oxidation number of A after oxidation is :
 a) 3 b) 3 - n c) n - 3 d) +n
199. The burning of hydrogen is called :
 a) Hydrogenation b) Hydration c) Oxidation d) reduction
200. Oxidation number of chlorine in chlorine heptaoxide is :
 a) +1 b) +4 c) +6 d) +7
201. The correct order of reducing power of halide ions is :
 a) Cl⁻ > Br⁻ > I⁻ > F⁻
 b) Cl⁻ > I⁻ > Br⁻ > F⁻
 c) Br⁻ > Cl⁻ > I⁻ > F⁻
 d) I⁻ > Br⁻ > Cl⁻ > F⁻
202. The reaction, $3\text{ClO}^-(aq) \rightarrow \text{ClO}_3^-(aq) + 2\text{Cl}^-(aq)$ is an example of :
 a) Oxidation reaction
 b) Reduction reaction
 c) Disproportionation reaction
 d) Decomposition reaction
203. The ox.no. of S in Na₂S₄O₆ is :
 a) + 2.5
 b) +2 and +3 (two S have +2 and other two have +3)
 c) +2 and +3 (three S have +2 and one S has +3)
 d) +5 and 0 (two S have +5 and the other two S have 0)
204. Oxidation is a process which involves :
 a) de-electronation b) Electronation c) Addition of hydrogen d) Addition of metal
205. A student states that heating of limestone is an oxidation process, the reason he gives that an oxide of the metal is produced on heating. Which one is correct?
 a) The statement and reason are true
 b) The statement and reason are wrong
 c) The statement is true but the reason is false
 d) None of the above
206. A sulphur containing species that cannot be an oxidising agent is :
 a) H₂SO₄ b) H₂S c) SO₂ d) H₂SO₃
207. KMnO₄ acts as indicator in its redox titrations.
 a) Self b) External c) Internal d) Not an
208. In a reaction between zinc and iodine in which zinc iodide is formed, which is oxidised?
 a) Zinc ions b) Iodide ions c) Zinc atom d) Iodine
209. The best oxidising agent of the oxygen family is:
 a) Tellurium b) Selenium c) Sulphur d) Oxygen
210. The oxidation state of iron in sodium nitroprusside is :

- a) +2 b) +1 c) Zero d) +3
211. A compound of Xe and F is found to have 53.3% Xe. Oxidation number of Xe in this compound is :
a) -4 b) Zero c) +4 d) +6
212. Which combination is odd with respect to oxidation numbers of S, Cr, N and H respectively:
a) $\text{H}_2\text{SO}_5, \text{H}_2\text{S}_2\text{O}_8, \text{H}_2\text{SO}_4, \text{SF}_6$
b) $\text{K}_2\text{Cr}_2\text{O}_7, \text{K}_2\text{CrO}_4, \text{CrO}_5, \text{CrO}_2\text{Cl}_2$
c) $\text{NH}_3, \text{NH}_4^+, \text{N}_3\text{H}, \text{NO}_2^-$
d) $\text{CaH}_2, \text{NaH}, \text{LiH}, \text{MgH}_2$
213. 0.2 g of a sample of H_2O_2 required 10 mL of N KMnO_4 in a titration in the presence of H_2SO_4 . Purity of H_2O_2 is :
a) 25% b) 85% c) 65% d) 95%
214. When KMnO_4 as oxidising agent and ultimately forms $\text{MnO}_4^{2-}, \text{Mn}_2\text{O}_3$ and Mn^{2+} , the number of electrons transferred per mole of KMnO_4 each case respectively is :
a) 4, 3, 1, 5 b) 1, 5, 3, 7 c) 1, 3, 4, 5 d) 1, 3, 8, 5
215. Titration of KI with H_2O_2 in presence of acid is a :
a) Clock reaction b) Redox reaction c) Intermolecular redox d) All of these
216. Oxidation state of nitrogen is incorrectly given for :
Compound **Oxidation state**
a) $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$ -3
b) NH_2OH -1
c) $(\text{N}_2\text{H}_5)_2\text{SO}_4$ +2
d) Mg_3N_2 -3
217. Fluorine exhibits only -1 oxidation state, while iodine exhibits oxidation states of -1, +1, +3, +5 and +7. This is due to :
a) Fluorine being a gas
b) Available d -orbitals in iodine
c) Non-availability of d -orbitals in iodine
d) None of the above
218. Elements which generally exhibit multiple oxidation states and whose ions are coloured are known as :
a) Metalloid b) Non-metals c) Metals d) Transition metals
219. The oxidation state of sulphur in sodium tetrathionate ($\text{Na}_2\text{S}_4\text{O}_6$) is
a) 2 b) 0 c) 2.5 d) 3.5
220. Which is strongest oxidising agent?
a) O_3 b) O_2 c) Cl_2 d) F_2
221. Sulphur has the highest oxidation state in :
a) SO_2 b) SO_3 c) H_2SO_3 d) H_2S
222. Nitrogen has fractional oxidation number in :
a) N_2H_4 b) NH_4 c) HN_3 d) N_2F_2
223. As the oxidation state for any metal increases, the tendency to show ionic nature:
a) Decreases b) Increases c) Remains same d) None of these
224. In acid medium Zn reduces nitrate ion to NH_4^+ ion according to the reaction
 $\text{Zn} + \text{NO}_3^- + \text{Zn}^{2+} + \text{NH}_4^+ + \text{H}_2\text{O}$ (unbalanced)
How many moles of HCl are required to reduce half a mole of NaNO_3 completely? Assume the availability of sufficient Zn.
a) 5 b) 4 c) 3 d) 2
225. Weight of FeSO_4 (mol. wt. = 152) oxidized by 200 mL of 1 N KMnO_4 solution is :
a) 30.4 g b) 15.2 g c) 60.8 g d) 158 g
226. In the ionic equation,
 $\text{BiO}_3^- + 6\text{H}^+ + xe^- \rightarrow \text{Bi}^{3+} + 3\text{H}_2\text{O}$
The values of x is

- a) Mn^{4+} b) Mn^{2+} c) Mn^{6+} d) Mn
245. In balancing the half reaction, $\text{S}_2\text{O}_3^{2-} \rightarrow \text{S}(s)$, the number of electrons that must be added is :
 a) 2 on the right b) 2 on the left c) 3 on the right d) 4 on the left
246. What volume of 0.1 M KMnO_4 is needed to oxidise 100 mg of FeC_2O_4 in acidic solution?
 a) 4.1 mL b) 8.2 mL c) 10.2 mL d) 4.6 mL
247. Which one is not a redox titration?
 a) FeSO_4 vs. $\text{K}_2\text{Cr}_2\text{O}_7$ b) CuSO_4 vs. hypo c) I_2 vs. hypo d) AgNO_3 vs. KCl
248. A 0.518 g sample of lime stone is dissolved in HCl and then the calcium is precipitated as CaC_2O_4 . After filtering and washing the precipitate, it requires 40.0 mL of 0.250 N KMnO_4 , solution acidified with H_2SO_4 to titrate is as, $\text{MnO}_4^- + \text{H}^+ + \text{C}_2\text{O}_4^{2-} \rightarrow \text{Mn}^{2+} + \text{CO}_2 + 2\text{H}_2\text{O}$. The percentage of CaO in the sample is :
 a) 54.0 % b) 27.1 % c) 42% d) 84%
249. The missing term in following equation is : $2\text{Fe}^{3+}(aq) + \text{Sn}^{2+}(aq) \rightarrow 2\text{Fe}^{2+}(aq) + ?$
 a) Sn^{4+} b) Sn^{2+} c) Sn d) None of these
250. Reaction of Br_2 with Na_2CO_3 in aqueous solution gives sodium bromide and sodium bromate with evolution of CO_2 gas. The number of sodium bromide molecules involved in the balanced chemical equation is
 a) 1 b) 3 c) 5 d) 7
251. Oxidation number of carbon in C_3O_2 , Mg_2C_3 are respectively :
 a) $-4/3, +4/3$ b) $+4/3, -4/3$ c) $-2/3, +2/3$ d) $-2/3, +4/3$
252. The reaction; $\text{KI} + \text{I}_2 \rightarrow \text{KI}_3$ shows :
 a) Oxidation b) Reduction c) Complex formation d) All of these
253. The oxidation state of Cr in chromium trioxide is
 a) +3 b) +4 c) +5 d) +6
254. Oxidation number of S in S_2Cl_2 is :
 a) +1 b) +6 c) Zero d) -1
255. In which of the following N has lowest oxidation number?
 a) NO b) NO_2 c) N_2O d) N_2O_5
256. 2 mole of FeSO_4 are oxidized by 'X' mole of KMnO_4 whereas 2 mole of FeC_2O_4 are oxidized by 'Y' mole of KMnO_4 . The ration f 'X' and 'Y' is :
 a) 1 : 3 b) 1 : 2 c) 1 : 4 d) 1 : 5
257. H_2S reacts with halogens, the halogens :
 a) Are oxidised b) Are reduced c) Form sulphur halides d) None of these
258. In an experiment 50 mL of 0.1 M solution of a salt reacted with 25 mL of 0.1 M solution of sodium sulphite. The half equation for the oxidation of sulphite ion is :
 $\text{SO}_3^{2-}(aq) + \text{H}_2\text{O}(l) \rightarrow \text{SO}_4^{2-}(aq) + 2\text{H}^+(aq) + 2e^-$
 If the oxidation number of metal in the salt was 3, what would be the new oxidation number of metal?
 a) Zero b) 1 c) 2 d) 4
259. The most stable oxidation state of copper is :
 a) +2 b) +1 c) +3 d) +4
260. White phosphorus reacts with caustic soda, the products are PH_3 and NaH_2PO_2 . This reaction is an example of
 a) Oxidation b) Reduction c) Disproportionation d) Neutralisation
261. When a sulphur atom becomes a sulphide ion :
 a) It gains two electrons
 b) The mass number changes
 c) There is no change in the composition of atom
 d) None of the above
262. Titre value is the volume of titrant used for a definite amount of unknown reagent at its :
 a) Equivalence point b) End point c) Neutralization point d) All of these
263. Oxidation states of X, Y, Z are +2, +5 and -2 respectively. Formula of the compound formed by these wii

- be
- a) X_2YZ_6 b) XY_2Z_6 c) XY_5 d) X_3YZ_4
264. In which compound, oxygen has an oxidation state of +2 ?
 a) H_2O_2 b) H_2O c) OF_2 d) CO
265. If equal volumes of 1M $KMnO_4$ and 1 M $K_2Cr_2O_7$ solutions are allowed to oxidise F^{2+} to F^{3+} in acidic medium volume of oxidant required for one mole of F^{2+} will be :
 a) $V_{KMnO_4} > V_{K_2Cr_2O_7}$
 b) $V_{KMnO_4} < V_{K_2Cr_2O_7}$
 c) $V_{KMnO_4} = V_{K_2Cr_2O_7}$
 d) Nothing can be predicted
266. How many gram of $KMnO_4$ should be taken to make up 250 mL of a solution of such strength that 1 mL is equivalent to 5.0 mg of Fe in $FeSO_4$?
 a) 1.414 g b) 0.70 g c) 3.16 g d) 1.58 g
267. The oxidation number of Cr in K_2CrO_4 is
 a) +3 b) -6 c) +6 d) -3
268. In the reaction, $2Na_2S_2O_3 + I_2 \rightarrow Na_2S_4O_6 + 2NaI$, the oxidation state of sulphur is :
 a) Decreased b) Increased c) Unchanged d) None of these
269. The equivalent weight of $KMnO_4$ (acidic medium) is (at. wt. of K = 39; Mn = 55) :
 a) 158 b) 15.8 c) 31.6 d) 3.16
270. The oxidation number of chromium in potassium dichromate is
 a) +2 b) +4 c) +6 d) +8
271. The equivalent weight of $MnSO_4$ is half of its molecular weight when it is converted to :
 a) Mn_2O_3 b) MnO_2 c) MnO_4^- d) Mn_4^{2-}
272. Aqueous solution of SO_2 reacts with H_2S to precipitate sulphur. Here SO_2 acts as :
 a) Catalyst b) Reducing agent c) Oxidizing agent d) Acid
273. Saline hydrides are :
 a) Strong oxidants
 b) Strong reductants
 c) Strong dehydrating agents
 d) Strong bleaching agents
274. State the oxidation number of carbonyl carbon in methanal and methanoic acid respectively
 a) 0 and 0 b) 0 and +2 c) +1 and +2 d) +1 and +3
275. The eq. wt. of I_2 in the change $I_2 \rightarrow IO_3^-$ is :
 a) 12.7 b) 63.5 c) 25.4 d) 2.54
276. Equivalent mass of oxidizing agent in the reaction is.
 $SO_2 + 2H_2S \rightarrow 3S + 2H_2O$
 a) 32 b) 64 c) 16 d) 8
277. In a conjugate pair of reductant and oxidant, the reductant has :
 a) Lower ox.no. b) Higher ox.no. c) Same ox.no. d) Either of these
278. In which of the following reactions, hydrogen is acting as an oxidising agent?
 a) With Li to form LiH b) With I_2 to give HI c) With S to give H_2S d) None of the above
279. The number of moles of Mohr's salt required per mole of dichromate ion is :
 a) 3 b) 4 c) 5 d) 6
280. For reducing one mole of Fe^{2+} ion to Fe, the number of faraday of electricity is :
 a) 2 b) 1 c) 1.5 d) 4
281. $Co(s) + Cu^{2+}(aq) \rightarrow Co^{2+}(aq) + Cu(s)$. This reaction is :
 a) Oxidation reaction b) Reduction reaction c) Redox reaction d) None of these
282. The oxidation state of I in $H_4IO_6^-$ is :
 a) +7 b) -1 c) +5 d) +1
283. The oxidation number of N in NH_3 is :

- a) -3 b) +3 c) Zero d) 5
284. Mn^{2+} can be converted into Mn^{7+} by reacting with
a) SO_2 b) Cl_2 c) PbO_2 d) SnCl_2
285. The oxidation number of Ni in $\text{K}_4[\text{Ni}(\text{CN})_4]$ is :
a) +1 b) +2 c) -1 d) 0
286. Which change occur when lead monoxide is converted into lead nitrate?
a) Oxidation
b) Reduction
c) Neither oxidation nor reduction
d) Both oxidation and reduction
287. How many mole of electron are involved in the reduction of one mole of MnO_4^- ion in alkaline medium to MnO_3^- ?
a) 2 b) 1 c) 3 d) 4
288. The oxidation number of Fe in $\text{K}_4\text{Fe}(\text{CN})_6$ is :
a) +2 b) +3 c) +4 d) +6
289. For the reaction, $\text{NH}_3 + \text{OCl}^- \rightarrow \text{N}_2\text{H}_4 + \text{Cl}^-$
occurring in basic medium, the coefficient of N_2H_4 in the balanced equation will be
a) 1 b) 2 c) 3 d) 4
290. In the reaction $\text{H}_2\text{O} + \text{H}_2\text{O}_2 \rightarrow \text{S} + 2\text{H}_2\text{O}$
a) H_2S is an acid and H_2O_2 is a base
b) H_2S is a base and H_2O_2 is an acid
c) H_2S is an oxidising agent and H_2O_2 is a reducing agent
d) H_2S is a reducing agent and H_2O_2 is an oxidising agent
291. When H_2SO_3 is converted into H_2SO_4 the change in the oxidation state of sulphur is from:
a) 0 to +2 b) +2 to +4 c) +4 to +2 d) +4 to +6
292. The oxidation number of nitrogen in NH_2OH is :
a) +1 b) -1 c) -3 d) -2
293. In the reaction, $2\text{CuSO}_4 + 4\text{KI} \rightarrow \text{Cu}_2\text{I}_2 + 2\text{K}_2\text{SO}_4 + \text{I}_2$ The ratio of equivalent weight of CuSO_4 to its molecular weight is :
a) $1/8$ b) $1/4$ c) $1/2$ d) 1
294. In the reaction between acidified $\text{K}_2\text{Cr}_2\text{O}_7$ and iron (II) ions shown by the equation : $\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 6\text{Fe}^{2+}(\text{aq}) + 14\text{H}^+(\text{aq}) \rightarrow 2\text{Cr}^{3+}(\text{aq}) + 7\text{H}_2\text{O}(\text{l}) + 6\text{Fe}^{3+}(\text{aq})$
a) The colour of the solution changes from green to blue
b) The iron (II) ions are reduced
c) The dichromate ions are reduced
d) Hydrogen ions are reduced
295. Which is the reducing agent in the reaction, $8\text{H}^+ + 4\text{NO}_3^- + 6\text{Cl}^- + \text{Sn}(\text{s}) \rightarrow \text{SnCl}_6^{2-} + 4\text{NO}_2 + 4\text{H}_2\text{O}$?
a) $\text{Sn}(\text{s})$ b) Cl^- c) NO_3^- d) $\text{NO}_2(\text{g})$
296. Which is a redox reaction?
a) $\text{H}_2\text{SO}_4 + 2\text{NaOH} \rightarrow \text{Na}_2\text{SO}_4 + 2\text{H}_2\text{O}$
b) $\text{BaCl}_2 + \text{H}_2\text{SO}_4 \rightarrow \text{BaSO}_4 + 2\text{HCl}$
c) $\text{CH}_3\text{COOH} + \text{C}_2\text{H}_5\text{OH} \rightarrow \text{CH}_3\text{COOC}_2\text{H}_5 + \text{H}_2\text{O}$
d) $2\text{FeCl}_3 + \text{SnCl}_2 \rightarrow 2\text{FeCl}_2 + \text{SnCl}_4$
297. Which one of the following reactions involves disproportionation?
a) $2\text{H}_2\text{SO}_4 + \text{Cu} \rightarrow \text{CuSO}_4 + 2\text{H}_2\text{O} + \text{SO}_2$ b) $\text{As}_2\text{O}_3 + 3\text{H}_2\text{S} \rightarrow \text{As}_2\text{S}_3 + 3\text{H}_2\text{O}$
c) $2\text{KOH} + \text{Cl}_2 \rightarrow \text{KCl} + \text{KOCl} + \text{H}_2\text{O}$ d) $\text{Ca}_3\text{P}_2 + 6\text{H}_2\text{O} \rightarrow 3\text{Ca}(\text{OH})_2 + 2\text{PH}_3$
298. The oxidation state of chromium in the final product formed by the reaction between KI and acidified potassium dichromate solution is
a) +3 b) +2 c) +6 d) +4
299. Which of the following acts as an oxidising as well as reducing agent?

- a) Na_2O b) Na_2O_2 c) NaNO_3 d) NaNO_2
300. Oxidation state of carbon in graphite is:
 a) Zero b) +1 c) +4 d) +2
301. Which compound has oxidation number of carbon equal to zero?
 a) C_6H_6 b) CH_3 c) C_2H_4 d) $\text{C}_6\text{H}_{12}\text{O}_6$
302. In the reaction, $2\text{KMnO}_4 + 16\text{HCl} \rightarrow 2\text{KCl} + 2\text{MnCl}_2 + 8\text{H}_2\text{O} + 5\text{Cl}_2$, the reduction product is :
 a) Cl_2 b) MnCl_2 c) KCl d) H_2O
303. The oxidation number of phosphorus in $\text{Mg}_2\text{P}_2\text{O}_7$ is :
 a) +5 b) -5 c) +6 d) -7
304. 1 mole of chlorine combines with a certain weight of a metal giving 111 g of its chloride. The atomic weight of the metal (assuming its valency to be 2) is :
 a) 40 b) 20 c) 80 d) None of these
305. Oxidation state of chromium
-
- a) +10 b) +6 c) +3 d) +2
306. Oxidation states of the metal in the minerals haematite and magnetite, respectively, are
 a) II, III in haematite and III in magnetite b) II, III in haematite and II in magnetite
 c) II in haematite and II, III in magnetite d) III in haematite and II, III in magnetite
307. The colour of $\text{K}_2\text{Cr}_2\text{O}_7$ changes from red-orange to lemon-yellow on treatment with $\text{KOH}(aq)$ because of :
 a) Reduction of Cr(VI) to Cr(III)
 b) Formation of chromium hydroxide
 c) Conversion of dichromate into chromate ion
 d) Oxidation of potassium hydroxide to potassium peroxide
308. How many electrons are involved in oxidation of KMnO_4 in basic medium?
 a) 1 b) 2 c) 5 d) 3
309. The oxidation state of nitrogen in NH_4NO_3 is :
 a) -3 and +5 b) +3 and +5 c) +5 d) +3
310. When Sn(IV) chloride is treated with excess HCl, the complex $[\text{SnCl}_6]^{2-}$ is formed. The oxidation state of Sn in this complex is:
 a) +6 b) -2 c) +4 d) -5
311. Oxidation number of chlorine in HOCl is :
 a) Zero b) -1 c) +1 d) +2
312. In the reaction, $\text{C} + 4\text{HNO}_3 \rightarrow \text{CO}_2 + 2\text{H}_2\text{O} + 4\text{NO}_2$, HNO_3 acts as :
 a) An oxidising agent
 b) An acid
 c) An acid as well as oxidising agent
 d) A reducing agent
313. Change of hydrogen into proton is :
 a) Oxidation of hydrogen
 b) Acid-base reaction
 c) Reduction of hydrogen
 d) Displacement reaction
314. 8 g of sulphur are burnt to form SO_2 which is oxidised by Cl_2 water. The solution is treated with BaCl_2 solution. The amount of BaSO_4 precipitated is :
 a) 1.0 mole b) 0.5 mole c) 0.24 mole d) 0.25 mole
315. The number of mole of ferrous oxalate oxidised by one mole of KMnO_4 is:
 a) 1/5 b) 3/5 c) 2/3 d) 5/3

316. Reactants react in the equal number of to give products.
 a) Mole b) Weights c) Equivalent d) All of these
317. Mole and millimole of reactants react in theas represented by balanced stoichiometric equation.
 a) Molar ratio b) Equal amount c) Both (a) and (b) d) None of these
318. The reaction of white phosphorus with aqueous NaOH gives phosphine along with another phosphorus containing compound. The reaction type the oxidation states of phosphorus in phosphine and the other product are respectively :
 a) Redox reaction; -3 and -5
 b) Redox reaction; $+3$ and $+5$
 c) Disproportionation reaction; -3 and $+1$
 d) Disproportionation reaction; -3 and $+3$
319. Which can act only as oxidising agent?
 a) Oxygen b) Fluorine c) Iodine d) H_2O_2
320. For the reaction : $N_2 + 3H_2 \rightarrow 2NH_3$; if E_1 and E_2 are equivalent masses of NH_3 and N_2 respectively, then $E_1 - E_2$ is :
 a) 1 b) 2 c) 3 d) 4
321. Bleaching action of SO_2 is due to :
 a) Reduction b) Oxidation c) Hydrolysis d) Acidic nature
322. In $N_2 + 2H_2O \rightarrow NH_4^+ + NO_2^-$; N is :
 a) Oxidised b) Reduced c) Both (a) and (b) d) None of these
323. If three electrons are lost by a metal ion M^{3+} , its final oxidation number will be :
 a) Zero b) $+6$ c) $+2$ d) $+4$
324. In the reaction, $NaH + H_2O \rightarrow NaOH + H_2$:
 a) H^- is oxidised
 b) Na^+ is reduced
 c) Both NaH and H_2O are reduced
 d) None of the above
325. Which of the following acts as an oxidizing agent?
 a) HNO_3 b) Cl_2 c) $FeCl_3$ d) All of these
326. How many gram of I_2 are present in a solution which requires 40 mL, of 0.11 N $Na_2S_2O_3$ to react with it, $S_2O_3^{2-} + I_2 \rightarrow S_4O_6^{2-} + 2I^-$?
 a) 12.7 g b) 0.558 g c) 25.4 g d) 11.4 g
327. The number of mole of $KMnO_4$ that will be needed to react with one mole of sulphite ion in acidic solution is :
 a) $2/5$ b) $3/5$ c) $4/5$ d) 1
328. What weight of HNO_3 is required to make 1 litre of 2 N solution to be used as an oxidising agent in the reaction? $3Cu + 8HNO_3 \rightarrow 3Cu(NO_3)_2 + 2NO + 4H_2O$
 a) 63 g b) 21 g c) 42 g d) 84 g
329. The oxidation state of two sulphur atoms in $H_2S_2O_8$
 a) -6 b) -2 c) $+6$ d) -4
330. In a conjugate pair of reductant and oxidant, the oxidant has :
 a) Higher ox.no. b) Lower ox.no. c) Same ox.no. d) Either of these
331. In the equation, $H_2S + 2HNO_3 \rightarrow 2H_2O + 2NO_2 + S$. The equivalent weight of hydrogen sulphide is :
 a) 17 b) 34 c) 68 d) 18
332. In which transfer of five electrons takes place?
 a) $MnO_4^- \rightarrow Mn^{2+}$ b) $CrO_4^{2-} \rightarrow Cr^{3+}$ c) $MNO_4^- \rightarrow MnO_2$ d) $Cr_2O_7^{2-} \rightarrow 2Cr^{3+}$
333. Oxidation number of nitrogen is highest in
 a) N_3H b) N_2O_4 c) NH_2OH d) NH_3
334. Starch gives blue colour with :
 a) KI b) I_2 c) Cl_2 d) None of these


335. The number of mole of potassium salt, *i. e.*, $\text{KHC}_2\text{O}_4 \cdot \text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$ oxidised by one mole of permanganate ion is :
- a) 2/5 b) 4/5 c) 1 d) 5/4
336. When an acidified solution of ferrous ammonium sulphate is treated with KMnO_4 solution, the ion which is oxidised is :
- a) Fe^{2+} b) SO_4^{2-} c) NH_4^+ d) MnO_4^-
337. Oxidation number of N in N_3H is :
- a) -3 b) +3 c) Zero d) -1/3
338. Hydrogen peroxide in aqueous solution decomposes on warming to give oxygen according to the equation, $2\text{H}_2\text{O}_2(\text{aq}) \rightarrow 2\text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g})$ under conditions where one mole of gas occupies 24 dm^3 , 100 cm^3 of *XM* solution of H_2O_2 produces 3 dm^3 of O_2 . Thus, *X* is :
- a) 2.5 b) 1 c) 0.5 d) 0.25
339. CuSO_4 and KI on mixing gives :
- a) $\text{CuI}_2 + \text{K}_2\text{SO}_4$ b) $\text{Cu}_2\text{I}_2 + \text{K}_2\text{SO}_4$ c) $\text{Cu}_2\text{I}_2 + \text{K}_2\text{SO}_4 + \text{I}_2$ d) $\text{CuI}_2 + \text{K}_2\text{SO}_4 + \text{I}_2$
340. Which metal exhibits more than one oxidation states?
- a) Na b) Mg c) Al d) Fe
341. Which of the following oxidation state is the most common among the lanthanoides :
- a) 4 b) 2 c) 5 d) 3
342. 13.5 g aluminium changes to Al^{3+} in solution by losing :
- a) 18×10^{23} electrons
b) 6.023×10^{23} electrons
c) 3.01×10^{23} electrons
d) 9×10^{23} electrons
343. In CH_2Cl_2 , the oxidation number of C is :
- a) -4 b) +2 c) Zero d) +4
344. In the compounds KMnO_4 and $\text{K}_2\text{Cr}_2\text{O}_7$, the highest oxidation state is of the element
- a) Mn b) K c) O d) Cr
345. The oxidation state of nitrogen varies from :
- a) -3 to +5 b) 0 to +5 c) -3 to 1 d) +3 to +5
346. The oxidation state of hydrogen in CaH_2 is :
- a) +1 b) -1 c) Zero d) +2
347. The most common oxidation state of an element is -2. The number of electrons present in its outermost shell is :
- a) 2 b) 4 c) 6 d) 8
348. A good indicator must possess the following characteristics :
- a) The colour change should be sharp
b) The colour change should be clear
c) It must be sensitive to the equivalent point
d) All of the above
349. The oxidation number of Xe in XeF_4 and XeO_2 is
- a) +6 b) +4 c) +1 d) +3
350. The oxidation number of arsenic in arsenate is :
- a) +5 b) +4 c) +6 d) +2
351. The reaction,
 $\text{Ag}^{+2}(\text{aq}) + \text{Ag}(\text{s}) \rightleftharpoons 2\text{Ag}^+(\text{aq})$
is an example of
- a) Reduction b) Oxidation c) Disproportionation d) None of these
352. During the presence of SO_3^{2-} and S^{2-} in a mixture, on addition of dil. H_2SO_4 , one notice that:
- a) SO_2 and H_2S are not formed
b) SO_2 and H_2S formed during change undergoes a redox change forming colloidal sulphur and thus, no

- smell
- c) A smell of burning sulphur
d) A smell of rotten egg
353. Which is not an oxidising agent?
a) KClO_3 b) O_2 c) $\text{C}_6\text{H}_{12}\text{O}_6$ d) $\text{K}_2\text{Cr}_2\text{O}_7$
354. The charge on cobalt in $[\text{Co}(\text{CN})_6]^{3-}$ is :
a) -6 b) $+3$ c) -3 d) $+6$
355. The most stable oxidation state of chromium is :
a) $+5$ b) $+3$ c) $+2$ d) $+4$
356. Arrange the following as increase in oxidation number
(i) Mn^{2+} (ii) MnO_2
(iii) KMnO_4 (iv) K_2MnO_4
a) (i) > (ii) > (iii) > (iv) b) (i) < (ii) < (iv) < (iii) c) (ii) < (iii) < (i) < (iv) d) (iii) > (i) > (iv) > (ii)
357. What mass of MnO_2 is reduced by 35 mL of 0.16 N oxalic acid in acidic solution? The skeleton equation is,
 $\text{MnO}_2 + \text{H}^+ + \text{H}_2\text{C}_2\text{O}_4 \rightarrow \text{CO}_2 + \text{H}_2\text{O} + \text{Mn}^{2+}$:
a) 8.7 g b) 0.24 g c) 0.84 g d) 43.5 g
358. Stronger is oxidising agent, more is;
a) Standard reduction potential of that species
b) The tendency to get itself oxidised
c) The tendency to lose electrons by that species
d) Standard oxidation potential of that species
359. How many g of KMnO_4 are needed to prepare 3.75 litre of 0.850 N solution if KMnO_4 is reduced as,
 $\text{MnO}_4^- + 8\text{H}^+ + 5\text{e} \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$?
a) 101 g b) 202 g c) 50.5 g d) 303.0 g KMnO_4
360. When KMnO_4 is reduced with oxalic acid in acid medium, the oxidation number of Mn changes from:
a) $+7$ to $+4$ b) $+6$ to $+4$ c) $+7$ to $+2$ d) $+4$ to $+2$
361. Addition of zinc powder to CuSO_4 solution precipitates copper due to :
a) Reduction of Cu^{2+} b) Reduction of SO_4^{2-} c) Reduction of Zn d) Hydrolysis of CuSO_4
362. Titrations in which liberated I_2 is estimated to carry out the volumetric estimations are known astitrations.
a) Iodometric b) Iodimetric c) Acidimetric d) Alkalimetric
363. In the course of chemical reaction, an oxidant :
a) Loses electron b) Gains electron c) Either of these d) None of these
364. In alkaline condition KMnO_4 reacts as follows :
 $2\text{KMnO}_4 + 2\text{KOH} \rightarrow 2\text{K}_2\text{MnO}_4 + \text{H}_2\text{O} + \text{O}$. The eq. wt. of KMnO_4 is :
a) 52.7 b) 158 c) 31.6 d) 79
365. Oxidation number of nitrogen in AgNO_3 is:
a) $+5$ b) -3 c) $+3$ d) -2
366. Total number of AlF_3 molecules in a sample of AlF_3 containing 3.01×10^{23} ions of F^- is :
a) 9×10^{24} b) 3×10^{24} c) 7×10^{23} d) 10^{23}
367. Oxidation number of N in NOCl is :
a) $+3$ b) $+2$ c) $+1$ d) $+4$
368. The oxidation state of chlorine is highest in the compound :
a) Cl_2 b) HCl c) Cl_2O d) Cl_2O_7
369. How many gram of KMnO_4 are contained in 4 litre of 0.05 N solution? The KMnO_4 is to be used as an oxidant in acidic medium :
a) 1.58 g b) 15.8 g c) 6.32 g d) 31.6 g
370. The reaction; $\text{H}_2\text{S} + \text{H}_2\text{O}_2 \rightarrow 2\text{H}_2\text{O} + \text{S}$ shows :
a) Acidic nature of H_2O_2
b) Alkaline nature of H_2O_2

- a) $A_2(BC)_2$ b) $A_2(BC_4)_3$ c) $A_3(BC_4)_2$ d) ABC
386. In an oxidation process for a cell $M_1 \rightarrow M_1^{n+} + ne$, the other metal (M_2) being univalent showing reduction takes up theelectrons to complete redox reaction.
- a) $(n - 1)$ b) 1 c) n d) 2
387. In which of the following reactions, chlorine acts as an oxidising agent?
- (i) $CH_3CH_2OH + Cl_2 \rightarrow CH_3CHO + HCl$
(ii) $CH_3CHO + Cl_2 \rightarrow CCl_3CHO + HCl$
(iii) $CH_4 + Cl_2 \rightarrow CH_3Cl + HCl$
- The correct answer is
- a) (i) only
b) (ii) only
c) (i) and (iii)
d) (i),(ii) and (iii)
388. During a redox change, the oxidant $K_2Cr_2O_7$ is always reduced to :
- a) Cr^{5+} b) Cr^{4+} c) Cr^{3+} d) Cr^{2+}
389. When potassium permanganate is titrated against ferrous ammonium sulphate, the equivalent weight of potassium permanganate is :
- a) Molecular weight/10 b) Molecular weight/5 c) Molecular weight/2 d) Molecular weight
390. Which conversion is an oxidation?
- a) $SO_4^{2-} \rightarrow SO_3^{2-}$ b) $Cu^{2+} \rightarrow Cu$ c) $H^+ \rightarrow H$ d) $H^- \rightarrow H$
391. In which case +1 oxidation state is stable than +3?
- a) Ga b) Al c) Tl d) B
392. In the reduction of dichromate by Fe(II), the number of electrons involved per chromium atom is :
- a) 3 b) 1 c) 2 d) 4
393. When $K_2Cr_2O_7$ is converted into K_2CrO_4 , the change in oxidation number of chromium is
- a) 0 b) 5 c) 7 d) 9
394. Which of the following acts as both an oxidizing as well as reducing agent?
- a) HNO_3 b) HNO_2 c) HI d) H_2SO_4
395. In which of the following compounds, nitrogen exhibits highest oxidation state?
- a) N_3H b) NH_2OH c) N_2H_4 d) NH_3
396. 1 mole of MnO_4^{2-} in neutral aqueous medium disproportionates to :
- a) $\frac{2}{3}$ mole of MnO_4^- and $\frac{1}{3}$ mole of MnO_2
b) $\frac{1}{3}$ mole of MnO_4^- and $\frac{2}{3}$ mole of MnO_2
c) $\frac{1}{3}$ mole of Mn_2O_7 and $\frac{1}{3}$ mole of MnO_2
d) $\frac{2}{3}$ mole of Mn_2O_7 and $\frac{1}{3}$ mole of MnO_2
397. Which one of the compound does not decolourised an acidified solution of $KMnO_4$?
- a) SO_2 b) $FeCl_3$ c) H_2O_2 d) $FeSO_4$
398. When one mole of $KMnO_4$ reacts with HCl, the volume of chlorine liberated at NTP will be:
- a) 11.2 litre b) 22.4 litre c) 44.8 litre d) 56.0 litre
399. What would happen when a small quantity of H_2O_2 is added to a solution of $FeSO_4$?
- a) Colour disappears
b) H_2 is evolved
c) An electron is added to Fe^{2+}
d) An electron is lost by Fe^{2+}
400. The oxidation state of I in IPO_4 is
- a) +1 b) +3 c) +5 d) +7
401. The number of moles of $KMnO_4$ reduced by one mole of KI in alkaline medium is
- a) 1 b) 5 c) $\frac{1}{2}$ d) $\frac{1}{5}$
402. A 0.50 M solution of KI reacts with excess of H_2SO_4 and KIO_3 solutions according to the equation, $6H^+ +$

- $5I^- + IO_3^- \rightarrow 3I_2 + 3H_2O$. Which of the following statements is true?
- 200 mL of the KI solution reacts with 0.10 mole KIO_3 .
 - 100 mL of the KI solution reacts with 0.060 M of H_2SO_4 .
 - 0.5 litre of the KI solution produces 0.15 mole of I_2
 - None of the above
403. Oxidation number of chromium in $K_2Cr_2O_7$ is :
- +2
 - +3
 - +6
 - 4
404. A standard solution is one whose :
- Concentration is 1 M
 - Concentration is unknown
 - Concentration is known
 - None of the above
405. In the reaction, $SO_2 + 2H_2S \rightarrow 3S + 2H_2O$, the substance oxidised is
- H_2S
 - SO_2
 - S
 - H_2O
406. Oxidation number of P in $HP_2O_7^-$ ion is
- +5
 - +6
 - +7
 - +3
407. The oxidation number that iron does not exhibit in its common compounds or in its elemental state is :
- Zero
 - +1
 - +2
 - +3
408. Oxidation number of Cl in $NOClO_4$ is :
- +7
 - 7
 - +5
 - 5
409. In which reaction is hydrogen acting as an oxidising agent?
- With iodine to give hydrogen iodide
 - With lithium to give lithium hydride
 - With nitrogen to give ammonia
 - With sulphur to give hydrogen sulphide
410. In presence of moisture SO_2 can :
- Gain electrons
 - Lose electrons
 - Act as oxidising agent
 - Does not act as reducing agent
411. The oxidation number of Mn in MnO_2 is :
- +4
 - +6
 - +2
 - 4
412. Which is not correct in case of Mohr's salt?
- It decolourises $KMnO_4$
 - It is primary standard
 - It is a double salt
 - Oxidation state of Fe is +3 in the salt
413. In the reduction of dichromate by Fe (II), the number of electrons involved per chromium atom is :
- 3
 - 1
 - 2
 - 4
414. Which of the following is a redox reaction?
- $NaCl + KNO_3 \rightarrow NaNO_3 + KCl$
 - $CaC_2O_4 + 2HCl \rightarrow CaCl_2 + H_2C_2O_4$
 - $Ca(OH)_2 + 2NH_4Cl \rightarrow CaCl_2 + 2NH_3 + 2H_2O$
 - $2K[Ag(CN)_2] + Zn \rightarrow 2Ag + K_2[Zn(CN)_4]$
415. What volume of 2N $K_2Cr_2O_7$ solution is required to oxidise 0.81 g of H_2S in acidic medium?
- 47.8 mL
 - 23.8 mL
 - 40 mL
 - 72 mL
416. Oxidation number of As atom in H_3AsO_4 is :
- +5
 - +6
 - +4
 - 3
417. In the following change, $3Fe + 4H_2O \rightarrow Fe_3O_4 + 4H_2$. If the atomic weight of iron is 56, then its equivalent weight will be :
- 42
 - 21
 - 63
 - 84
418. In permonosulphuric acid (H_2SO_5), the oxidation number of sulphur is
- +8
 - +4
 - +5
 - +6

419. The reaction,



is an example of

- a) Reduction b) Oxidation c) Comproportionation d) Disproportionation

420. Amount of oxalic acid present in a solution can be determined by its titration with KMnO_4 solution in the presence of H_2SO_4 . The titration gives unsatisfactory result when carried out in the presence of HCl , because HCl :

- a) Oxidises oxalic acid to carbon dioxide and water
b) Gets oxidized by oxalic acid to chlorine
c) Furnishes H^+ ions in addition to those from oxalic acid
d) Reduces permanganate to Mn^{2+}

421. Which is not a redox change?

- a) $\text{CaCO}_3 \rightarrow \text{CaO} + \text{CO}_2$
b) $2\text{H}_2 + \text{O}_2 \rightarrow 2\text{H}_2\text{O}$
c) $\text{Na} + \text{H}_2\text{O} \rightarrow \text{NaOH} + \frac{1}{2}\text{H}_2$
d) $\text{MnCl}_3 \rightarrow \text{MnCl}_2 + \frac{1}{2}\text{Cl}_2$

422. Sulphurous acid can be used as :

- a) Oxidising agent b) Reducing agent c) Bleaching agent d) All of these

SMART ACHIEVERS LEARNING PVT. LTD.

REDOX REACTIONS



CHEMISTRY

: ANSWER KEY :

1)	c	2)	c	3)	a	4)	d	177)	b	178)	d	179)	c	180)	a
5)	a	6)	d	7)	d	8)	b	181)	a	182)	c	183)	d	184)	a
9)	c	10)	d	11)	c	12)	d	185)	c	186)	a	187)	c	188)	d
13)	d	14)	b	15)	d	16)	b	189)	a	190)	c	191)	a	192)	b
17)	d	18)	a	19)	c	20)	a	193)	c	194)	a	195)	a	196)	b
21)	b	22)	a	23)	b	24)	c	197)	b	198)	b	199)	c	200)	d
25)	a	26)	a	27)	d	28)	a	201)	d	202)	c	203)	d	204)	a
29)	a	30)	d	31)	c	32)	a	205)	b	206)	b	207)	a	208)	c
33)	a	34)	a	35)	d	36)	c	209)	d	210)	a	211)	d	212)	c
37)	a	38)	a	39)	b	40)	a	213)	b	214)	c	215)	d	216)	c
41)	a	42)	d	43)	c	44)	a	217)	b	218)	d	219)	c	220)	d
45)	c	46)	c	47)	a	48)	b	221)	b	222)	c	223)	a	224)	a
49)	b	50)	b	51)	b	52)	c	225)	a	226)	b	227)	b	228)	b
53)	a	54)	a	55)	d	56)	c	229)	a	230)	b	231)	c	232)	b
57)	a	58)	a	59)	b	60)	c	233)	c	234)	c	235)	a	236)	a
61)	a	62)	a	63)	a	64)	c	237)	d	238)	c	239)	c	240)	c
65)	d	66)	c	67)	b	68)	b	241)	b	242)	a	243)	d	244)	b
69)	a	70)	b	71)	a	72)	b	245)	d	246)	a	247)	d	248)	a
73)	d	74)	a	75)	c	76)	b	249)	a	250)	c	251)	b	252)	d
77)	d	78)	b	79)	b	80)	d	253)	a	254)	a	255)	c	256)	a
81)	a	82)	d	83)	c	84)	a	257)	b	258)	c	259)	a	260)	c
85)	d	86)	b	87)	c	88)	a	261)	a	262)	d	263)	b	264)	c
89)	c	90)	b	91)	a	92)	b	265)	a	266)	b	267)	c	268)	b
93)	b	94)	a	95)	d	96)	c	269)	c	270)	c	271)	b	272)	c
97)	b	98)	c	99)	b	100)	b	273)	b	274)	b	275)	c	276)	c
101)	c	102)	b	103)	a	104)	a	277)	a	278)	a	279)	d	280)	a
105)	a	106)	b	107)	b	108)	a	281)	c	282)	a	283)	a	284)	c
109)	b	110)	a	111)	b	112)	d	285)	d	286)	c	287)	a	288)	a
113)	a	114)	a	115)	d	116)	d	289)	a	290)	d	291)	d	292)	b
117)	c	118)	b	119)	d	120)	b	293)	d	294)	c	295)	a	296)	d
121)	b	122)	a	123)	b	124)	a	297)	c	298)	a	299)	d	300)	a
125)	c	126)	a	127)	a	128)	a	301)	d	302)	b	303)	a	304)	c
129)	d	130)	a	131)	a	132)	b	305)	b	306)	d	307)	c	308)	a
133)	c	134)	b	135)	a	136)	d	309)	a	310)	c	311)	c	312)	a
137)	b	138)	a	139)	c	140)	a	313)	a	314)	d	315)	d	316)	c
141)	c	142)	b	143)	b	144)	b	317)	a	318)	c	319)	b	320)	a
145)	a	146)	c	147)	a	148)	d	321)	a	322)	c	323)	b	324)	a
149)	c	150)	c	151)	b	152)	b	325)	d	326)	b	327)	a	328)	c
153)	d	154)	a	155)	b	156)	a	329)	c	330)	a	331)	a	332)	a
157)	a	158)	a	159)	c	160)	d	333)	b	334)	b	335)	d	336)	a
161)	a	162)	d	163)	a	164)	a	337)	d	338)	a	339)	c	340)	d
165)	a	166)	a	167)	d	168)	a	341)	d	342)	d	343)	c	344)	a
169)	c	170)	b	171)	a	172)	c	345)	a	346)	b	347)	c	348)	d
173)	c	174)	a	175)	b	176)	c	349)	b	350)	a	351)	d	352)	b

353) c	354) b	355) b	356) b
357) b	358) a	359) a	360) c
361) a	362) a	363) b	364) b
365) a	366) d	367) a	368) d
369) c	370) c	371) a	372) c
373) d	374) b	375) a	376) b
377) a	378) b	379) d	380) b
381) a	382) a	383) a	384) d
385) c	386) c	387) d	388) c
389) b	390) d	391) c	392) a
393) a	394) b	395) a	396) a
397) b	398) d	399) d	400) b
401) a	402) c	403) c	404) c
405) a	406) b	407) b	408) a
409) b	410) b	411) a	412) d
413) a	414) d	415) b	416) a
417) b	418) d	419) c	420) d
421) a	422) d		

SMART ACHIEVERS LEARNING PVT. LTD.



Hence, changes in oxidation number are 5,1,3,4.


25 (a)

Alkaline earth metals have only +2 ox.no. in combined state.


26 (a)

Alkali metals are strongest reducing agents.


27 (d)



28 (a)



29 (a)





30 (d)

$$a + (-2) = 0$$

$$\therefore a = +2$$

31 (c)



32 (a)

Meq. of oxidant = Meq. of reductant

$$0.5 \times V = 2 \times 2000$$

$$\therefore V = 8 \text{ litre}$$

33 (a)



Oxygen shows -1 oxidation state in H_2O_2 .

$$2(+1) + 2x = 0$$

$$2x = -2$$

$$x = -1$$


34 (a)



On adding Eq. (i) and (ii), we get





To balance O atom, add $6\text{H}_2\text{O}$ molecules on RHS and 12H^+ on LHS, then



35 (d)

Cl has +7 ox.no. in KClO_4 .

36 (c)




37 (a)

Oxidation number in elemental form is zero. Covalency is two because of S-S-S-S-chain.

38 (a)



\therefore Total charge on cation or anion = +6



$$\therefore E = \frac{112}{6} \text{ or } \frac{56}{3}$$



39 (b)

C_3O_2 is carbon sub-oxide.

$$\text{Thus, } 3a - (2 \times 2) = 0$$

$$a = +\frac{4}{3}$$


40 (a)



41 (a)

Oxidation state of oxygen in H_2O_2 is -1. -1 is the intermediate oxidation state of oxygen.

42 (d)



S of H_2SO_4 is reduced.

43 (c)

$$1 + a + 3 \times (-2) = -1$$

$$\therefore a = +4$$


44 (a)

$$1 + a \times 2 = 0$$

$$\therefore a = -\frac{1}{2}$$



Ox.no. of alkali metals is always +1.

45 (c)






Only this reaction involves oxidation and reduction.

46 (c)




47 (a)




\therefore 3 mole of $\text{KMnO}_4 = 5$ mole of FeC_2O_4

48 (b)



$$E_{\text{IO}_4^-} = \frac{M}{7}$$

49 (b)



$$\therefore E_{\text{Fe}_3\text{O}_4} = \frac{M}{\text{No. of electrons lost or gained by one molecule}} = \frac{M}{1}$$

50 (b)

$$\text{Meq. of HNO}_3 = \text{Meq. of Fe}^{2+}$$

$$(\text{Eq. wt. of HNO}_3 = M/3)$$

$$\text{Or } 3 \times 3 \times V = \frac{8}{56} \times 1000$$

$$\therefore V = 15.87 \text{ mL}$$

51 (b)

The oxidation state of N are +5, +2, 0 and -3 in HNO₃, NO, N₂ and NH₄Cl respectively.

52 (c)

The oxidation state of iodine in HIO₄ is +7 as

$$1 + x + 4(-2) = 0$$

$$x = +7$$

The oxidation state of iodine in H₃IO₅ is +7 as

$$3 + x + 5(-2) = 0$$

$$x = +7$$

The oxidation state of iodine in H₅IO₆ is +7 as


$$5 + x + 6(-2) = 0$$

$$x = +7$$

53 (a)

Ag⁺ is reduced to Ag.

54 (a)





S of H₂SO₄ is reduced.

55 (d)

The characteristics of oxidant. Note these.

56 (c)



57 (a)

Meq. of bleaching powder = Meq. of Cl₂ = Meq. of hypo

$$\frac{w}{35.5} \times 1000 = 50 \times \frac{1}{10} \quad \therefore w_{\text{Cl}_2}$$

$$= 0.1775 \text{ g}$$




$$\therefore \text{Per cent Cl}_2 = \frac{0.1775}{5} \times 100 = 3.55 \%$$

58 (a)

$$a + 5 \times (-1) = 0$$




$$\therefore a = +5$$

59 (b)




$$\therefore E = M/3$$

60 (c)



61 (a)

Caro's acid is H₂SO₅. It has a peroxide linkage so, oxidation state of S is



Let the oxidation state of S is x .

H₂SO₅ (one peroxide bond)

$$+2 + x + 3(-2) + 1(-2) = 0$$

$$2 + x - 6 - 2 = 0$$

$$x - 6 = 0$$

$$x = 6$$

62 (a)

The formula for Eq. wt. of reductant or oxidant.

63 (a)





$$\therefore E = M/5$$

64 (c)

No doubt oxygen is taken in respiration, but oxidant-reduction occur simultaneously.

65 (d)



66 (c)

Both Os and Ru show +8 ox.no.


67 (b)

Two oxygen atom have peroxide linkage, (*i. e.*, -1 oxidation number) and six have -2 ox.no.

$$\text{Thus, } 2 \times 1 + 2 \times a + 6 \times (-2) + 2 \times (-1) = 0$$

$$\therefore a = +6$$

68 (b)



$$x + 2 = 5$$

$$\therefore x = 5 - 2 = +3$$

69 (a)

Meq. of K₂Cr₂O₇ = Meq. of FeSO₄

$$1 \times V = \frac{10}{152/1} \times 1000$$

- $\therefore V = 65.78 \text{ mL}$
- 70 **(b)**
 $\text{Cl}_2 + \text{H}_2\text{O} \rightarrow 2\text{HCl} + \text{O}$; thus, matter is oxidised by liberated oxygen.
- 72 **(b)**
 $\text{SnCl}_2 + 2\text{HgCl}_2 \rightarrow \text{Hg}_2\text{Cl}_2 + \text{SnCl}_4$
- 73 **(d)**
 Addition of KI to CuSO_4 makes it dark brown.
- 74 **(a)**
 Mn is stronger oxidising agent in +7 oxidation state. e.g., KMnO_4 .
- 75 **(c)**

$$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + n\text{Fe}^{2+} \rightarrow 2\text{Cr}^{3+} + n\text{Fe}^{3+} + 7\text{H}_2\text{O}$$

$$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6e^- \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$$
 (reduction)...(i)

$$\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + e^-$$
 (oxidation)...(ii)
 Eq.(ii) is multiplied by 6

$$6\text{Fe}^{2+} \rightarrow 6\text{Fe}^{3+} + 6e^-$$
 Thus, balanced equation is

$$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{Fe}^{2+} \rightarrow 2\text{Cr}^{3+} + 6\text{Fe}^{3+} + 7\text{H}_2\text{O}$$
 Hence, the value of 'n' is 6.
- 76 **(b)**
 $4e + \text{N}^{5+} \rightarrow \text{N}^+$
 \therefore Possible product is N_2O .
- 77 **(d)**
 Find oxidation number of P in each.
- 78 **(b)**
 $\text{CrO}_4^{2-} + \text{SO}_3^{2-} \rightarrow \text{Cr}(\text{OH})_4^- + \text{SO}_4^{2-}$
 Let the oxidation number of Cr is x in CrO_4^{2-}
 $x + 4(-2) = -2$
 $x = 6$
 and in $\text{Cr}(\text{OH})_4^-$ the oxidation number of Cr is y
 $y + 4(-2) + 4(1) = -1$
 $y - 8 + 4 = -1$
 $y = 3$
 Hence, oxidation number of Cr changes from +6 to +3.
- 79 **(b)**
 Find oxidation no. in each.
- 80 **(d)**
 $\text{Mn}^{7+} + e \rightarrow \text{Mn}^{6+}$
 $\therefore E = M/1$
- 81 **(a)**
 The sum of oxidation states of all elements in an ion is equal to charge on it.
 Let the oxidation state of S in $\text{SO}_4^{2-} = x$
- 90 **(b)**

- $\therefore x + (-2 \times 4) = -2$
 Or $x = +6$
- 82 **(d)**
 $\text{Sn}^{2+} \rightarrow \text{Sn}^{4+} + 2e$; $2e + \text{Hg}^{2+} \rightarrow \text{Hg}^0$
- 83 **(c)**
 Oxidation number of iodine in given species is as follows
 O.N. of iodine in $\text{IF}_3 = +3$
 O.N. of iodine in $\text{I}_3^- = -\frac{1}{3}$
 O.N. of iodine in $\text{IF}_5 = +5$
 O.N. of iodine in $\text{IF}_7 = +7$
- 84 **(a)**
 $1 + a + 3 \times (-2) = 0$
 $\therefore a = +5$
- 85 **(d)**
 KO_3 **Na_2O_2**
 Suppose O.N. of O = x suppose
 O.N. of O = x
 $+1 + 3x = 0$ $2 \times 1 + 2x = 0$
 $3x = -1$ $2 + 2x = 0$
 $x = -\frac{1}{3}$ $2x = -2$
 $x = -0.33$ $x = -\frac{2}{2}$
 $x = -1$
- 86 **(b)**
 $\text{I}_2^0 \rightarrow 2\text{I}^- + 2e$
- 87 **(c)**
 Os and Ru show +8 oxidation number.
- 88 **(a)**
 Meq. of $\text{AgNO}_3 = 100 \times 1 = 100$
 Meq. of $\text{CuSO}_4 = 100 \times 1 \times 2 = 200$
 Thus, H_2S is needed in the same Meq. ratio.
- 89 **(c)**
 $\text{Na}_2\text{S}_2\text{O}_3$,
 $2(+1) + 2x + 3(-2) = 0$
 $2 + 2x - 6 = 0$
 $x = +2$
 $\text{Na}_2\text{S}_4\text{O}_6$
 $2(+1) + 4(x) + 6(-2) = 0$
 $2 + 4x - 12 = 0$
 $4x = +10$
 $x = +2.5$

Meq. of $K^+ = \text{Meq. of } KMnO_4$

$$= \frac{1}{5} \times 1000 = 200$$

$$\therefore \text{Eq. of } K^+ = \frac{200}{1000} = 0.2$$

$$\text{Also, mole of } K^+ = \frac{0.2}{5} \left[\overset{\text{Valence factor} = 5}{Mn^{7+} + 5e \rightarrow Mn^{2+}} \right] = 0.04$$


$$\therefore \text{No. of } K^+ = \frac{0.2}{5} \times 6.023 \times 10^{23} = 2.4 \times 10^{22}$$

92 (b) $4e + Br^{5+} \rightarrow Br^{1+}$; Thus, BrO_3^- is to be reduced by a reducing agent.

93 (b) $6e + (N^0)_2 \rightarrow 2N^{-3}$
 $\therefore E_{N_2} = \frac{28}{6}; E_{NH_3} = \frac{17}{3}$


94 (a) F_2 is oxidant; ClO_4^- and MnO_4^- are also oxidant.

96 (c) None of elements in reaction (c) undergoes a change in oxidation number, therefore reaction (c) is not a redox reaction



It is a double decomposition reaction

97 (b) $Na_2S_4O_6$ is salt of $H_2S_4O_6$ which has the following structure



\Rightarrow Difference in oxidation number of two types of sulphur = 5

98 (c) Sum of oxidation no. of atoms in it is zero.

99 (b) $Sn^{2+} \rightarrow Sn^{4+} + 2e$
 $\therefore E = M/2 = \frac{119 + 71}{2} = 95$

100 (b) $2 \times 1 + a + 4 \times (-2) = 0$
 $\therefore a = +6$

101 (c) Electronic configuration of $Mn : 1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^5, 4s^2$ } More stable due to

$Mn^{2+} : 1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^5$ half filled d

$Mn^{7+} : 1s^2, 2s^2 2p^6, 3s^2 3p^6$

102 (b) Meq. of $H_2O_2 = 25 \times 0.5 \times 2 = 25$;
 Meq. of $KMnO_4 = 50 \times 0.2 \times 5 = 50$;
 $\therefore 25$ Meq. or 5 milli mole of $KMnO_4$ are left.

103 (a) $K - C \equiv N$
 N is more electronegative and thus, has -3 oxidation number as it involves three covalent bonds.
 Thus, $1 + a + (-3) = 0$
 $\therefore a = +2$

104 (a) Ox.no. of Ni is equal to zero.

105 (a) $Mn^{7+} + 1e \rightarrow Mn^{6+}$
 $\therefore E = M/1$

106 (b) Mn has $+6$ ox.no. in K_2MnO_4 and $+2$ ox.no. in $MnSO_4$.

107 (b) In reaction $0 \quad +1 \quad -1$
 $H_2O + Br_2 \rightarrow HOBr + HBr$

The oxidation number of bromine increases from 0 to $+1$ and decreases from 0 to -1 , so due to this reason bromine is both oxidised as well as reduced in the above reaction.

108 (a) $1 + 2 \times (+1) + a + 2 \times (-2) = 0$
 $\therefore a = +1$

109 (b) H in $LiAlH_4$ has -1 ox.no. and thus, easily oxidized.

110 (a) NO in iron complex has $+1$ ox.no.
 Thus, $a + 5 \times (0) + 1 + 1 \times (-2) = 0$
 $\therefore a = +1$

111 (b)



Let the oxidation state of Fe in $\text{Fe}_3\text{O}_4 = x$

$$\therefore 3x + 4 \times (-2) = 0$$


$$\text{Or } 3x - 8 = 0$$

$$\therefore x = \frac{8}{3}$$


112 (d)



113 (a)



114 (a)



$$E_{\text{NO}_3^-} = \frac{M}{8} = \frac{62}{8}$$

$$E_{\text{NH}_4^+} = \frac{M}{8} = \frac{18}{8}$$

115 (d)


$$1 + a + 4 \times (-2) = 0$$

$$\therefore a = +7$$

116 (d)

Find oxidation number of iodine in each.

117 (c)






118 (b)

$$3 \times 1 + a + 6 \times (-1) = 0$$


$$\therefore a = +3$$

119 (d)



Hence, number of moles of I_2 produced = 3

120 (b)



MnO_2 is itself reduced.

121 (b)

$$\text{Meq. of } \text{O}_2 = \text{Meq. of } \text{KMnO}_4 = 100 \times 0.5$$

$$\frac{w}{8} \times 1000 = 50$$

$$\therefore w_{\text{O}_2} = 0.4 \text{ g}$$

$$\therefore V_{\text{O}_2} = \frac{224 \times 0.4}{32} = 0.28 \text{ litre}$$

122 (a)



Oxidation involves loss of electrons and reduction involves gain of electrons, hence in case of oxidation-reduction reactions (redox reactions) charge remains conserved

123 (b)

$\text{Ni} \rightarrow \text{Ni}^{2+} + 2e$; Ni is oxidized and thus, reductant.

124 (a)


$$-1 \quad +6 \quad 0 \quad +4$$




oxidising

agent

125 (c)



126 (a)



The oxidation number of S increases from -2 to 0 in elemental sulphur and hence, H_2S gets oxidized

127 (a)

S_8 has zero oxidation state of S.

$$\text{In } \text{S}_2\text{F}_2: 2 \times a + 2 \times (-1) = 0; \therefore a = +1$$

$$\text{In } \text{H}_2\text{S}: 2 \times 1 + a = 0; \therefore a = -2$$

128 (a)

Cr in CrO_2Cl_2 has $+6$ and Mn in MnO_4^- has $+7$ oxidation number respectively, the highest value for them.

129 (d)

F is more electronegative than oxygen.

130 (a)

Oxidation number of Cl in ClO_3^- .


$$\text{ClO}_3 = -1$$

$$x + 3(-2) = -1$$

$$x = +6 - 1$$

$$x = +5$$

131 (a)



$$N = \frac{15}{74.5/2 \times 1} = 0.40$$

132 (b)

In ionic hydrides, H has -1 ox.no.

133 (c)

Let the oxidation number of Xe is x in XeOF_2 .

$$x + (-2) + 2(-1) = 0$$


$$x - 2 - 2 = 0$$

$$x = +4$$

134 (b)

No change in ox.no. of any species.

135 (a)



136 (d)

Both are same.

137 (b)

$$2 \times 1 + 2 \times 1 + 4 \times (-2) = 0$$

$$\therefore a = +3$$

138 (a)

Let oxidation state of P in $\text{Ba}(\text{H}_2\text{PO}_2)_2$ is x , then

$$2(+1) + 2[2(+1) + x + 2(-2)] = 0$$

$$2 + 2(2 + x - 4) = 0$$

- $\text{SO}_2 + 2\text{H}_2\text{S} \rightarrow 2\text{H}_2\text{O} + 3\text{S}$
- 160 (d)
 F^- can be oxidized to F_2 only by electrolysis.
- 161 (a)
 $2\text{S}_2^{2+} \rightarrow \text{S}_4^{(5/2)+} + 2e \therefore \text{Eq. wt. of Na}_2\text{S}_2\text{O}_3 = \frac{M}{1}$
 $\text{I}_2^0 + 2e \rightarrow 2\text{I}^-$
- 162 (d)
 The same species in each reaction is oxidized and reduced as well to give disproportionation reaction.
- 163 (a)
 N_3H (hydrazoic acid)
 $+3(x) + 1 = 0$
 $3x + 1 = 0$
 $x = -\frac{1}{3}$
- 164 (a)
 $\text{Fe}^{2+} + \text{Ce}^{4+} \rightarrow \text{Fe}^{3+} + \text{Ce}^{3+}$
 $5\text{Fe}^{2+} + \text{MnO}_4^- + 8\text{H}^+ \rightarrow 5\text{Fe}^{3+} + \text{Mn}^{2+} + 4\text{H}_2\text{O}$
 $\therefore \frac{\text{Moles of ceric ammonium sulphate}}{\text{Moles of potassium permanganate}} = \frac{1}{1/5}$
 $= 5.0$
- 165 (a)
 $\text{N}^{3-} \rightarrow \text{N}^{2+} + 5e$
 $\therefore E_{\text{NH}_3} = \frac{17}{5}$
- 166 (a)
 C has -4 ox.no. in CH_4 ,
 In rest all it has $+4$ ox.no.
- 167 (d)
 $[\text{Cr}_2^{6+} + 6e \rightarrow 2\text{Cr}^{3+}] \times 1; [\text{Sn}^{2+} \rightarrow \text{Sn}^{4+} + 2e]$
 $\times 3$
- 168 (a)
 Milliequivalent $[(W/\text{Eq. wt.}) \times 1000]$ or
 millimole $[(\frac{W}{M}) \times 1000]$ do not change on
 dilution.
- 169 (c)
 Cr^{3+} ion is green; $\text{Cr}_2^{6+} + 6e \rightarrow 2\text{Cr}^{3+}$.
- 170 (b)
 $\text{Hg} + \text{O}_3 \rightarrow \text{HgO} + \text{O}_2$
- 171 (a)
 $a + (4 \times 0) + 2 \times (-1) = 1$
 $\therefore a = +3$
- 172 (c)
 $2\text{V}^{2+} \rightarrow \text{V}_2^{5+} + 6e$
- 173 (c)
 $\text{Cr}^{6+} + 3e \rightarrow \text{Cr}^{3+}$
 $\therefore E = M/3$
- 174 (a)
 Ox.no. of Cr on both side is $+6$

- 175 (b)
 $\text{S}^{2-} \rightarrow \text{S}^{4+} + 6e$
 $\therefore \text{Eq.} = \text{mole} \times 6$
- 176 (c)
 $2 \times 1 + a + 4 \times (-2) = 0$
 $\therefore a = +6$
- 177 (b)
 Iodine has -1 (minimum ox.no.) and $+7$ (maximum ox.no.).
- 178 (d)
 These are formulae of Meq.
- 179 (c)
 $\text{CuSO}_4 + \text{H}_2\text{S} \rightarrow \text{CuS} + \text{H}_2\text{SO}_4$
- 180 (a)
 I in KIO_4 has $+7$ ox.no.
- 181 (a)
 $2 \times a + 3 \times (-2) = -2$
 $\therefore a = +2$
- 182 (c)
 $6e + \text{Cr}_2^{6+} \rightarrow 2\text{Cr}^{3+}$;
 $\text{Eq. wt. of Cr} = \frac{\text{at. wt.}}{3}$
- 183 (d)
 $\text{H}_2\text{O}_2 + \text{Na}_2\text{C O}_3 \rightarrow \text{Na}_2\text{O}_2 + \text{CO}_2 + \text{H}_2\text{O}$
 $\begin{matrix} -1 & +2 & +4 & -2 & & +1 & -1 & +4 & -2 & & -2 \\ \text{H}_2\text{O}_2 & + & \text{Na}_2\text{C O}_3 & \rightarrow & \text{Na}_2\text{O}_2 & + & \text{CO}_2 & + & \text{H}_2\text{O} \end{matrix}$
 None of the elements changes its oxidation number
- 184 (a)
 Usually burettes have least count of 0.1 mL.
- 185 (c)
 The oxidation state of N in NH_3 is
 $x + 3(+1) = 0$
 $x = -3$
 The oxidation state of N in HNO_3 is
 $1 + x + 3(-2) = 0$
 $x = 5$
 The oxidation state in N in NaN_3 is
 $+1 + 3x = 0$
 $x = -1/3$
 The oxidation state of N in Mg_3N_2 is
 $3(2) + 2x = 0$
 $6 + 2x = 0$
 $x = -3$
 Hence, three molecules
 (*i. e.*, NH_3 , NaN_3 , Mg_3N_2) have negative oxidation state.
- 186 (a)
 Fe in $\text{Fe}(\text{CO})_5$ has zero oxidation no., *i. e.*, the lowest for metals.
- 187 (c)
 The weight of rider used is 0.0002 g.

- 188 (d)
Ions are hydrated on dissolution of salt in water.
- 189 (a)
Ox.no. of each element on two sides is same.
- 190 (c)
 $10e + 2\text{Br}^{5+} \rightarrow \text{Br}_2^0 \quad \therefore \text{Eq. wt. of KBrO}_3 = \frac{M}{5}$
 $2\text{Br}^- \rightarrow \text{Br}_2 + 2e$
- 191 (a)
Corrosion involves oxidation of species.
- 192 (b)
 $\text{MnO}_4^- + 8\text{H}^+ + 5e^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$
 $[\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + e^-]^5$

 $\text{MnO}_4^- + 8\text{H}^+ + 5\text{Fe}^{2+} \rightarrow \text{Mn}^{2+} + 5\text{Fe}^{3+} + 4\text{H}_2\text{O}$
 \therefore Five electrons gets transferred.
- 193 (c)
 $1 + a + 3 \times (-2) = 0$
 $\therefore a = +5$
- 194 (a)
The species present in solution but does not take part in the reaction and are also omitted while writing the potential redox change are called spectator ion.
- 195 (a)
It is the formula of turns bull's blue.
- 196 (b)
Si has 4 electrons in its valence shell. When it reacts with strongly electropositive metal like Na, Mg, K etc., it gives 4 electrons and its oxidation state in this case is -4 .
- 197 (b)
Oxygen in H_2O_2 has ox.no. -1 which can increase or decrease.
- 198 (b)
 $A^{n-} \rightarrow A^{a+} + (a+n)e$
 $\text{Cr}_2^{6+} + 6e \rightarrow 2\text{Cr}^{3+}$
 Also, Meq of A = Meq. of $\text{K}_2\text{Cr}_2\text{O}_7$
 $3.26 \times 10^{-3}(a+n) = 1.68 \times 10^{-3} \times 6$
 Or $a+n = 3$
 $\therefore a = 3 - n$
- 199 (c)
 $\text{H}_2^0 \rightarrow \text{H}_2^+ + 2e$ (H_2O is formed)
- 200 (d)
 $2 \times a + 7 \times (-2) = 0$
 $\therefore a = +7$
- 201 (d)
Due to higher E_{OP}^0 order.
- 202 (c)
Cl atom is oxidised ($\text{Cl}^{1+} \rightarrow \text{Cl}^{5+} + 4e$) as well as Cl atom is reduced ($\text{Cl}^{1+} + 2e \rightarrow \text{Cl}^-$). Such

reactions are called auto redox or disproportionation reactions.

- 203 (d)
Ox.no. of S in $\text{Na}_2\text{S}_4\text{O}_6$ is no doubt 2.5 but it is average of two values, i. e.,
 $\frac{2 \times (+5) + 2 \times 0}{4} = +5/2$
- 204 (a)
De-electronation is loss of electrons, i.e. $M \rightarrow M^{4+} + 4e$
- 205 (b)
 $\text{CaCO}_3 \xrightarrow{\Delta} \text{CaO} + \text{CO}_2$; This is simple decomposition and not a redox change.
- 206 (b)
 S^{2-} has minimum ox.no. and thus, can act only as reducing agent.
- 207 (a)
It imparts its colour at end point.
- 208 (c)
 $\text{Zn}^0 \rightarrow \text{Zn}^{2+} + 2e$
- 209 (d)
Oxygen has highest electron affinity in its family.
- 210 (a)
 $\text{Na}_2[\text{Fe}(\text{CN})_5\text{NO}]$
- 211 (d)
The formula is obtained by taking an account of g atoms.
 $\text{Xe} = \frac{53.3}{131} = 0.4, \text{F} = \frac{46.5}{20} = 2.325,$
 i. e., 1 : 6 or XeF_6
- 212 (c)
N in $\text{NH}_3, \text{NH}_4^+, \text{N}_3\text{H}$ and NO_2^- has $-3, -3, -1/3$ and $+3$ oxidation number respectively.
- 213 (b)
Meq. of $\text{H}_2\text{O}_2 = \text{Meq. of KMnO}_4$
 $\frac{w}{34/2} \times 1000 = 10 \times 1$
 $\therefore w_{\text{H}_2\text{O}_2} = 0.17$
 $\therefore \text{Per cent purity} = \frac{0.17}{0.2} \times 100 = 85\%$
- 214 (c)
 $\text{Mn}^{7+} + e \rightarrow \text{Mn}^{6+} \quad (\text{MnO}_4^{2-})$
 $\text{Mn}^{7+} + 3e \rightarrow \text{Mn}^{4+} \quad (\text{MnO}_2)$
 $2\text{Mn}^{7+} + 8e \rightarrow (\text{Mn}^{3+})_2 \quad (\text{Mn}_2\text{O}_3)$
 $\text{Mn}^{7+} + 5e \rightarrow \text{Mn}^{2+} \quad (\text{MnO}_2)$
- 215 (d)
The reaction involves :
 $\text{H}_2\text{O}_2 + 2\text{I}^- + 2\text{H}^+ \rightarrow \text{I}_2 + 2\text{H}_2\text{O}(l)$
 $2\text{Na}_2\text{S}_2\text{O}_3 + \text{I}_2 \rightarrow \text{Na}_2\text{S}_4\text{O}_6 + 2\text{NaI}$
 The reaction gives blue colour only after all the

$\text{Na}_2\text{S}_2\text{O}_3$ is used. The reaction is carried out with adjusted amount of $\text{Na}_2\text{S}_2\text{O}_3$ so that only a fraction of H_2O_2 and KI reaction occurs before the blue colour of starch— I_2 appears, however the slow redox reaction of H_2O_2 — I_2 continues. The appearance of blue colour is like clock alarm and in such reactions time for the appearance of blue colour is noticed. The phenomenon is used in studying rate of reaction. If time taken for blue colour appearance is longer, the reaction is slow and *vice – versa*.

216 (c)

N in $(\text{N}_2\text{H}_5)_2\text{SO}_4$ has -2 ox.no.

217 (b)


The $5p$ –electrons of outermost shell in iodine are unpaired during their excitation to $5d$ –subshell.

218 (d)

A characteristic property of transition elements.

219 (c)

Let the oxidation state of sulphur in $\text{Na}_2\text{S}_4\text{O}_6$ is x .



$$1 \times 2 + 4 \times x + (-2) \times 6 = 0$$

$$2 + 4x - 12 = 0$$

$$4x - 10 = 0$$

$$4x = 10$$

$$x = \frac{10}{4} = 2.5$$

220 (d)

F_2 is strongest oxidant among all the species.

221 (b)

S has $+6$ ox. no. in SO_3

222 (c)



$$3 \times a + 1 \times 1 = 0$$

$$\therefore a = -1/3$$

223 (a)

Tendency to lose more electron for cation decreases.

224 (a)



\therefore 1 mole of NO_3^- (Or NaNO_3) is reduced by

= 10 moles of HCl

$\therefore \frac{1}{2}$ mole of NO_3^- will be reduced by

= $10 \times \frac{1}{2}$ moles of HCl

= 5 moles of HCl


225 (a)

Meq. of FeSO_4 = Meq. of KMnO_4

$$\frac{w}{152/1} \times 1000 = 200 \times 1$$


$$\therefore w = 30.4 \text{ g}$$

226 (b)



$$x = 2$$

227 (b)



228 (b)

Meq. of $\text{Na}_2\text{S}_2\text{O}_3$ = Meq. of CuSO_4


$$\therefore V \times 0.4 \times 1 = 50 \times 0.2 \times 1$$

$$\therefore V = 25 \text{ mL}$$



229 (a)

$$N = \frac{47.5}{189.7/2 \times 2.25} = 0.222 \text{ N}$$

230 (b)



231 (c)



232 (b)

Oxidation no. of N in NO^+ is

$$(1 \times x) + 1 \times (-2) = +1$$

$$\therefore x = +3$$

Oxidation no. of Cl in ClO_4^- is

$$(1 \times x) + 4 \times (-2) = -1$$

$$x = +7$$

233 (c)

1. Sulphurous acid H_2SO_3

$$2 + x + (-2 \times 3) = 0$$

$$x - 4 = 0$$

$$\therefore x = 4$$

2. Pyrosulphuric acid ($\text{H}_2\text{S}_2\text{O}_7$)

$$2 + 2x + (-2 \times 7) = 0$$

$$\text{or } 2x = 12$$

$$\therefore x = 6$$

3. Thiosulphuric acid ($\text{H}_2\text{S}_2\text{O}_3$)

$$2 + 2x + (-2 \times 3) = 0$$

$$\text{or } 2x = 4$$

$$x = 2$$


4. Dithionous acid ($\text{H}_2\text{S}_2\text{O}_4$)

$$2 + 2x + (-2 \times 4) = 0$$

$$2x = 6$$


$$\therefore x = 3$$

234 (c)



CN^- also acts as reducing agent.

235 (a)



236 (a)


Meq. of oxalic acid = Meq. of KMnO_4

$$V \times 0.1 \frac{250 \times 8}{100 \times 31.6} \times 1000 = 6.3 \text{ litre}$$

237 (d)

H_3PO_3 is phosphorous acid.

238 (c)



239 (c)

$$\text{H}_4\text{P}_2\text{O}_5 : 4 \times 1 + 2 \times a - 5 \times 2 = 0$$

$$a = +3$$


$$\text{H}_4\text{P}_2\text{O}_6 : 4 \times 1 + 2 \times a - 6 \times 2 = 0$$

$$a = +4$$

$$\text{H}_4\text{P}_2\text{O}_7 : 4 \times 1 + 2 \times a - 7 \times 2 = 0$$

$$a = +5$$

240 (c)



241 (b)

Meq. of oxalate = Meq. of KMnO_4

$$\frac{w}{88/2} \times 1000 = 90 \times \frac{1}{20}$$

$$\therefore w_{\text{oxalate ion}} = 0.198 \text{ g}$$

$$\therefore \% \text{ of oxalate ion} = \frac{0.198}{0.3} \times 100 = 66\%$$

242 (a)

Meq. of Cl_2 = Meq. of KMnO_4

$$\frac{w}{71/2} \times 1000 = \frac{10}{31.6} \times 1000$$


$$\therefore w_{\text{Cl}_2} = 11.23 \text{ g}$$

$$\therefore V_{\text{Cl}_2} = \frac{22.4 \times 11.23}{71} = 3.54 \text{ litre}$$



243 (d)

$$N = \frac{15.8 \times 1000}{158/5 \times 100} = 5$$

244 (b)




245 (d)



246 (a)

Meq. of KMnO_4 = Meq. of FeC_2O_4



$$0.1 \times 5 \times V = \frac{100 \times 10^{-3}}{144/3} \times 1000$$

$$\therefore V = 4.1 \text{ mL}$$

247 (d)

It is precipitation reaction.

248 (a)

Meq. of lime stone = Meq. of CaC_2O_4
= Meq. of KMnO_4

= Meq. Of CaO

$$\therefore 40 \times 0.250 = \frac{w}{56/2} \times 1000$$

$$\therefore w_{\text{CaO}} = 0.28$$

$$\therefore \text{per cent of CaO} = \frac{0.28 \times 100}{0.518} = 54\%$$


249 (a)

Equate charge on both side, $2 \times 3 + 2 = 2 \times 2 + a$

$\therefore a = +4$; Thus, Sn^{4+} is choice.

250 (c)

Br_2 is disproportionated in basic medium as



251 (b)

Carbon has negative oxidation no. in Mg_3C_2 and positive oxidation number in C_3O_2 ; O is more electronegative than C. Mg is more electropositive than C.

252 (d)

It is a complexation reaction involving reduction of I_2 and oxidation of KI.

253 (a)

Oxidation state of Cr in Cr_2O_3 is



$$2x + (-2)3 = 0$$

$$2x - 6 = 0$$

$$2x = 6$$

$$x = +3$$

254 (a)



$$2 \times a + 2 \times (-1) = 0$$

$$\therefore a = +1$$

255 (c)





N has + 1 ox.no.


256 (a)



5 mole FeSO_4 = 1 mole KMnO_4


$$'X' = \frac{2}{3} \text{ mole}$$





$$\therefore 'Y' = \frac{3 \times 2}{5}$$

257 (b)




258 (c)

Meq. of salt = Meq. Of Na_2SO_3

$$50 \times 0.1 \times n = 25 \times 0.1 \times 2$$

$$\therefore n = 1 \text{ (change in ox. no.)}$$




259 (a)

Cu^{2+} is more stable than Cu^+ although later, has $3d^{10}$ configuration. In Cu^+ 18 electron core is not held properly by nuclear charge and thus, Cu^+ is readily converted to Cu^{2+} .


260 (c)

\therefore In this reaction phosphorus is simultaneously oxidised and reduced.

\therefore It is disproportionation reaction.



261 (a)



262 (d)

All terms have same meaning.

263 (b)

The sum of the oxidation states is always zero in neutral compound.

The oxidation state of X, Y, and Z are +2, +5 and -2 respectively.

5. In X_2YZ_6

$$2 \times 2 + 5 + 6(-2) \neq 0$$

6. In XY_2Z_6

$$2 + 5 \times 2 + 6(-2) = 0$$

7. In XY_5

$$2 + 5 \times 5 \neq 0$$

8. In X_3YZ_4

$$3 \times 2 + 5 + 4(-2) \neq 0$$

Hence, the formula of the compound is XY_2Z_6 .




264 (c)

F is most electronegative element and thus, has -1 ox.no.

$$\text{Thus, } a + (-2) = 0$$

$$\therefore a = +2$$

265 (a)



Meq. of Fe^{2+} = Meq. of KMnO_4 = Meq. of $\text{K}_2\text{Cr}_2\text{O}_7$

$$1 \times 5 \times V_{\text{KMnO}_4} = 1 \times 6 \times V_{\text{K}_2\text{Cr}_2\text{O}_7}$$

$$\therefore V_{\text{KMnO}_4} = \frac{6}{5} V_{\text{K}_2\text{Cr}_2\text{O}_7}$$

266 (b)

Meq. of KMnO_4 in 1 mL = Meq. of Fe = $\frac{5 \times 10^{-3}}{56/1} \times 10^3$

$$\therefore \text{Meq. if } \text{KMnO}_4 \text{ in 250 mL} = \frac{5 \times 250}{56/1}$$

$$\text{Thus, } \frac{w}{31.6} \times 1000 = \frac{5 \times 250}{56/1} = 0.7 \text{ g}$$

267 (c)


Let the oxidation number of Cr in K_2CrO_4 is x.

$$2(+1) + x + 4(-2) = 0$$


$$2 + x - 8 = 0$$

$$x = +6$$

268 (b)



269 (c)



$$\therefore E = M/5$$

270 (c)

Let the oxidation number of Cr be x

\therefore For $\text{K}_2\text{Cr}_2\text{O}_7$


$$+1 \times 2 + 2x + 7(-2) = 0$$

$$2 + 2x - 14 = 0$$

$$2x = 12$$

$$x = 6$$


271 (b)



272 (c)

$\text{S}^{4+} + 4e \rightarrow \text{S}^0$; SO_2 is reduced and thus, oxidant.

273 (b)



274 (b)

Let the oxidation number of carbonyl carbon in methanal (HCHO) and methanoic acid (HCOOH) is x and y is respectively.

In HCHO ,

$$2(+1) + x + (-2) = 0$$

$$2 + x - 2 = 0$$

$$x = 0$$


In HCOOH ,

$$2(+1) + y + 2(-2) = 0$$

$$2 + y - 4 = 0$$


$$y = 2$$

275 (c)




$$\therefore E = \frac{M}{10} = \frac{254}{10} = 25.4$$

276 (c)




$$\therefore E_{SO_2} = \frac{64}{4} = 16$$

277 (a)





278 (a)



Oxidation number of hydrogen is decreasing from 0 to -1. So, H_2 is acting as oxidising agent in this reaction.


279 (d)

Mohr's salt is $FeSO_4 \cdot (NH_4)_2 SO_4 \cdot 6H_2O$



280 (a)

1 faraday of electricity involves change of one mole electron.



281 (c)

Oxidation of Co and reduction of Cu^{2+} is taking place.

282 (a)

$$4 \times 1 + a + 6 \times (-2) = -1$$


$$\therefore a = +7$$

283 (a)

$$a + 3 \times (+1) = 0$$

$$\therefore a = -3$$

284 (c)



285 (d)


$$4 \times 1 + a + 4 \times (-1) = 0$$

$$\therefore a = 0$$

286 (c)

Ox. no. of each species remains same.

287 (a)




288 (a)


$$4 \times 1 + a + 6 \times (-1) = 0$$

$$\therefore a = +2$$

289 (a)




290 (d)



H_2S - Oxidation, Reducing agent.

H_2O_2 - Reduction, Oxidising agent.

291 (d)




292 (b)

$$a + 2 \times 1 - 1 = 0$$


$$\therefore a = -1$$

293 (d)



$$\therefore E = \frac{M}{1}$$

294 (c)




$Cr_2O_7^{2-}$ is reduced.

295 (a)




296 (d)




297 (c)

The reactions, in which the same element is oxidised as well as reduced, are called disproportionation reactions.



In this reaction, the same element, *ie.*, Cl_2 is oxidised as well as reduced, so it is an example of disproportionation reaction.

298 (a)




$Cr_2O_7^{2-}$ is reduced to Cr^{3+} .

Thus, final state of Cr is +3. Hence, (a)


299 (d)

$NaNO_2$ (Sodium nitrite) acts both as oxidising as well as reducing agent because in it N-atom is in +3 oxidation state (intermediate oxidation state).

Oxidising property



Reducing property



300 (a)


Graphic is uncombined state of carbon.

301 (d)

$$6 \times a + 12 \times 1 + 6 \times (-2) = 0$$

$$\therefore a = 0$$

302 (b)



303 (a)

$$2 \times 2 + 2 \times a + 7 \times (-2) = 0$$

$$\therefore a = +5$$

304 (c)


Eq. of Cl_2 = eq. of chloride

$$1 \times 2 = \frac{111}{E + 35.5}$$

$$\therefore E = 40$$


$$\therefore M = 40 \times 2 = 80 \text{ (Metal is bivalent.)}$$

305 (b)



It is chromium peroxide.

Let the oxidation number of Cr is "x".



$$x + (-1)2 + (-1)2 + (-2)1 = 0$$

$$x - 6 = 0$$

$$x = +6$$


Hence, the oxidation state of Cr is +6.

306 (d)

Haematite is Fe_2O_3 , in which oxidation number of iron is III.



Magnetite is Fe_3O_4 which is infact a mixed oxide ($\text{FeO} \cdot \text{Fe}_2\text{O}_3$), hence iron is present in both II and III oxidation state.

307 (c)



308 (a)

In basic medium



Change in oxidation number

$$= 7 - 6 = +1$$

So, electrons involved = $1e^-$

309 (a)

In NH_4^+ , N has ox.no. -3 and in NO_3^- , N has ox.no. +5.

310 (c)

$$a + 6 \times (-1) = -2$$

$$\therefore a = +4$$

311 (c)


$$1 + 1 \times (-2) + a = 0$$

$$\therefore a = +1$$

312 (a)

$e + \text{N}^{5+} \rightarrow \text{N}^{4+}$; Thus, HNO_3 is oxidant.



313 (a)



314 (d)

$\text{S} \xrightarrow{\text{O}_2} \text{SO}_2 \xrightarrow{\text{Cl}_2} \text{SO}_4^{2-} \xrightarrow{\text{BaCl}_2} \text{BaSO}_4$ One mole of S will give one mole of BaSO_4 . Thus, mole of BaSO_4 formed = mole of S = $\frac{8}{32} = \frac{1}{4}$

315 (d)



316 (c)

Equal equivalent of species react together.


317 (a)

It is a fact.

318 (c)

The balanced disproportionation reaction involving white phosphorus with aq. NaOH is

Oxidation of P^0 to P^{+1} state





Reduction of P^0 to P^{-3} state

319 (b)

F can have only -ve ox.no., i.e., $2e + \text{F}_2^0 \rightarrow 2\text{F}^{1-}$ or F_2 can be reduced only.

320 (a)




$$E_{\text{N}_2} = \frac{28}{6}; E_{\text{NH}_3} = \frac{17}{3}$$

321 (a)

$\text{SO}_2 + 2\text{H}_2\text{O} \rightarrow \text{H}_2\text{SO}_4 + 2\text{H}$; thus, matter is reduced by liberated hydrogen.

322 (c)

N_2 undergoes oxidation and reduction as well;



323 (b)






324 (a)



325 (d)

All these substances can accept electrons and can decrease their oxidation number and hence, all these act as oxidation agent





326 (b)

$$\text{Meq. of I}_2 = \text{Meq. of Na}_2\text{S}_2\text{O}_3 = 40 \times 0.11$$

$$\therefore \frac{w}{254/2} \times 1000 = 40 \times 0.11$$

$$w_{\text{I}_2} = 0.558 \text{ g}$$

327 (a)



328 (c)


$$\text{Meq. of HNO}_3 = 1000 \times 2 = 2000$$

$$\therefore \frac{w}{63/3} \times 1000 = 2000$$

$$\therefore w = 42 \text{ g}$$

329 (c)

The chemical structure of $\text{H}_2\text{S}_2\text{O}_8$ is as follows



$$2 \times (+1) + 2 \times x + 6 \times (-2) + 2 \times (-1) = 0$$

for H for S for O for O

$$+2 + 2x - 12 - 2 = 0$$


$$2x = +12$$

$$x = +6$$

330 (a)



331 (a)



$$\therefore E = M/2 = \frac{34}{2} = 17$$

332 (a)



333 (b)

In N_3H

$$\text{Oxidation number of N} = -\frac{1}{3}$$

In N_2O_4 Oxidation number of N = +4

In NH_2OH Oxidation number of N = -1



In NH_3 Oxidation number of N = -3

Hence, in N_2O_4 the oxidation number of nitrogen is highest.


334 (b)

Starch + $\text{I}_2 \rightarrow$ Blue

335 (d)



336 (a)



337 (d)

$$3 \times a + (+1) = 0$$

$$\therefore a = -1/3$$

338 (a)

$$\text{Mole of O}_2 \text{ formed} = \frac{3}{24} = \frac{1}{8}$$

$$\therefore \text{Mole of H}_2\text{O}_2 = \frac{1}{8} \times 2 = \frac{1}{4}$$

$$\therefore 100 \times X = \frac{1}{4} \times 1000 \text{ (m mole} = M \times V)$$

$$\therefore X = 2.5$$

339 (c)



340 (d)

+2 and +3

341 (d)

It is a fact.

342 (d)



Thus, 27 g Al forms Al^{3+} by losing 3N electrons

$$\therefore 13.5 \text{ g Al will lose } \frac{3N \times 13.5}{27} = \frac{3}{2} N \text{ electrons}$$

343 (c)

$$a + 2 \times 1 + 2 \times (-1) = 0$$

$$\therefore a = 0$$

344 (a)

Mn has +7 oxidation state in KMnO_4 .

$$1 + x + 4(-2) = 0$$

$$1 + x - 8 = 0$$

$$x = +7$$

345 (a)

Minimum ox.no. = group no. - 8.

Maximum ox.no. = group no.

346 (b)

H possesses negative one value of oxidation number in ionic hydrides.

347 (c)

Due to -ve oxidation number it should be non-metal having six electrons in outer shell.

348 (d)

These are characteristics of indicator.

349 (b)

The oxidation state of Xe in both XeO_2 and XeF_4 is 4.

$$\begin{array}{ccc} \text{XeO}_2 & & \text{XeF}_4 \\ x + 2(-2) = 0 & & x + 4(-1) = 0 \\ x = 4 & & x = 4 \end{array}$$

350 (a)


Na_3AsO_4 is sodium arsenate

Or AsO_4^{3-} is arsenate.

Thus, $a + 4 \times (-2) = -3$


$\therefore a = +5$

351 (d)



Hence, those reactions in which two or more species undergo oxidation as well as reduction are called comproportionation.

352 (b)



353 (c)

Glucose is reducing agent.

354 (b)

$a + 6 \times (-1) = -3$

$\therefore a = +3$

355 (b)

It is a fact.

356 (b)

i. Oxidation state of Mn in $\text{Mn}^{2+} = +2$

ii. Let oxidation state of Mn in $\text{MnO}_2 = x$

$\therefore x + (2 \times -2) = 0$

$\therefore x = +4$

(iii) Let the oxidation state of Mn in $\text{KMnO}_4 = x$

$\therefore +1 + x + (-2 \times 4) = 0$

$\therefore x = +7$

iv) Let oxidation state of Mn in $\text{K}_2\text{MnO}_4 = x$

$\therefore (+1 \times 2) + x + (-2 \times 4) = 0$

$\therefore x = +6$

\therefore Increasing order of oxidation states is

(i) < (ii) < (iv) < (iii)

357 (b)

Meq. of $\text{MnO}_2 = \text{Meq. of oxalic acid}$

$= 0.16 \times 35 = 56$

$\therefore \frac{w}{87/2} \times 1000 = 56$

$w_{\text{MnO}_2} = 0.24 \text{ g}$

358 (a)

More is E_{RP}^0 , more is the tendency to get itself reduced or more is oxidising power.


359 (a)

Meq. of $\text{KMnO}_4 = 3750 \times 0.85$


$\therefore \frac{w}{31.6} \times 1000 = 3750 \times 0.85$

$\therefore w = 100.7 \text{ g}$

360 (c)




361 (a)




362 (a)

It is definition of iodimetric titrations.

363 (b)



364 (b)



$\therefore E = M/1$

365 (a)

$1 + a + 3 \times (-2) = 0$

$\therefore a = +5$

366 (d)

\therefore 3 ions of F^- from 1 molecule of AlF_3

$\therefore 3 \times 10^{23}$ ions of F^- from 10^{23} molecules of AlF_3

367 (a)

Calculate ox.no. by taking NO^+ in NOCl

368 (d)

Cl has +7 ox.no. in Cl_2O_7 .

369 (c)

Meq. of $\text{KMnO}_4 = 4000 \times 0.05$

$\therefore \frac{w}{31.6} \times 1000 = 4000 \times 0.05$


$w = 6.32 \text{ g}$

370 (c)

H_2O_2 oxidises S^{2-} to S^0 .

371 (a)

Following is balanced redox reaction.



So, coefficients of

MnO_4^- , $\text{C}_2\text{O}_4^{2-}$ and H^+ are 2, 5, and 16 respectively.

372 (c)


$2 \times a + 1 \times (-2) = 0$

$\therefore a = +1$

373 (d)


Oxidation-reduction takes place simultaneously.

374 (b)



$$\therefore \text{Eq. wt.} = \frac{\text{mol. wt.}}{6}$$

375 (a)




376 (b)

F_2 shows only -1 ox.no.

377 (a)

Reduction (oxidation number decreases)



Oxidation (oxidation number is increases)


The reactions in which the same substance undergoes oxidation as well as reduction, are called disproportionation reactions.

So, the above reaction is an example of disproportionation reaction.

378 (b)

It is definition of iodimetric titrations.

379 (d)



380 (b)

$+2$ oxidation state due to $1s^2, 2s^2, 2p^2$ configuration having 2 unpaired electrons in $2p$ -subshell. $+4$ oxidation state due to $1s^2, 2s^1 2p^3$ configuration in excited state having four unpaired electrons.

381 (a)

Meq. of $\text{SnCl}_2 = \text{Meq. of HgCl}_2$

$$0.5 \times V = 600 \times 0.1$$

$$\therefore V = 120 \text{ mL}$$

382 (a)

Meq. of $\text{FeSO}_4 = \text{Meq. of KMnO}_4 = 200 \times 1$

$$\therefore \frac{w}{152/1} \times 1000 = 200$$

$$\therefore w = 30.4 \text{ g}$$





383 (a)

Meq. of $\text{Fe} = \text{Meq. of K}_2\text{Cr}_2\text{O}_7$

$$\frac{w}{56/1} \times 1000 = 1 \times 0.1055$$

$$\therefore w = 5.9 \times 10^{-3} \text{ g} = 5.9 \text{ mg}$$

384 (d)



$$\therefore 3 \text{ mole MnO}_4^- \equiv 5 \text{ mole FeC}_2\text{O}_4$$


385 (c)

The sum of oxidation number is zero.



386 (c)

Electrons released at anode = Electrons used at cathode.

388 (c)



389 (b)



390 (d)



Loss of an electron or increase in oxidation number is oxidation process.



391 (c)

Due to inert pair effect which is more predominant in T1.

392 (a)



Thus, electrons involved per Cr atom = 3.

393 (a)

Let oxidation state of Cr in $\text{K}_2\text{Cr}_2\text{O}_7 = x$

$$(+1 \times 2) + 2x + (-2 \times 7) = 0$$

$$\text{or } +2 + 2x - 14 = 0$$

$$\therefore x = +6$$


Let oxidation state of Cr in $\text{K}_2\text{CrO}_4 = x$

$$+1 \times 2 + x + (-2 \times 4) = 0$$

$$2 + x - 8 = 0$$

$$x = 6$$

\therefore Change in oxidation state of Cr is zero when it changes from






394 (b)

In HNO_2 , the oxidation number of N is $+3$ which is less than the maximum possible, oxidation number *ie*, $+5$ and more than the minimum possible oxidation number *ie*, -3 , therefore, it can act both as an oxidizing as well as reducing agent

395 (a)

Ox. No. of N in $\text{N}_3\text{H}, \text{NH}_2\text{OH}, \text{N}_2\text{H}_4, \text{NH}_3$ are $-\frac{1}{3}, -1, -2, -3$ respectively.

396 (a)



397 (b)

FeCl_3 cannot be oxidised because Fe has highest oxidation state.

398 (d)


Meq. of $\text{KMnO}_4 = \text{Meq. Of Cl}_2$

$$1 \times 5 \times 1000 = \frac{w}{(71/2)} \times 1000$$

$$\therefore w = 177.5 \text{ g}$$

$$\therefore V_{Cl_2} = 56 \text{ litre at NTP}$$

399 (d)



400 (b)

Let oxidation state of I in $IPO_4 = 'x'$.


$$x + (-3) = 0$$

(PO_4^{3-} ion has charge equal to -3)

$$x = +3$$

401 (a)

In alkaline medium



Hence, moles of KI = moles of $KMnO_4$.

402 (c)

5 mole I^- gives 3 mole I_2

403 (c)


$$2 \times 1 + 2 \times a + 7 \times (-2) = 0$$

$$\therefore a = +6$$

404 (c)


The concentration of standard solution is known.

405 (a)



$\therefore H_2S$ is oxidised in this reaction.

406 (b)




407 (b)

Iron usually shows zero, +2, +3 oxidation state.

408 (a)


Calculate ox.no. of Cl in $NOClO_4$ by assuming ClO_4^- and NO^+ .

409 (b)



H_2 is reduced and thus, oxidant.

410 (b)



411 (a)


$$a + 2 \times (-2) = 0$$

$$\therefore a = +4$$


412 (d)

Ox.no. Fe in Mohr's salt, $[FeSO_4 \cdot (NH_4)_2SO_4 \cdot 6H_2O]$ is +2.

413 (a)



414 (d)



415 (b)

Meq. of $K_2Cr_2O_7$ = Meq. of H_2S

$$2 \times V = \frac{0.81}{34/2} \times 1000$$


$$\therefore V = 23.8 \text{ mL}$$

416 (a)

$$3 \times 1 + a + 4 \times (-2) = 0$$

$$\therefore a = +5$$

417 (b)



$$\therefore E = \frac{M}{8/3} = \frac{56 \times 3}{8} = 21$$

418 (d)

Permonosulphuric acid (H_2SO_5) has two oxygen atoms in peroxide linkage, hence,


$$2(+1) + x + 2(-1) + 3(-2) = 0$$

$$2 + x - 2 - 6 = 0$$

$$x = +6$$






419 (c)

The reaction, in which two or more species undergo reduction as well as oxidation to give a single species are called comproportionation reaction. This is reverse of disproportionation reaction.



420 (d)

HCl is also oxidised along with oxalic acid by $KMnO_4$.



421 (a)

No change in oxidation no. in any of the species.

422 (d)

S in H_2SO_3 is in +4 oxidation state. It lies in between its maximum and minimum oxidation state, i.e., +6 and -2 and thus, S can increase or

decrease its ox.no.as the case may be.

SMART ACHIEVERS LEARNING PVT. LTD.

REDOX REACTIONS

CHEMISTRY

Assertion - Reasoning Type

This section contain(s) 0 questions numbered 1 to 0. Each question contains STATEMENT 1(Assertion) and STATEMENT 2(Reason). Each question has the 4 choices (a), (b), (c) and (d) out of which **ONLY ONE** is correct.

- a) Statement 1 is True, Statement 2 is True; Statement 2 **is** correct explanation for Statement 1
- b) Statement 1 is True, Statement 2 is True; Statement 2 **is not** correct explanation for Statement 1
- c) Statement 1 is True, Statement 2 is False
- d) Statement 1 is False, Statement 2 is True

1

Statement 1: Change in colour of acidic solution of potassium dichromate by breath is used to test drunk drivers.

Statement 2: Change in colour is due to the complexation of alcohol with potassium dichromate.

2

Statement 1: $\text{Mg(s)} + \text{F}_2\text{(s)} \rightarrow \text{MgF}_2\text{(s)}$: Magnesium loses electrons and acts as a reducing agent.

Statement 2: Reduction in general means acceptance of electrons by a reactant.

3

Statement 1: Bromide ion is serving as a reducing agent in a reaction.

Statement 2: Oxidation number of Br increases from -1 to 5.

4

Statement 1: Nitrous acid may act as an oxidizing agent as well as a reducing agent.

Statement 2: The oxidation number of nitrogen remains the same in all the compounds.

5

Statement 1: Both oxygen atoms in O_2 or O_3 have an oxidation number of -2.

Statement 2: Oxygen is assigned an oxidation number of -2 in almost all its compounds.

6

Statement 1: Oxidation number of Ni is zero.

Statement 2: Nickel is bonded to neutral ligand carbonyl.

REDOX REACTIONS

CHEMISTRY

: ANSWER KEY :

1)	c	2)	a	3)	a	4)	c
5)	d	6)	a				

SMART ACHIEVERS LEARNING PVT. LTD.

REDOX REACTIONS

CHEMISTRY

: HINTS AND SOLUTIONS :

- 1 **(c)**
Potassium dichromate react with alcohol and the reduction of potassium dichromate takes place and dichromate (orange red) changes to Cr^{3+} (green).
- 2 **(a)**
In general, oxidation is the loss of electrons and the reactant like, magnesium that loses electrons acts as a reductant or reducing agent.
- 3 **(a)**
+7
$$2 \text{MnO}_4^-(aq) + \text{Br}^-(aq) + \text{H}_2\text{O}(l) \rightarrow$$
$$\text{MnO}_2(aq) + \text{BrO}_3^-(aq) + 2\text{OH}^-(aq)$$

+ 4 + 5
Oxidation number is changes as -1 to + 5 (so loss of $6e^-$).
- 4 **(c)**
Oxidation number of N is changed according to compounds
(-1 to +5) N has five types of oxides as
- 5 **(d)**
Each atom in an elemental form is assigned an oxidation number of zero. For example, hydrogen atom in H_2 and oxygen atom in O_2 or O_3 , carbon in diamond and graphite, all have oxidation number equal to zero.
- 6 **(a)**
Oxidation number of N CO=0 (zero) as it a neutral ligand.
Oxidation number of Ni in $[\text{Ni}(\text{CO})_4]$ is also zero.