

## Tangent & Normals

### Single Correct Answer Type

1. The points of contact of the tangents drawn from the origin to the curve  $y = x^2 + 3x + 4$  are

1. (2, 14), (-2, 12)      2. (2, 12), (-2, 2)      3. (2, 14), (-2, 2)      4. (2, 12), (-2, 14)

Key. 3

Sol. Let  $P(x_1, y_1)$  be a point on the curve  $y = x^2 + 3x + 4$

$$\Rightarrow y_1 = x_1^2 + 3x_1 + 4 \quad \dots(1)$$

$$\left(\frac{dy}{dx}\right)_{at(x_1, y_1)} = 2x_1 + 3$$

Equation of tangent is :  $y - y_1 = m(x - x_1)$

It is passes through (0, 0)

$$\text{Then } y_1 = 2x_1^2 + 3x_1 \quad \dots(2)$$

From (1) & (2)  $x_1 = \pm 2$

$\therefore$  the points are (2,14)&(-2, 2)

2. If  $3x + 2y = 1$  acts as a tangent to  $y = f(x)$  at  $x = 1/2$  and if

$$p = \lim_{x \rightarrow 0} \frac{x(x-1)}{f\left(\frac{e^{2x}}{2}\right) - f\left(\frac{e^{-2x}}{2}\right)}, \text{ then, } \sum_{r=1}^{\infty} p^r = \underline{\hspace{2cm}}$$

- a) 1/2      b) 1/3      c) 1/6      d) 1/7

Key. A

Sol. slope of  $3x + 2y = 1$  is  $\frac{-3}{2}$

$$\Rightarrow f'\left(\frac{1}{2}\right) = \frac{-3}{2}$$

$$p = \lim_{x \rightarrow 0} \frac{x(x-1)}{f\left(\frac{e^{2x}}{2}\right) - f\left(\frac{e^{-2x}}{2}\right)} \left(\frac{0}{0}\right) = \frac{-1}{f'\left(\frac{1}{2}\right) + f'\left(\frac{1}{2}\right)} = \frac{1}{3}$$

$$\therefore \sum_{r=1}^{\infty} p^r = \frac{1}{3} + \frac{1}{3^2} + \dots \dots \infty = \frac{1/3}{1-1/3} = \frac{1/3}{2/3} = \frac{1}{2}$$

3. If the tangent drawn at  $P\left(t = \frac{\pi}{4}\right)$  to the curve  $x = \sec^2 t, y = \cot t$  meets the curve again at R, then, PR = \_\_\_\_\_

- a)  $\frac{3\sqrt{5}}{2}$       b)  $\frac{2\sqrt{5}}{3}$       c)  $\frac{5\sqrt{5}}{4}$       d)  $\frac{4\sqrt{5}}{5}$

Key. A

Sol. At  $t = \frac{\pi}{4}, x = 2, y = 1 \Rightarrow P$  is  $(2,1)$

$$\frac{dy}{dx} \Big|_{t=\frac{\pi}{4}} = \frac{-\operatorname{cosec}^2 t}{2 \operatorname{sect} . \operatorname{sect} . \operatorname{tant}} = -1/2$$

$$\therefore \text{tangent at } P(2,1) \text{ is, } y = \frac{4-x}{2}$$

$$\text{Eliminating 't' curve equation is, } x = 2, 5 \Rightarrow R(5, -1/2) \Rightarrow PR = \frac{3}{2}\sqrt{5}$$

4. If the points of contact of tangents to  $y = \sin x$ , drawn from origin always lie on  $\frac{a}{y^2} - \frac{b}{x^2} = c$ , then,
- a) a,b,c are in AP, but not in GP and HP
  - b) a,b,c are in GP, but not in HP and AP
  - c) a,b,c are in HP, but not in AP and GP
  - d) a,b,c are in AP, GP and HP

Key. D

Sol. Let  $P(h,k)$  be any point on  $y = \sin x$   
 $\Rightarrow k = \sinh$ . tangent P is  $y - k = \cosh(x - h)$

$$(0,0) \Rightarrow -k = \cosh.(0 - h) \Rightarrow \cosh = \frac{k}{h}$$

$$\Rightarrow \frac{1}{y^2} - \frac{1}{x^2} = 1 \Rightarrow a = 1, b = 1, c = 1$$

5.  $A(1,0), B(e,1)$  are two points on the curve  $y = \log_e x$ . If P is a point on the curve at which the tangent to the curve is parallel to the chord AB, then, abscissa of P, is

- a)  $\frac{e-1}{2}$
- b)  $\frac{e+1}{2}$
- c)  $e-1$
- d)  $e+1$

Key. C

Sol. By LMVT, applied to  $f(x) = \log_e x$  on  $[1, e], \exists \text{an } x_0 \in (1, e) \ni f'(x_0) = \frac{f(e) - f(1)}{e - 1}$   
 $\Rightarrow x_0 = e - 1$

6. The abscissa of the points. Where the tangent to the curve  $y = x^3 - 3x^2 - 9x + 5$  is parallel to x-axis is

- 1) 0 and 0
- 2)  $x=1$  and  $-1$
- 3)  $x=1$  and  $-3$
- 4)  $x=-1$  and  $3$

Key. 4

Sol.  $\Rightarrow \frac{dy}{dx} = 0 \Rightarrow x = -1, 3$   
 Tangent is parallel to x-axis

7. Co-ordinates of a point on the curve  $y = x \log x$  at which the normal is parallel to the line  $2x - 2y = 3$ , are

- 1) (0,0)                      2) (e,e)                      3)  $(e^{-2}, 2e^{-2})$                       4)  $(e^{-2}, -2e^{-2})$

Key. 4

Sol. Slope of the normal  $= \frac{-1}{1+\log x} \Rightarrow \frac{-1}{1+\log x} = 1 \Rightarrow x = e^{-2}$

8. If the point on  $y = x \tan \alpha - \frac{ax^2}{2u^2 \cos^2 \alpha}$  ( $0 < \alpha < \frac{\pi}{2}$ ) where the tangent is parallel to  $y=x$  has an ordinate  $\frac{u^2}{4a}$  then the value of  $\alpha$  is

- 1)  $\frac{\pi}{2}$                       2)  $\frac{\pi}{6}$                       3)  $\frac{\pi}{3}$                       4)  $\frac{\pi}{4}$

Key. 3

Sol. Given  $m=1 \Rightarrow \tan \alpha - \frac{ax}{u^2 \cos^2 \alpha} = 1 \Rightarrow x = \frac{(\tan \alpha - 1) u^2 \cos^2 \alpha}{a}$  substitute x and y values in

given equation  $\frac{u^2}{4a} = \frac{u^2}{a} \left[ \sin^2 \alpha - \frac{1}{2} \right] \Rightarrow \alpha = \frac{\pi}{3}$

9. If at each point of the curve  $y = x^3 - ax^2 + x + 1$  the tangent is inclined at an acute angle with the positive direction of the x-axis, then a lies in the interval

- 1)  $[-3, 3]$                       2)  $[-2, 2]$                       3)  $[-\sqrt{3}, \sqrt{3}]$                       4) R

Key. 3

Sol.  $\frac{dy}{dx} = 3x^2 - 2ax + 1, \frac{dy}{dx} > 0 \Rightarrow 3x^2 - 2ax + 1 > 0$

10. The number of tangents to the curve  $x^{\frac{3}{2}} + y^{\frac{3}{2}} = a^{\frac{3}{2}}$ , where the tangents are equally inclined to the axes, is

- 1) 2                      2) 1                      3) 0                      4) 4

Key. 2

Sol.  $\Rightarrow \frac{dy}{dx} = -\frac{x^{1/2}}{y^{1/2}}$

$\therefore \left(\frac{dy}{dx}\right)_{\alpha, \beta} = 1 \Rightarrow \alpha^{1/2} + \beta^{1/2} = 0$

$\alpha^{3/2} + \beta^{3/2} = a^{3/2}$  ( $\because (\alpha, \beta)$  is on the curve)

$\left(\frac{dy}{dx}\right)_{\alpha, \beta} = -1 \Rightarrow \alpha^{1/2} = \beta^{1/2}$

$\therefore \alpha = \beta = \frac{a}{2^{2/3}}$

there is only one point

11. The tangent at any point on the curve  $x = a \cos^3 \theta, y = a \sin^3 \theta$  meets the axes in P and Q. The locus of the mid point of PQ is

1)  $x^2 + y^2 = a^2$     2)  $2(x^2 + y^2) = a^2$     3)  $4(x^2 + y^2) = a^2$     4)  $x^2 + y^2 = 4a^2$

Key. 3

Sol. Equation of tangent at  $\theta$  is  $\Rightarrow P = (a \cos \theta, 0), Q = (0, a \sin \theta)$ . Locus of midpoint of PQ is  $4(x^2 + y^2) = a^2$

12. If the curves  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  and  $\frac{x^2}{l^2} - \frac{y^2}{m^2} = 1$  cut each other orthogonally then .

1)  $a^2 + b^2 = l^2 + m^2$     2)  $a^2 - b^2 = l^2 - m^2$     3)  $a^2 - b^2 = l^2 + m^2$     4)  $a^2 + b^2 = l^2 - m^2$

Key. 3

Sol. If the curves  $a_1x^2 + b_1y^2 = 1, a_2x^2 + b_2y^2 = 1$  cut each other orthogonally then apply

$$\frac{1}{a} - \frac{1}{b} = \frac{1}{a_1} - \frac{1}{b_1}$$

13. If the relation between the sub-normal and sub-tangent at any point on the curve

$y^2 = (x+a)^3$  is  $p(S.N) = q(S.T)^2$  then  $\frac{p}{q} =$

1)  $\frac{8}{27}$

2)  $\frac{27}{8}$

3)  $\frac{4}{9}$

4)  $\frac{9}{4}$

Key. 1

Sol. Length of sub normal =  $|y_1 m|$

Length of sub tangent =  $\left| \frac{y_1}{m} \right|$

14. The sum of the lengths of subtangent and tangent to the curve

$x = c \left[ 2 \cos \theta - \log (\operatorname{cosec} \theta + \cot \theta) \right], y = c \sin 2\theta$  at  $\theta = \frac{\pi}{3}$  is

1)  $\frac{c}{2}$

2)  $2c$

3)  $\frac{3c}{2}$

4)  $\frac{5c}{2}$

Key. 3

Sol. Length of tangent =  $\left| \frac{y_1 \sqrt{1+m^2}}{m} \right|$

Length of sub-tangent =  $\left| \frac{y_1}{m} \right|$

15. The curves  $C_1 : y = x^2 - 3; C_2 : y = kx^2, k < 1$  intersect each other at two different points. The tangent drawn to  $C_2$ , at one of the points of intersection  $A = (a, y_1) (a > 0)$  meets  $C_1$  again at  $B(1, y_2)$ . ( $y_1 \neq y_2$ ). Then value of  $a = \underline{\hspace{1cm}}$ ?

a) 4

b) 3

c) 2

d) 1

Sol : ans: b  
solving

$C_1 \& C_2 \Rightarrow A \left( \sqrt{\frac{3}{1-k}}, \frac{3k}{1-k} \right) = (a, ka^2) \equiv (a, a^2 - 3)$ .

tangent 1 to  $C_2$  at A is  $y + a^2 - 3 = 2kx$  ----- (1)

$\Rightarrow B = (1, -2) (A \neq 1)$ .

from expression (1)  $-2 + a^2 - 3 = 2a \left( 1 - \frac{3}{a^2} \right)$ .

$\Rightarrow a = 3, a = -2, a = 1$

$\therefore a = 3$

16. Let  $f\left(\frac{x+y}{2}\right) = \frac{1}{2}(f(x)+f(y))$  for real  $x$  and  $y$ . If  $f'(0)$  exists and equals to  $-1$  and  $f(0)=1$  then the value of  $f(2)$  is
- a) 1                                      b)  $-1$                                       c)  $\frac{1}{2}$                                       d) 2

KEY : B

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\frac{f(2x) + f(2h)}{2} - f(x)}{h}$$

$$f'(x) = -1 \quad ; \quad f(2x) = 2f(x) - 1$$

$$\Rightarrow f(x) = 1 - x$$

17. If the length of subnormal is equal to length of sub-tangent at point  $(3,4)$  on the curve  $y = f(x)$  and the tangent at  $(3,4)$  to  $y = f(x)$  meets the coordinate axes at A and B, then maximum area of the  $\Delta OAB$  where O is origin, is
- (A)  $\frac{45}{2}$  squnits                                      (B)  $\frac{49}{2}$  squnits
- (C)  $\frac{51}{2}$  squnits                                      (D)  $\frac{81}{2}$  squnits

KEY : B

Sol : Length of subnormal = length of subtangent

$$\Rightarrow \left| y_1 \left( \frac{dy}{dx} \right)_{(x_1, y_1)} \right| = \left| \frac{y_1}{\left( \frac{dy}{dx} \right)_{(x_1, y_1)}} \right|$$

$$\Rightarrow \left( \frac{dy}{dx} \right)_{(x_1, y_1)} = \pm 1$$

$$\text{If } \left( \frac{dy}{dx} \right)_{(x_1, y_1)} = 1$$

Then the equation of tangent is  $y - x = 1$  and area of  $\Delta OAB = \frac{1}{2} \times 1 \times 1 = \frac{1}{2}$

$$\text{If } \left( \frac{dy}{dx} \right)_{(x_1, y_1)} = -1$$

Then the equation of tangent is  $x + y = 7$  and area of  $\Delta OAB = \frac{1}{2} \times 7 \times 7 = \frac{49}{2}$

18. The equation of normal to the curve  $x + y = x^y$ , where it cuts the x-axis is
- (A)  $y = x - 1$                                       (B)  $x + y = 1$
- (C)  $12x + y + 2 = 0$                                       (D)  $3x + y = 3$

Key: A

Sol: At x-axis,  $y = 0 \Rightarrow x = 1$

$$x + y = x^y \Rightarrow \ln(x + y) = y \ln x$$

$$\frac{1}{x + y} \left( 1 + \frac{dy}{dx} \right) = \frac{y}{x} + \frac{dy}{dx} \ln x$$

$$\left( \frac{dy}{dx} \right) (1, 0) = -1$$

So equation of normal  $y - 0 = x - 1$ .

19. Maximum no. of parallel tangents of curves  $y = x^3 - x^2 - 2x + 5$  and  $y = x^2 - x + 3$  is

(A) 2

(B) 3

(C) 4

(D) none of these

Key: D

Sol: Let  $m$  be slope is common tangent

$$\text{Then } m = 2x - 1 \text{ and } m = 3x^2 - 2x - 2,$$

So, infinite common tangents

20. The equation of the straight lines which are both tangent and normal to the curve  $27x^2 = 4y^3$  are

a)  $x = \pm\sqrt{2}(y - 2)$

b)  $x = \pm\sqrt{3}(y - 2)$

c)  $x = \pm\sqrt{2}(y - 3)$

d)  $x = \pm\sqrt{3}(y - 3)$

Key: A

Sol.  $x = 2t^3, y = 3t^2 \Rightarrow$  tangent at  $t$  is  $x - yt = -t^3$  Normal at  $t_1$  is,  $xt_1 + y = 2t_1^4 + 3t_1^2$

$$\Rightarrow \frac{1}{t_1} = -t = \frac{-t^3}{2t_1^4 + 3t_1^2} \Rightarrow t^6 - 3t^2 - 2 = 0 \Rightarrow t^2 = 2 \Rightarrow t = \pm\sqrt{2}$$

$\therefore$  lines are  $x = \pm\sqrt{2}(y - 2)$

21. If  $f(x) + f(y) = f(x)f(y) + f(xy), f(1) = 0, f'(1) = -2$  then, equation to the tangent, drawn to the curve  $y = f(x)$  at  $x = \sqrt{2}$  is,

a)  $2\sqrt{2}x - y - 3 = 0$

b)  $2\sqrt{2}x + y - 3 = 0$

c)  $2\sqrt{2}x + y + \sqrt{3} = 0$

d)  $2\sqrt{2}x + 2y - 3 = 0$

Key: B

Sol. Clearly  $f(x) = 1 - x^2$  at  $x = \sqrt{2}, y = -1 \Rightarrow$  tangent at  $(\sqrt{2}, -1)$  is,

$$y + 1 = -2\sqrt{2}(x - \sqrt{2})$$

22. Let  $f(x)$  be a polynomial of degree 5. When  $f(x)$  is divided by  $(x - 1)^3$ , the remainder 33, and when  $f(x)$  is divided by  $(x + 1)^3$ , the remainder is  $-3$ . Then, equation to the tangent drawn to  $y = f(x)$  at  $x = 0$  is

a)  $135x + 4y + 60 = 0$

b)  $135x - 4y - 60 = 0$

c)  $135x - 4y + 60 = 0$

d)  $135x - 4y + 75 = 0$

Key: C

Sol.  $f(x) = \frac{27x^5}{4} - \frac{45x^3}{2} + \frac{135x}{4} + 15$  at  $x = 0, y = 15 \Rightarrow f'(0) = \frac{135}{4}$

$$\Rightarrow \text{tangent equation is } y - 15 = \frac{135}{4}(x) \Rightarrow 135x - 4y + 60 = 0$$

23. If the equation  $x^{5/3} - 5x^{2/3} = K$  has exactly one positive root, then, the complete solution set of K is,

- a)  $(-\infty, \infty)$                       b)  $(-\infty, 0)$                       c)  $(3, \infty)$                       d)  $(0, \infty)$

Key. D

Sol. Sketch  $y = x^{5/3} - 5x^{2/3}$  and  $y = K$

24. The equation of the straight lines which are both tangent and normal to the curve  $27x^2 = 4y^3$  are

- a)  $x = \pm\sqrt{2}(y - 2)$                       b)  $x = \pm\sqrt{3}(y - 2)$   
 c)  $x = \pm\sqrt{2}(y - 3)$                       d)  $x = \pm\sqrt{3}(y - 3)$

Key. A

Sol.  $x = 2t^3, y = 3t^2 \Rightarrow$  tangent at t is  $x - yt = -t^3$  Normal at  $t_1$  is,  $xt_1 + y = 2t_1^4 + 3t_1^2$

$$\Rightarrow \frac{1}{t_1} = -t = \frac{-t^3}{2t_1^4 + 3t_1^2} \Rightarrow t^6 - 3t^2 - 2 = 0 \Rightarrow t^2 = 2 \Rightarrow t = \pm\sqrt{2}$$

$$\therefore \text{lines are } x = \pm\sqrt{2}(y - 2)$$

25. If  $f(x) + f(y) = f(x)f(y) + f(xy), f(1) = 0, f'(1) = -2$  then, equation to the tangent, drawn to the curve  $y = f(x)$  at  $x = \sqrt{2}$  is,

- a)  $2\sqrt{2}x - y - 3 = 0$                       b)  $2\sqrt{2}x + y - 3 = 0$   
 c)  $2\sqrt{2}x + y + \sqrt{3} = 0$                       d)  $2\sqrt{2}x + 2y - 3 = 0$

Key. B

Sol. Clearly  $f(x) = 1 - x^2$  at  $x = \sqrt{2}, y = -1 \Rightarrow$  tangent at  $(\sqrt{2}, -1)$  is,  
 $y + 1 = -2\sqrt{2}(x - \sqrt{2})$

26. Let  $f(x)$  be a polynomial of degree 5. When  $f(x)$  is divided by  $(x - 1)^3$ , the remainder 33, and when  $f(x)$  is divided by  $(x + 1)^3$ , the remainder is -3. Then, equation to the tangent drawn to  $y = f(x)$  at  $x = 0$  is

- a)  $135x + 4y + 60 = 0$                       b)  $135x - 4y - 60 = 0$   
 c)  $135x - 4y + 60 = 0$                       d)  $135x - 4y + 75 = 0$

Key. C

Sol.  $f(x) = \frac{27x^5}{4} - \frac{45x^3}{2} + \frac{135x}{4} + 15$  at  $x = 0, y = 15 \Rightarrow f'(0) = \frac{135}{4}$

$$\Rightarrow \text{tangent equation is } y - 15 = \frac{135}{4}(x) \Rightarrow 135x - 4y + 60 = 0$$

27. Two runners A and B start at the origin and run along positive x-axis, with B running three times as fast as A. An observer, standing one unit above the origin, keeps A and B in view. Then the maximum angle of sight 'θ' between the observes view of A and B is

- a)  $\pi/8$                       b)  $\pi/6$                       c)  $\pi/3$                       d)  $\pi/4$


Key. B



Sol.  $\tan \theta = \tan(\theta_2 - \theta_1) \Rightarrow \tan \theta = \frac{3x - x}{1 + 3x \cdot x} = \frac{2x}{1 + 3x^2}$

let  $y = \frac{2x}{1 + 3x^2}$   $\frac{dy}{dx} = \frac{2(1 - 3x^2)}{(1 + 3x^2)^2}$

$\frac{dy}{dx} = 0 \Rightarrow x = \frac{1}{\sqrt{3}}$  and  $\frac{d^2y}{dx^2} = \frac{-24x}{(1 + 3x^2)^3} < 0$  for  $x = 1/\sqrt{3}$   
 $\Rightarrow \theta = \pi \setminus 6$



28. If the line joining the points (0,3) and (5,- 2) is a tangent to the curve  $y = \frac{c}{x+1}$ , then value of c is  
 A) 1                                      B) -2                                      C) 4                                      D) -4.

Key. 3

Sol. Eqn. of the line joining given points is  $(y + 2) = \frac{-2 - 3}{5 - 0}(x - 5)$ .

$\Rightarrow y + x = 3$ .

29. The number of points on the curve  $y^3 - 3xy + 2 = 0$  where the tangent is either horizontal or vertical is  
 A) 0                                      B) 1                                      C) 2                                      D) > 2.

Key. 2

Sol.  $3yy' - 3y - 3xy' = 0 \Rightarrow y' = \frac{y}{y^2 - x}$ .

$y' = 0 \Rightarrow y = 0$ , no real x

$y' = \infty \Rightarrow y^2 = x \Rightarrow y^3 = 1 \Rightarrow y = 1$ .

The point is (1,1).

30. The tangent to the curve  $y = \frac{1+3x^2}{3+x^2}$  drawn at the points for which  $y = 1$ , intersect at

- A) (0,0)                                      B) (0,1)                                      C) (1,0)                                      D) (1,1)

Key. 1

Sol.  $y = 1 \Rightarrow x = \pm 1$  points are (1,1), (-1,1)  $\Rightarrow \frac{dy}{dx} = \frac{16x}{(3+x^2)^2}$ ,  $\left(\frac{dy}{dx}\right)_{(1,1)} = 1$ ,  $\left(\frac{dy}{dx}\right)_{(-1,1)} = -1$

Eq. of tangent at (1,1) is  $y - 1 = (x - 1) \Rightarrow x - y = 0$

Eq. of tangent at (-1, 1)  $y - 1 = -1(x + 1) \Rightarrow x + y = 0$

Both tangents pass through origin.

31. The equation of the normal to the curve  $x + y = x^y$ , where it cuts x-axis is



Comparing we get,  $-t^3 = 2t_1^3 + 3t_1, t_1 = \frac{1}{t}$ . Eliminating  $t_1$ , we get  $t^6 = 2 + 3t^2$ .

Ⓡ  $t^2 = 2, t = \pm\sqrt{2}$

35. The tangent at any point P on the curve  $x^{2/3} + y^{2/3} = 4$  meets the coordinate axes at A and B Then  $AB =$   
 A) 2                                      B) 4                                      C) 8                                      D) 16

Key. 3

Sol.  $x = 8\cos^3 q, y = 8\sin^3 q, \frac{dy}{dx} = -\frac{\sin q}{\cos q}$ .

Tangent at  $q, y - 8\sin^2 q = -\frac{\sin q}{\cos q}(x - 8\cos^3 q)$

$x\sin q + y\cos q = 8\sin q\cos q$

$OA = 8\cos q, OB = 8\sin q$

$AB = \sqrt{OA^2 + OB^2} = 8$ .

36. If the tangent to the curve  $x = 1 - 3t^2, y = t - 3t^3$  at the point  $P(-2, 2)$  meets the curve again at Q, the angle between the tangents at P and Q is

- A)  $\frac{\pi}{6}$                                       B)  $\frac{\pi}{4}$                                       C)  $\frac{\pi}{3}$                                       D)  $\frac{\pi}{2}$ .

Key. 4

Sol.  $\frac{dy}{dx} = \frac{9t^2 - 1}{6t}$

$x = -2, y = 2$  Ⓡ  $t = -1, \frac{dy}{dx} = -\frac{4}{3}$

The tangent at P,  $y - 2 = -\frac{4}{3}(x + 2)$  Ⓡ  $4x + 3y = -2$ .

$4(1 - 3t^2) + 3(t - 3t^3) = -2$

Ⓡ  $(t + 1)^2(3t - 2) = 0$

$t = \frac{2}{3}$ , Slope of tangent at Q is  $\frac{dy}{dx} = \frac{9\left(\frac{2}{3}\right)^2 - 1}{6\left(\frac{2}{3}\right)} = \frac{3}{4}$ .

(1), (2) Ⓡ The tangents are perpendicular.

37. The curves  $x^3 - 3xy^2 = a$  and  $3x^2y - y^3 = b$  intersect at an angle of

- A)  $\frac{\pi}{4}$                                       B)  $\frac{\pi}{3}$                                       C)  $\frac{\pi}{2}$                                       D)  $\frac{\pi}{6}$ .

Key. 3

Sol. Clearly  $m_1 m_2 = -1$ .

38. The cosine of the angle of intersection of curves  $f(x) = 2^x \log_e x$  and  $g(x) = x^{2x} - 1$  is

- A) 1                                      B) 0                                      C)  $\frac{1}{2}$                                       D)  $\frac{\sqrt{3}}{2}$ .

Key. 1

Sol. Clearly, (1,0) is the point of intersection of the given curves.

Now,  $f'(x) = \frac{2^x}{x} + 2^x (\log_e 2)(\log_e x)$

\ Slope of tangent to the curve  $f(x)$  at (1,0),  $m_1 = 2$ .

$g'(x) = \frac{d}{dx}(e^{2x \log x} - 1) = x^{2x} \left( \frac{2}{x} + 2 \log_e x \right)$

\ Slope of tangent to the curve  $g(x)$  at (1,0),  $m_2 = 2$ .

Since  $m_1 = m_2 = 2$ .

\ Two curves touch each other, so the angle between them is 0.

Hence,  $\cos q = \cos 0 = 1$ .

39. The curves  $\frac{x^2}{a} + \frac{y^2}{b} = 1$  and  $\frac{x^2}{a_1} + \frac{y^2}{b_1} = 1$  will cut orthogonally if

- A)  $ab = a_1b_1$                       B)  $\frac{a}{b} = \frac{a_1}{b_1}$                       C)  $a + b = a_1 + b_1$                       D)

$a - b = a_1 - b_1$

Key. 4

Sol.  $\frac{x^2}{a} + \frac{y^2}{b} = 1 \dots (1)$

$\frac{x^2}{a_1} + \frac{y^2}{b_1} = 1 \dots (2)$

(1)-(2)  $\otimes \quad x^2 \left( \frac{1}{a} - \frac{1}{a_1} \right) + y^2 \left( \frac{1}{b} - \frac{1}{b_1} \right) = 0$ .

$\otimes \quad \frac{x^2(a_1 - a)}{a_1 a} = - \frac{y^2(b_1 - b)}{b_1 b} \dots (3)$

Differentiating (1),  $\frac{x}{a} + \frac{ym_1}{b} = 0$

$\therefore m_1 = - \frac{bx}{ay}, m_2 = - \frac{b_1x}{a_1y}$

$m_1 m_2 = \frac{b_1 b x^2}{a a_1 y^2} = - \frac{a_1 b_1 - b \frac{a}{a_1}}{a_1 - \frac{a}{a_1}} = -1$ .

40. The value of n in the equation of curve  $y = a^{-n} x^n$ , so that the sub-normal may be of constant length is

- A) 2                      B)  $\frac{3}{2}$                       C)  $\frac{1}{2}$                       D) 1

Key. 3

Sol. Taking log and differentiating both sides, we get  $\frac{dy}{dx} = \frac{ny}{x} \dots (1)$

Length of sub-normal =  $na^{2-2n} \cdot x^{2n-1}$

$$n = \frac{1}{2}$$

41. Let  $f(x) = x^2 + xg'(1) + g''(2)$  and  $g(x) = f(1)x^2 + xf'(x) + f''(x)$ , then  $f(3) + g(3) =$   
 A) 7                                      B) -7                                      C) 0                                      D) 6

Key. 2

Sol. Let  $g'(1) = a, g''(1) = b$  then  $f(x) = x^2 + ax + b$  then  $f(1) = 1 + a + b$

$$g(x) = (1 + a + b)x^2 + x(2x + a) + b$$

$$g'(x) = 2x(3 + a + b) + a$$

$$g'(1) = a \Rightarrow a + b + 3 = 0, \quad g''(2) = b \Rightarrow 2a + b = -6$$

42. Tangents are drawn from origin to the curve  $y = \sin x + \cos x$ . Then their points of contact lie on the curve

a)  $\frac{1}{x^2} + \frac{2}{y^2} = 1$                       b)  $\frac{2}{x^2} - \frac{1}{y^2} = 1$                       c)  $\frac{2}{x^2} + \frac{1}{y^2} = 1$                       d)  $\frac{2}{y^2} - \frac{1}{x^2} = 1$

Key. D

Sol.  $y_1 = \sqrt{2} \sin\left(x_1 + \frac{\pi}{4}\right), \frac{dy}{dx} = \frac{y_1}{x_1}$  where  $(x_1, y_1)$  is point on the curve

$$\frac{y_1^2}{x_1^2} = 2 \cos^2\left(x_1 + \frac{\pi}{4}\right) + \frac{y_1^2}{4x_1^2} = 2 \cos^2\left(x_1 + \frac{\pi}{4}\right) + \frac{y_1^2}{4x_1^2}$$

$$\Rightarrow \text{Locus of } (x_1, y_1) \text{ is } \frac{2}{y^2} - \frac{1}{x^2} = 1$$

43. The abscissa of two points on  $y = (2010)x^2 + (2011)x - 2011$  are 2010 and 2012. If the chord joining those two points is parallel to tangent at P on the curve then the ordinate of P is equal to

- a) (2009)(2010)(2011)    b) (2010)(2011)(2012)  
 c) (2011)(2012)(2013)    d) none

Key. B

Sol. Apply LMVT with  $a = 2010, b = 2012$

$$f(x) = 2010x^2 + 2011x - 2011.$$

$$\frac{f(b) - f(a)}{b - a} = f'(c) \Rightarrow c = 2011, f(c) = (2010)(2011)(2012)$$

44. Tangent at  $P_1$  other than origin on the curve  $y = x^3$  meets the curve again at  $P_2$ . The tangent at  $P_2$  meets the curve again at  $P_3$  and so on ..... then  $\frac{\text{area of } DP_1P_2P_3}{\text{area of } DP_2P_3P_4}$  equals

- a) 1 : 20                                      b) 1 : 16                                      c) 1 : 8                                      d) 1 : 2

Key. B

Sol. Let  $P_1 = (t_1, t_1^3), P_2 = (t_2, t_2^3), P_3 = (t_3, t_3^3) \dots$

Solving tangent equation at  $P_1$  with the curve again we get  $t_2 = -2t_1$ . Repeating the process

we have  $t_3 = 4t_1$        $t_4 = -8t_1 \dots$

$$\therefore \frac{\Delta P_1 P_2 P_3}{\Delta P_2 P_3 P_4} = \frac{\begin{vmatrix} t_1 & t_1^3 & 1 \\ t_2 & t_2^3 & 1 \\ t_3 & t_3^3 & 1 \end{vmatrix}}{\begin{vmatrix} t_2 & t_2^3 & 1 \\ t_3 & t_3^3 & 1 \\ t_4 & t_4^3 & 1 \end{vmatrix}} = \frac{1}{16}$$

45. The value of parameter  $t$  so that the line  $(4-t)x + ty + (a^3 - 1) = 0$  is normal to the curve  $xy = 1$  may lie in the interval

- A)  $(1, 4)$       B)  $(-\infty, 0) \cup (4, \infty)$       C)  $(-4, 4)$       D)  $[3, 4]$

Key. B

Sol. Slope of line  $(4-t)x + ty + (a^3 - 1) = 0$

is  $\frac{-(4-t)}{t}$  (or)  $\frac{t-4}{t}$

$\therefore xy = 1$

$\therefore \frac{dy}{dx} = \frac{-y}{x} = \frac{-1}{x^2}$

$\therefore$  slope of normal  $= x^2 = \frac{t-4}{t}$

$\therefore x^2 > 0$

$\frac{t-4}{t} > 0$

$t \in (-\infty, 0) \cup (4, \infty)$

46. The angle of intersection of curves  $y = \lceil |\sin x| + |\cos x| \rceil$  and  $x^2 + y^2 = 5$ , where  $\lceil \cdot \rceil$  denotes greatest integral function is

- A)  $\tan^{-1}(2)$       B)  $\tan^{-1}(\sqrt{2})$       C)  $\tan^{-1}(\sqrt{3})$       D)  $\tan^{-1}(3)$

Key. A

Sol. We know that  $1 \leq |\sin x| + |\cos x| \leq \sqrt{2}$

$\therefore y = \lceil |\sin x| + |\cos x| \rceil = 1$

Let P and Q be the points of intersection of given curves clearly the given curves meet at points where  $y = 1$ , so we get

$x^2 + 1 = 5$

$\Rightarrow x = \pm 2$

$\therefore P(2, 1)$  and  $Q(-2, 1)$

Now  $x^2 + y^2 = 5$

$\Rightarrow x = \pm 2$

$\therefore P(2,1)$  and  $Q(-2,1)$

Now  $x^2 + y^2 = 5$

$\Rightarrow \frac{dy}{dx} = \frac{-x}{y}, \left(\frac{dy}{dx}\right)_{(2,1)} = -2, \left(\frac{dy}{dx}\right)_{(-2,1)} = 2$

Clearly the slope of a line  $y = 1$ , is 0 and the slope of tangent at P and Q are -2 and 2 respectively.

$\therefore$  The angle of intersection is  $\tan^{-1}(2)$

47. If the tangent at (1, 1) on  $y^2 = x(2-x)^2$  meets the curve again at P, then P is  
 a) (4, 4)                                      b) (-1, 2)                                      c) (9/4, 3/8) d) (9/5, 3/8)

Key. C

Sol.  $2y \frac{dy}{dx} = (2-x)^2 - 2x(2-x)$ , so  $\left.\frac{dy}{dx}\right|_{(1,1)} = -\frac{1}{2}$  Therefore, the equation of tangent at

(1, 1) is

$y - 1 = -\frac{1}{2}(x - 1)$

$\Rightarrow y = \frac{-x + 3}{2}$

The intersection of the tangent and the curve is given by  $(1/4)(-x+3)^2 = x(4+x^2-4x)$

$\Rightarrow x^2 - 6x + 9 = 16x + 4x^3 - 16x^2$

$\Rightarrow 4x^3 - 17x^2 + 22x - 9 = 0$

$\Rightarrow (x-1)(4x^2 - 13x + 9) = 0 \Rightarrow (x-1)^2(4x-9) = 0$

Since  $x = 1$  is already the point of tangency,  $x = 9/4$  and  $y^2 = \frac{9}{4}\left(2 - \frac{9}{4}\right)^2 = \frac{9}{24}$ . Thus the required point is  $(9/4, 3/8)$ .

48. The equation of the normal to the curve parametrically represented by  $x = t^2 + 3t - 8$  and  $y = 2t^2 - 2t - 5$  at the point P(2, -1) is  
 a)  $2x + 3y - 1 = 0$                                       b)  $6x - 7y - 11 = 0$   
 c)  $7x + 6y - 8 = 0$                                       d)  $3x + y - 1 = 0$

Key. C

Sol. 
$$\left. \begin{aligned} t^2 + 3t - 8 = 2 &\Rightarrow t = 2, -5 \\ 2t^2 - 2t - 5 = -1 &\Rightarrow t = 2, -1 \end{aligned} \right\} \Rightarrow t = 2, \frac{dy}{dx} = \frac{4t-2}{2t+3} \Rightarrow \left( \frac{dy}{dx} \right)_{t=2} = \frac{6}{7}$$

*equation of normal*  $y + 1 = \frac{-7}{6}(x - 2)$

49. Tangents are drawn from origin to the curve  $y = \cos x$ , their points of contact lie on the curve

a)  $x^2 + y^2 = x^2 y^2$

b)  $y^2 - x^2 = x^2 y^2$

c)  $x^2 + y^2 = 1$

d)  $x^2 - y^2 = x^2 y^2$

Key. D

Sol. Let point of contact is (h, k)

$\Rightarrow k = \cos h$  --- (1) *eq. of tangent at (h, k)*  $y - k = -\sin h(x - h)$ , it passes through origin  $\Rightarrow -k = h \sin h$  --- (2)

$\cos^2 h + \sin^2 h = k^2 + \frac{k^2}{h^2} \Rightarrow 1 = y^2 + \frac{y^2}{x^2}$  is the locus of point of contact

50. The angle between tangents at the point of intersection of two curves

$x^3 - 3xy^2 + 2 = 0, 3x^2y - y^3 = 2$  is

a)  $\frac{\pi}{6}$

b)  $\frac{\pi}{4}$

c)  $\frac{\pi}{3}$

d)  $\frac{\pi}{2}$

Key. D

Sol. Let the point of intersection is (x, y)

$x^3 - 3xy^2 + 2 = 0 \Rightarrow \frac{dy}{dx} = \frac{x^2 - y^2}{2xy}, 3x^2y - y^3 = 2 \Rightarrow \frac{dy}{dx} = \frac{2xy}{y^2 - x^2}, m_1 m_2 = -1 \Rightarrow \theta = 90^\circ$

51. Let the equation of a curve in parametric form be  $x = a(\theta + \sin \theta), y = a(1 - \cos \theta)$ . The

angle between the tangent drawn at the point  $\theta = \frac{\pi}{3}$  and normal drawn at the point

$\theta = \frac{2\pi}{3}$  is

a)  $\frac{\pi}{6}$

b)  $\frac{\pi}{4}$

c)  $\frac{\pi}{3}$

d)  $\frac{\pi}{2}$

Key. C

Sol.  $\frac{dy}{dx} = \frac{a \sin \theta}{a(1 + \cos \theta)} = \tan \frac{\theta}{2}$



$$m_1 = \tan \frac{\pi}{3} = \frac{1}{\sqrt{3}}, m_2 = -\frac{1}{\tan \frac{\theta}{2}} = -\frac{1}{\tan \frac{\pi}{3}} = -\frac{1}{\sqrt{3}}, \tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right| = \sqrt{3} \Rightarrow \theta = 60^\circ$$

52. Let the equation of a curve be  $\frac{x^2}{4} + \frac{y^2}{3} = 1$  where  $(2 \cos \theta, \sqrt{3} \sin \theta)$  is a general point on the curve. If the tangent to the given curve intersects the co-ordinate axes at points A, B, then the locus of midpoint of AB is

- a)  $2x^2 + \sqrt{3}y^2 = 4$                       b)  $3x^2 + 4y^2 = 4x^2y^2$   
 c)  $3x^2 + 4y^2 = x^2y^2$                       d)  $4x^2 + 3y^2 = 4x^2y^2$

Key.

B

Sol. Equation of tangent is

$$y - \sqrt{3} \sin \theta = \frac{-\sqrt{3}}{2} \cdot \cot \theta (x - 2 \cos \theta) \Rightarrow x \text{ intercept } (x_0) = \frac{2}{\cos \theta} \Rightarrow \cos \theta = \frac{2}{x_0},$$

$$y \text{ intercept } (y_0) = \frac{\sqrt{3}}{\sin \theta} \Rightarrow \sin \theta = \frac{\sqrt{3}}{y_0}, \text{ if mid point is } (h, k)$$

$$h = \frac{x_0}{2}, k = \frac{y_0}{2}, \cos \theta = \frac{1}{h}, \sin \theta = \frac{\sqrt{3}}{2k} \Rightarrow \frac{1}{h^2} + \frac{3}{4k^2} = 1$$

53. If the curve  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  and  $\frac{x^2}{\alpha^2} - \frac{y^2}{\beta^2} = 1$  cut each other orthogonally, then

- a)  $a^2 + b^2 = \alpha^2 + \beta^2$                       b)  $a^2 - b^2 = \alpha^2 - \beta^2$   
 c)  $a^2 - b^2 = \alpha^2 + \beta^2$                       d)  $a^2 + b^2 = \alpha^2 - \beta^2$

Key.

C

Slope of  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  at  $P(x_0, y_0)$  is  $-\frac{b^2 x_0}{a^2 y_0}$ , Slope of  $\frac{x^2}{\alpha^2} - \frac{y^2}{\beta^2} = 1$  at  $P(x_0, y_0)$  is  $\frac{\beta^2 x_0}{\alpha^2 y_0}$

Sol.

$$\therefore M_1 M_2 = -1 \Rightarrow b^2 \beta^2 x_0^2 = a^2 \alpha^2 y_0^2 \text{ ---- (1)}$$

now solving the curves

$$x_0^2 \left( \frac{1}{a^2} - \frac{1}{\alpha^2} \right) = -y_0^2 \left( \frac{1}{b^2} + \frac{1}{\beta^2} \right) \text{ ---- (2)}$$

from (1) & (2)

$$\frac{\frac{1}{a^2} - \frac{1}{\alpha^2}}{\frac{1}{b^2} + \frac{1}{\beta^2}} = \frac{b^2 \beta^2}{a^2 \alpha^2} \Rightarrow a^2 - b^2 = \alpha^2 + \beta^2$$

54. The rate of change of  $\sqrt{x^2 + 16}$  with respect to  $\frac{x}{x-1}$  at  $x = 3$  is

a) 1b)  $\frac{11}{5}$

c)  $-\frac{12}{5}$

d) -3

Key. C

Sol.  $u = \sqrt{x^2 + 16} \frac{du}{dx} = \frac{2x}{2\sqrt{x^2 + 16}} = \frac{x}{\sqrt{x^2 + 16}}, V = \frac{x}{x-1} \Rightarrow \frac{dv}{dx} = \frac{-1}{(x-1)^2}$

Sol.

$\frac{du}{dv} = \frac{du/dx}{dv/dx} = \frac{-12}{5}$

55. A curve represented parametrically by the equation  $x = t^3 - 4t^2 - 3t$  and  $y = 2t^2 + 3t - 5$  where  $t \in R$ . If H denotes the number of point(s) on the curve where the tangent is horizontal and V is the number of point(s) where the tangent is vertical then

a) H = 2, V = 1

b) H = 1, V = 2

c) H = 2, V = 2

d) H = 1, V = 1

Key. B

Sol.  $\frac{dy}{dt} = 4t + 3, \frac{dx}{dt} = 3t^2 - 8t - 3$  Tangents are horizontal if  $\frac{dy}{dx} = 0$

$\Rightarrow \frac{dy}{dt} = 0, \frac{dx}{dt} \neq 0, 4t + 3 = 0 \Rightarrow t = -\frac{3}{4}$

Tangents are vertical if  $\frac{dx}{dy} = 0, \frac{dx}{dt} = 0, \frac{dy}{dt} \neq 0, 3t^2 - 8t - 3 = 0 \Rightarrow t = 3, -\frac{1}{3}$

56. The tangents to the curve  $y = \frac{1+3x^2}{3+x^2}$  drawn at the points for which  $y = 1$ , intersect at

a) (0, 0)

b) (0, 1)

c) (1, 0)

d) (1, 1)

Key. A

Sol.  $y = 1 \Rightarrow x = \pm 1$  points are (1,1), (-1,1)  $\Rightarrow \frac{dy}{dx} = \frac{16x}{(3+x^2)^2}, \left(\frac{dy}{dx}\right)_{(1,1)} = 1, \left(\frac{dy}{dx}\right)_{(-1,1)} = -1$

Eq. of tangent at (1,1) is  $y - 1 = (x - 1) \Rightarrow x - y = 0$

Eq. of tangent at (-1, 1)  $y - 1 = -1(x + 1) \Rightarrow x + y = 0$

Both tangents pass through origin.

57. A cyclist moving on a level road at 4 m/s stops, pedalling and free wheels to rest. The retardation of the cycle has two components, a constant  $0.08 \text{ m/s}^2$  due to friction in the working parts, and resistance of  $0.02 \text{ v}^2/\text{s}^2$  where v is speed in meter per second. The distance traversed by the cycle before it comes to rest (approximately) is

a)  $40\frac{1}{4} \text{ mts}$

b)  $40\frac{1}{2} \text{ mts}$

c)  $20\frac{1}{2} \text{ mts}$

d)  $20\frac{1}{4} \text{ mts}$

Key. A

Sol. Let x be the displacement of the particle and let its acceleration of particle at P is a

$v = \frac{dx}{dt}$  and  $a = \frac{dv}{dt} = \frac{d^2x}{dt^2} = v \frac{dv}{dx}$

By data retardation =  $0.08 + 0.02 v^2 = 0.02(4 + v^2)$

$$v \frac{dx}{dt} = -0.02(4 + v^2) \Rightarrow \int_0^{x^1} dx = \frac{-1}{0.04} \int_4^0 \frac{2v}{4 + v^2} dv \Rightarrow x^1 = \frac{\log 5}{0.04} \approx 40 \frac{1}{4} \text{ mts}$$

58. For  $x = t^2 - 1, y = t^2 - t$ , the tangent line is perpendicular to x-axis when

- A)  $t = 0$                       B)  $t = \infty$                       C)  $t = 1/\sqrt{3}$                       D)  $t = -1/\sqrt{3}$

Key. A

Sol.  $\frac{dy}{dx} = \tan \theta = \infty \Rightarrow \frac{dx}{dy} = 0 \quad \frac{dy}{dt} \neq 0$   
 $2t = 0$  and  $2t - 1 \neq 0 \Rightarrow t = 0$  and  $t \neq 1/2$

59. The acute angle between the curves  $y = |x^2 - 1|$  and  $y = |x^2 - 3|$  at their points of intersection is

- a)  $\pi/4$     b)  $\tan^{-1}(4\sqrt{2}/7)$   
 c)  $\tan^{-1}(4\sqrt{7})$     d)  $\tan^{-1}(2\sqrt{2}/7)$

Key. B

Sol. The point of intersection is  $x^2 = 2, y = 1$ . The given equations represent four parabolas.

$$y = \pm(x^2 - 1) \text{ and } y = \pm(x^2 - 3)$$

The curves intersect when  $1 < x^2 < 3$  or  $1 < x < \sqrt{3}$  or  $-\sqrt{3} < x < -1$

$$\therefore y = x^2 - 1 \text{ and } y = -(x^2 - 3)$$

The points of intersection are  $(\pm\sqrt{2}, 1)$

$$\text{At } (\sqrt{2}, 1), m_1 = 2x = 2\sqrt{2}, m_2 = -2x = -2\sqrt{2}$$


$$\therefore \tan \theta = \left| \frac{4\sqrt{2}}{1-8} \right| = \frac{4\sqrt{2}}{7} \Rightarrow \theta = \tan^{-1} \left( \frac{4\sqrt{2}}{7} \right)$$

60. A man of height 2m walks directly away from a lamp at a height of 5m, on a level road at 3m/s. The rate at which the length of his shadow is increasing is

- a) 1m/s                      b) 2m/s                      c) 3m/s                      d) 4m/s

Key. B

Sol. Let L be the lamp and PQ be the man and OQ = x metre be his shadow and let MQ = y metre.



$$\therefore \frac{dy}{dt} = \text{speed of the man} \\ = 3 \text{ m/s (given)}$$

$\therefore \Delta OPQ$  and  $\Delta OLM$  are similar

$$\therefore \frac{OM}{OQ} = \frac{LM}{PQ}$$

$$\Rightarrow \frac{x+y}{x} = \frac{5}{2}$$

$$\Rightarrow y = \frac{3}{2}x$$

$$\therefore \frac{dy}{dt} = \frac{3}{2} \frac{dx}{dt}$$

$$\Rightarrow 3 = \frac{3}{2} \frac{dx}{dt}$$

$$\Rightarrow \frac{dx}{dt} = 2 \text{ m/s.}$$

61. The parametric equations of a curve are  $x = t^2$  and  $y = t^3$ .  $A(t_1), B(t_2)$  are points on the curve. If  $t_1 = 1, t_2 = 3$  then the abscissa of the point P on the curve the tangent at which is parallel to chord AB is  
 A) 13 / 6                      B) 169 / 36                      C) 17 / 36                      D) 27 / 4

Key. B

Sol.  $(t^2, t^3)$  is a/pt on the curve.

$$\frac{dx}{dt} = 2t \text{ and } \frac{dy}{dt} = 3t^2$$

$$\frac{dy}{dx} = \frac{3t^2}{2t} = \frac{3}{2}t$$

A=(1,1) and B=(9,27)

$$\text{Slope of AB} = \frac{27-1}{9-1} = \frac{26}{8} = \frac{13}{4}$$

$$\frac{3}{2}t = \frac{13}{4} \Rightarrow t = \frac{13}{6}$$

$\therefore Tgt$  at  $\left(\left(\frac{13}{6}\right)^2, \left(\frac{13}{6}\right)^3\right)$  is parallel to AB

62. The sum of the coordinates of the point on the graph of  $f(x) = x^3 + 4x$  the tangent at which is parallel to the chord joining the points (-2, -16) and (1, 5) is  
 A) -6                      B) 4                      C) -8                      D) 5/2

Key. A

$$\text{Sol. Slope of chord} = \frac{5 - (-16)}{1 - (-2)} = \frac{21}{3} = 7$$

$$f'(x) = 3x^2 + 4$$

By L.M.V.T  $\exists C \in (-2, 1)$  such that  $f'(C) = 7$

$$3c^2 + 4 = 7 \Rightarrow C = \pm 1$$

$$\therefore C = -1$$

Point  $= (C, f(C)) = (-1, -5)$

63. The maximum value of the sum of the intercepts made by any tangent to the curve  $(a \sin^2 \theta, 2a \sin \theta)$  with the axes is

- (a)  $2a$                       (b)  $a/4$                       (c)  $a/2$                       (d)  $a$

Key. A

Sol. Equation of tangent  $\frac{y - 2a \sin \theta}{x - a \sin^2 \theta} = \frac{1}{\sin \theta}$

$$\Rightarrow \frac{x}{-a \sin^2 \theta} + \frac{y}{a \sin \theta} = 1$$

Sum of intercepts  $= a(\sin^2 \theta + \sin \theta)$

which is maximum when  $\sin \theta = 1$

(sum of intercepts) $_{\max} = 2a$

64. The tangent to the curve  $y = e^x$  drawn at the point  $(c, e^c)$  intersects the line joining the points  $(c-1, e^{c-1})$  and  $(c+1, e^{c+1})$


- (a) on the left of  $x = c$                       (b) on the right of  $x = c$   
 (c) at no point                      (d) at all points

Key. A

Sol. Slope of AB  $= \frac{e^{c+1} - e^{c-1}}{2}$

Slope of tangent is  $e^c$

$$\frac{e^{c+1} - e^{c-1}}{2} > e^c \left( \because e - \frac{1}{e} > 2 \right)$$



y-coordinate of straight line AB at  $x = c$  will be more than y-coordinate of the tangent at  $x = c$  for this graph.

Also rate of increasing of AB is more than tangent. So already these two lines had interested before  $x = c$ .